


 Cite this: *RSC Adv.*, 2021, **11**, 29130

# A review on convenient synthesis of substituted coumarins using reuseable solid acid catalysts

 Susheel Gulati, \* Rajvir Singh and Suman Sangwan

Due to growing concern about chemicals and their impact on the environment, cleaner reaction conditions are needed to be incorporated into chemical synthetic procedures. Recently, the use of heteropolyacid catalysts, mainly reuseable solid acid catalysts, has gained a leading role in organic synthesis due to their environmental and economic considerations and industrial utilization. The high catalytic activity, moisture sensitivity, reusability and inexpensive makes solid supported reagents attractive substituents to conventional Lewis acids. Nowadays synthesis of coumarins and their derivatives has attracted considerable attention from organic and medicinal chemists for many years as a large number of natural products contain this heterocyclic nucleus. In continuation with our investigations into the synthesis of substituted coumarins and due to several advantages of heterogeneous catalysts *viz.* cost-effective, no side products, high yield of desired products and no toxic waste material, here we report a new approach for the synthesis of substituted coumarins using solid acid catalysts.

 Received 23rd June 2021  
 Accepted 19th August 2021

DOI: 10.1039/d1ra04887b

[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

## Introduction

Coumarins as a major class of natural and synthetic products exhibit a variety of pharmacological and biological activities.<sup>1–3</sup> There is growing curiosity for coumarins and their derivatives due to their anti-HIV, anti-oxidant, anti-fungal, antihelmintic and antibacterial properties.<sup>4–8</sup> They are used in food and cosmetic industries as additive and also found applications as insecticides, optical brighteners, fluorescent and laser dyes.<sup>9–13</sup> Masesane *et al.*<sup>14</sup> reported the synthesis of chromane derivatives through the reaction of salicylaldehyde and enolates and they found that reactions of salicylaldehyde and enolates give nearly optically pure chromane derivatives. Coumarins can also be prepared by various methods *viz.* Pechmann condensation, Perkin, Knoevenagel and Reformatsky reactions.<sup>15</sup> Pechmann condensation has been most popularly method for coumarin synthesis, since it proceeds from simple substrate *viz.* phenol and  $\beta$ -ketoester and gives excellent yields of coumarins. Pechmann condensation utilizes various catalysts *viz.* sulphuric acid, trifluoroacetic acid, phosphorous pentoxide,  $ZrCl_4$ ,  $TiCl_4$  and ionic liquids, which have many drawbacks such as long reaction time, use of hazardous solvents, creates side products and salt waste due to acid neutralization.<sup>16</sup> There has been some effort to find alternative, eco-friendly synthetic methods. Nowadays, the use of heterogeneous solid acid catalysts has fascinated significant attention. These catalysts have some advantages such as ease of product work-up, recyclability, strong safety and tolerance for wide range of temperature and

pressures.<sup>17–21</sup> Naikwadi and his coworkers<sup>22</sup> reported the catalytic reaction of active methylene compounds with cyclic enol ethers and aryl acetals through oxonium intermediate under solvent-free conditions using heterogeneous solid acid catalysts and they found that Amberlyst-15 gave excellent yields of alkylated products. Therefore, there is a propensity to replace the classic homogeneous catalysts by heterogeneous solid acid catalysts. Due to several benefits of heterogeneous catalysts, in this review we encapsulate synthesis of substituted coumarins using solid acid catalyst.

## Synthesis of substituted coumarins using solid acid catalysts

An efficient and facile synthesis of novel class of coumarin-containing secondary benzamide derivatives (**4**) has been developed *via* one-pot condensation of 5,7-dihydroxy coumarins (**1**), substituted aldehydes (**2**) and benzamide (**3**) using tungstate sulphuric acid by Karami and his coworkers (Scheme 1).<sup>23</sup> To standardize the reaction conditions, a reaction between 5,7-dihydroxy-4-methylcoumarin, benzaldehyde and benzamide were chosen as a model reaction. The model reaction was screened under various conditions. After conducting several experiments, they found that the desired reaction took place efficiently using 1 mol% of tungstate sulphuric acid (TSA) at 120 °C under solvent-free conditions. The proposed mechanism of the formation of desired products is shown in Fig. 1. According to proposed mechanism, first there is formation of adduct (**I**) by the condensation reaction of substituted aldehyde and benzamide in the presence of TSA as an efficient proton source. Then C-8 of coumarin attacks on adduct (**I**) and gives

Department of Chemistry, Chaudhary Charan Singh Haryana Agricultural University, Hisar, 125004, India. E-mail: [sgbhuna108@gmail.com](mailto:sgbhuna108@gmail.com)/[sgbhuna@hau.ac.in](mailto:sgbhuna@hau.ac.in)





Scheme 1 TSA-catalyzed synthesis of coumarin-containing secondary benzamides.

intermediate (II). Finally by tautomerization desired product obtained. They also found that tungstate sulphuric acid is reuseable heterogeneous catalyst, which make this procedure mild, convenient and eco-friendly. Simplicity of procedure, use of safe and recyclable catalysts, high yields and short reaction times are some beauties of present methodology.

Khaligh *et al.* found that poly(4-vinylpyridinium) hydrogen sulfate solid acid was efficient catalyst for the synthesis of

substituted coumarins (7) *via* Pechmann condensation reaction between substituted phenols (5) and  $\beta$ -ketoester (6) using ultrasound irradiation at ambient temperature. Simplicity in operation, avoid use of toxic catalysts and solvents, excellent yield of desired products, reuse of catalyst are some merits of present methodology. First they standardized the reaction conditions by exploring model reaction between resorcinol and ethylacetoacetate (Scheme 2)<sup>24</sup> in presence of different solvents



Fig. 1 Possible mechanism for synthesis of coumarin-containing secondary benzamides.








Fig. 2 Proposed mechanism for Pechmann reaction of resorcinol with ethyl acetoacetate at room temperature under ultrasonic irradiation.



Scheme 3 Synthesis of bis-coumarin derivatives in Fe<sub>3</sub>O<sub>4</sub>@sulfosalicylic acid MNPs as catalyst under microwave irradiation in water.

with an  $\alpha,\beta$ -unsaturated intermediate to give the final polyhydroquinoline product. Finally, a tautomeric proton shift produces the desired product. Table 4 presented the results

from the synthesis of bis-coumarin by reaction of benzaldehyde and 4 hydroxycoumarin in the presence of Fe<sub>3</sub>O<sub>4</sub>@sulfosalicylic acid magnetic nanoparticles which has

Table 3 Optimization of the model reaction

| Entry | Catalyst (g)  | Power | Time (min) | Yield (%) |
|-------|---|-------|------------|-----------|
| 1     | Sulfosalicylic acid (0.01)                                  | 180   | 15         | 75        |
| 2     | FeCl <sub>3</sub> ·6H <sub>2</sub> O (0.05)                 | 180   | 15         | 43        |
| 3     | Bulk-Fe <sub>3</sub> O <sub>4</sub> (0.05)                  | 180   | 15         | 50        |
| 4     | Nano-Fe <sub>3</sub> O <sub>4</sub> (0.05)                  | 180   | 15         | 68        |
| 5     | Fe <sub>3</sub> O <sub>4</sub> @sulfosalicylic acid (0.03)  | 180   | 20         | 89        |
| 6     | Fe <sub>3</sub> O <sub>4</sub> @sulfosalicylic acid (0.03)  | 300   | 10         | 92        |
| 7     | Fe <sub>3</sub> O <sub>4</sub> @sulfosalicylic acid (0.05)  | 100   | 10         | 89        |
| 8     | Fe <sub>3</sub> O <sub>4</sub> @sulfosalicylic acid (0.05)  | 180   | 10         | 96        |
| 9     | Fe <sub>3</sub> O <sub>4</sub> @sulfosalicylic acid (0.08)  | 180   | 10         | 96        |
| 10    | Fe <sub>3</sub> O <sub>4</sub> @sulfosalicylic acid (0.015) | 180   | 10         | 80        |
| 11    | —   | 180   | 20         | 30        |





Fig. 3 Proposed mechanism for the synthesis of biscoumarin derivatives in the presence of  $\text{Fe}_3\text{O}_4$ @sulfosalicylic acid magnetic nanoparticles.

Table 4 Comparison of efficiency of present catalyst with other catalysts reported in literature

| Entry | Catalyst/condition   | Time (min) | Yield | References |
|-------|--|------------|-------|------------|
| 1     | Ionic liquids, reflux  | 260        | 84    | 26         |
| 2     | Choline hydroxide, reflux  | 240        | 86    | 27         |
| 3     | No catalyst/trifluoroethanol, reflux                                       | 360        | 80    | 28         |
| 4     | $\text{Fe}_3\text{O}_4$ @sulfosalicylic acid/ $\text{H}_2\text{O}$ , $M_w$ | 10         | 96    | 25         |

been compared with the other methods reported in literature. The results show that the present method is preferable because of its reaction times and efficiency.

Samiei *et al.* reported the green synthesis coumarin derivatives (7) *via* Pechmann condensation reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in excellent yield under solvent-free conditions in presence of novel sulfonated carbon-coated magnetic nanoparticles (Scheme 4).<sup>29</sup>

For optimization of reaction conditions, first the model reaction was explored between resorcinol and ethyl acetoacetate to produce 7-hydroxy-4-methylcoumarin. The reaction was also optimized with respect to various parameters *viz.* catalyst loading, different temperatures and various solvents as shown in Table 5. It was observed from Table 5 that lack of catalyst and also with a catalyst loading of  $\text{Fe}_3\text{O}_4$  NPs, CCMNPs ( $\text{Fe}_3\text{O}_4$ @C) led to no product even after 6 h, while the use of SCCMNPs ( $\text{Fe}_3\text{O}_4$ @C@OSO<sub>3</sub>H) could produce related 4*H*-coumarin in



Scheme 4 Synthesis of substituted coumarins.



**Table 5** The effect of various solvents, temperature and catalyst loadings for the synthesis of substituted coumarins through Pechmann condensation

| Entry | Catalyst loading   | Solvent                         | T (°C)     | Time (min) | Yield (%)   |
|-------|--|---------------------------------|------------|------------|-------------|
| 1     | —  | —                               | 120        | 360        | No reaction |
| 2     | Fe <sub>3</sub> O <sub>4</sub> NPs (6.5 mol%)                              | —                               | 120        | 360        | No reaction |
| 3     | CCMNPs (Fe <sub>3</sub> O <sub>4</sub> @C) (6.5 mol%)                      | —                               | 120        | 360        | No reaction |
| 4     | SCCMNPs (Fe <sub>3</sub> O <sub>4</sub> @C@OSO <sub>3</sub> H) (3.25 mol%) | —                               | 120        | 30         | 86          |
| 5     | <b>SCCMNPs (Fe<sub>3</sub>O<sub>4</sub>@C@OSO<sub>3</sub>H) (6.5 mol%)</b> | —                               | <b>120</b> | <b>20</b>  | <b>98</b>   |
| 6     | SCCMNPs (Fe <sub>3</sub> O <sub>4</sub> @C@OSO <sub>3</sub> H) (13 mol%)   | —                               | 120        | 20         | 98          |
| 7     | SCCMNPs (Fe <sub>3</sub> O <sub>4</sub> @C@OSO <sub>3</sub> H) (6.5 mol%)  | —                               | 100        | 30         | 87          |
| 8     | SCCMNPs (Fe <sub>3</sub> O <sub>4</sub> @C@OSO <sub>3</sub> H) (6.5 mol%)  | —                               | 90         | 40         | 83          |
| 9     | SCCMNPs (Fe <sub>3</sub> O <sub>4</sub> @C@OSO <sub>3</sub> H) (6.5 mol%)  | H <sub>2</sub> O                | Reflux     | 360        | Trace       |
| 10    | SCCMNPs (Fe <sub>3</sub> O <sub>4</sub> @C@OSO <sub>3</sub> H) (6.5 mol%)  | Toluene                         | Reflux     | 360        | No reaction |
| 11    | SCCMNPs (Fe <sub>3</sub> O <sub>4</sub> @C@OSO <sub>3</sub> H) (6.5 mol%)  | CH <sub>2</sub> Cl <sub>2</sub> | Reflux     | 360        | No reaction |
| 12    | SCCMNPs (Fe <sub>3</sub> O <sub>4</sub> @C@OSO <sub>3</sub> H) (6.5 mol%)  | EtOH                            | Reflux     | 360        | No reaction |
| 13    | SCCMNPs (Fe <sub>3</sub> O <sub>4</sub> @C@OSO <sub>3</sub> H) (6.5 mol%)  | CH <sub>3</sub> CN              | Reflux     | 360        | No reaction |

**Table 6** Comparison of activity of some heterogeneous solid acid catalysts for the synthesis of substituted coumarins

| Entry | Catalyst  | Catalyst (mol%) | T (°C)     | Time (min) | Yield (%) | References |
|-------|---|-----------------|------------|------------|-----------|------------|
| 1     | Fe <sub>3</sub> O <sub>4</sub> -DABCO                                 | 1               | 100        | 40         | 93        | 30         |
| 2     | γ-Fe <sub>3</sub> O <sub>4</sub> @HAP-Ag                              | 10              | 80         | 20         | 95        | 31         |
| 3     | Fe <sub>3</sub> O <sub>4</sub> @SiO <sub>2</sub> @PrSO <sub>3</sub> H | 2               | 130        | 25         | 96        | 32         |
| 4     | CMK-15-SO <sub>3</sub> H  | 3               | 130        | 20         | 95        | 33         |
| 5     | Random pore carbon-SO <sub>3</sub> H                                  | 7               | 130        | 60         | 90        | 34         |
| 6     | Fe <sub>3</sub> O <sub>4</sub> @SiO <sub>2</sub> @EtSO <sub>3</sub> H | 75              | 90         | 90         | 93        | 35         |
| 7     | SnCl <sub>4</sub> -SiO <sub>2</sub>                                   | 5               | 120        | 35         | 90        | 36         |
| 8     | SBA-15-Ph-Pr-SO <sub>3</sub> H  | 7               | 130        | 60         | 90        | 36         |
| 9     | ZrW <sub>2</sub>  | 20              | 12         | 120        | 94        | 37         |
| 10    | SnW <sub>2</sub>  | 20              | 120        | 120        | 88        | 37         |
| 11    | Nanosponge MFI zeolite  | 0.5             | 130        | 120        | 94        | 38         |
| 12    | TiZnO   | 10              | 110        | 180        | 85        | 39         |
| 13    | Fe <sub>3</sub> O <sub>4</sub> @Boehmite-NH <sub>2</sub> -CoII        | 6.6             | 90         | 30         | 95        | 40         |
| 14    | <b>SCCMNPs</b>  | <b>6.5</b>      | <b>120</b> | <b>20</b>  | <b>98</b> | 29         |

a good yield during the short time. Hence, SCCMNP with the sulfonic acid moiety on the surface of MNP was introduced as an effective catalyst in the Pechmann condensation. They also found that 6.5 mol% catalyst loading was identified as an optimized concentration in the model reaction at 120 °C under solvent-free condition.

The comparison of catalytic activity of present catalyst with other catalysts reported in literature was shown in Table 6.

Khan and his coworkers reported the synthesis of coumarins (7) *via* Pechmann condensation reaction between substituted phenols (5) and β-ketoesters (6) in presence of zirconia-based heterogeneous catalyst (Scheme 5).<sup>41</sup> First of all model

reaction was carried out between resorcinol and ethyl acetoacetate without a catalyst at 80 °C, but there will be no formation of product as shown in Table 7. They also observed that excellent yield of product was obtained when electron releasing group linked with substituted phenols, while poor yield of product was obtained when electron withdrawing group linked with substituted phenols. They also studied reaction between resorcinol and ethyl acetoacetate with 50 mg of the catalyst ZrO<sub>2</sub>-TiO<sub>2</sub> in polar solvent *viz.* ethanol and non-polar solvent *viz.* toluene by varying the temperature condition as shown in Table 8. The plausible mechanism for the reaction is depicted in Fig. 4.


**Scheme 5** Synthesis of substituted coumarins.

Table 7 The reaction for synthesis of substituted coumarins in solvent-free condition at room temperature

| Entry | Reactant                                  | Catalyst                           | Temperature (°C) | Time (min) | %Yield     |
|-------|---|------------------------------------|------------------|------------|------------|
| 1     | Resorcinol + ethylacetoacetate            | ZrO <sub>2</sub> -TiO <sub>2</sub> | RT               | 180        | 97         |
| 2     | Resorcinol + ethylacetoacetate            | ZrO <sub>2</sub> -ZnO              | RT               | 240        | 63         |
| 3     | Resorcinol + ethylacetoacetate            | ZrO <sub>2</sub> /cellulose        | RT               | 180        | Nil        |
| 4     | Catechol + ethylacetoacetate              | ZrO <sub>2</sub> -TiO <sub>2</sub> | 80               | 240        | 55         |
| 5     | <i>o</i> -Nitrophenol + ethylacetoacetate | ZrO <sub>2</sub> -TiO <sub>2</sub> | 80               | 240        | Nil        |
| 6     | <b>Resorcinol + ethylacetoacetate</b>     | <b>Without catalyst</b>            | <b>80</b>        | <b>240</b> | <b>Nil</b> |

Table 8 Comparison of efficiency ZrO<sub>2</sub>-TiO<sub>2</sub> with reported catalysts

| Entry | Catalyst                               | Time (min) | Temperature (°C) | Solvent             | Yield     | References |
|-------|--|------------|------------------|---------------------|-----------|------------|
| 1     | Zeolite BEA                            | 240        | 130              | PhNO <sub>2</sub>   | 63        | 42         |
| 2     | PFPAT                                  | 180        | 110              | Toluene             | 90        | 43         |
| 3     | MFRH                                   | 50         | 80               | Solvent-free        | 65        | 43         |
| 4     | Nanoreactors                           | 60         | 130              | Solvent-free        | 30        | 43         |
| 5     | CMK-5-SO <sub>3</sub> H                | 20         | 130              | Solvent-free        | 95        | 44         |
| 6     | CMK-5                                  | 60         | 130              | Solvent-free        | 10        | 44         |
| 7     | <b>ZrO<sub>2</sub>-TiO<sub>2</sub></b> | <b>180</b> | <b>RT</b>        | <b>Solvent-free</b> | <b>97</b> | <b>41</b>  |
| 8     | <b>ZrO<sub>2</sub>-TiO<sub>2</sub></b> | <b>110</b> | <b>60</b>        | <b>Toluene</b>      | <b>95</b> | <b>41</b>  |
| 9     | <b>ZrO<sub>2</sub>-TiO<sub>2</sub></b> | <b>150</b> | <b>60</b>        | <b>Ethanol</b>      | <b>92</b> | <b>41</b>  |

Kumbar and his coworkers developed efficient and facile methodology for synthesis of class of chromeno-3-substituted derivatives (**10a–10l**) in excellent yields in presence of solid-supported heterogeneous silica sulphuric acid as a reusable catalyst (Scheme 6).<sup>45</sup>

They found that use of silica sulphuric acid as catalyst provide good to excellent yields of desired products as shown in Table 9. The reaction was also optimized with respect to polar protic and aprotic solvents *viz.* acetonitrile, ethanol, DMF, dioxane, THF and DMSO as summarized in Table 10. The

plausible mechanism of reaction was presented in Fig. 5. First there is nucleophilic attack of aniline on the carbonyl carbon of coumarin. Then in next step protonation occurs from silica sulphuric acid, forming itself as a nucleophile in the reaction mixture. Then nucleophilic SSA abstracts protons from nitrogen and gains stability by the formation of double bond between C and N and subsequent dehydration give desired product.

Moghaddam and Hoda designed magnetic graphene oxide coated with cysteic acid as an efficient and reusable catalyst for the synthesis of 4*H*-chromene derivatives (**13**) *via* one-pot





Fig. 4 Plausible mechanism for the synthesis of substituted coumarins in presence of zirconia-based heterogeneous catalyst.





Scheme 6 Synthesis of novel coumarin Schiff bases in presence of silica sulphuric acid as reuseable catalyst.

multicomponent reaction between enolizable compound (11), malononitrile (12), substituted aldehydes (2) or isatin and a mixture of water-ethanol as a green solvent (Scheme 7).<sup>46</sup> Excellent yield of desired products, less reaction time, mild reaction conditions and eco-friendly approach are some merits of present methodology.

An efficient and facile method for the one-pot synthesis of dihydropyrano[3,2-*c*]chromene derivatives (14) have been reported *via* reaction between substituted aldehydes (2), 4-hydroxycoumarin (8) and malononitrile (12) in presence of poly(4-vinyl-pyridine) as a cheap, efficient and recyclable catalyst (Scheme 8). They also reported the synthesis of biscoumarin derivatives (9) *via* one-pot reaction between substituted aldehydes (2) and 4-hydroxycoumarin (8) in presence of same catalyst (Scheme 9).<sup>47</sup>

Table 9 Physical and analytical data of synthesized coumarin derivatives

| Products | R                          | Yield (%) | Time (min) | Melting point (°C) |
|----------|----------------------------|-----------|------------|--------------------|
| 10a      | H                          | 78        | 180        | 165–167            |
| 10b      | <i>p</i> -Cl               | 62        | 210        | 193–195            |
| 10c      | <i>p</i> -Br               | 61        | 190        | 182–184            |
| 10d      | <i>p</i> -OH               | 67        | 195        | 198–200            |
| 10e      | <i>p</i> -OCH <sub>3</sub> | 62        | 210        | 205–208            |
| 10f      | <i>p</i> -CH <sub>3</sub>  | 71        | 240        | 202–204            |
| 10g      | 2,6-Dimethyl               | 58        | 220        | 188–190            |
| 10h      | <i>m</i> -Cl               | 68        | 210        | 197–200            |
| 10i      | <i>m</i> -Br               | 69        | 190        | 178–181            |
| 10j      | <i>m</i> -OH               | 62        | 195        | 184–186            |
| 10k      | <i>m</i> -OCH <sub>3</sub> | 59        | 200        | 208–210            |
| 10l      | <i>m</i> -CH <sub>3</sub>  | 73        | 225        | 212–214            |

To optimize the reaction conditions, a model reaction was explored between 4-chlorobenzaldehyde, malononitrile and 4-hydroxycoumarin in presence of different concentration of P<sub>4</sub>VPy. The effect of different solvents *viz.* CH<sub>3</sub>CN, CH<sub>2</sub>Cl<sub>2</sub>, H<sub>2</sub>O and EtOH and temperature in the synthesis of dihydropyrano[3,2-*c*] chromene derivatives in the presence of P<sub>4</sub>VPy summarized in Table 11. They found that best result was obtained using 20 mg of P<sub>4</sub>VPy at 70 °C in a mixture of H<sub>2</sub>O and ethanol. They also observed that aldehydes containing electron-withdrawing as well as electron-donating groups such as Cl, Br, CH<sub>3</sub>, OCH<sub>3</sub>, NO<sub>2</sub> and OH in the *ortho*, *meta* and *para* positions can be easily converted to the corresponding dihydropyrano[3,2-*c*] chromenes in less reaction times with excellent yield.

After most favourable results of P<sub>4</sub>VPy in the synthesis of dihydropyrano[3,2-*c*]chromene derivatives, they were interested to study the efficiency of this polymeric reagent in the synthesis

Table 10 Optimization of reaction conditions

| Entry | Solvent      | SSA    | Time (h) | Temperature (°C) | Yield (%) |
|-------|--------------|--------|----------|------------------|-----------|
| 1     | Acetonitrile | 1.0    | 4        | 25               | 35        |
| 2     | Ethanol      | 1.0    | 3        | 25               | 78        |
| 3     | DMF          | 1.0    | 12       | 25               | Nil       |
| 4     | Dioxane      | 1.0    | 6        | 25               | 38        |
| 5     | THF          | 1.0    | 12       | 25               | Trace     |
| 6     | DMSO         | 1.0    | 12       | 25               | Nil       |
| 7     | Acetone      | 1.0    | 12       | 25               | Nil       |
| 8     | Acetonitrile | 2.0    | 12       | 40               | 42        |
| 9     | Ethanol      | 0.0    | 12       | 25               | Nil       |
| 10    | Ethanol      | Silica | 12       | 25               | 45        |







Fig. 5 Proposed reaction pathway for the synthesis of substituted coumarins.



Scheme 7 One-pot three-component reaction of enolizable compound, active methylene nitriles, and aldehydes catalyzed by MNPs·GO-CysA in water : ethanol.



Scheme 8 Synthesis of dihydropyrano[3,2-c] chromene derivatives.





Scheme 9 Synthesis of biscoumarin derivatives.

of biscoumarins. For standardization of reaction conditions, first model reaction was carried out between 4-chlorobenzaldehyde and 4-hydroxycoumarin in the presence of  $P_4Vpy$  at different reaction conditions as shown in Table 12. They observed that best reaction conditions for the synthesis of the biscoumarin derivatives are use of 20 mg of the  $P_4Vpy$  in water at 90 °C. They also found that aldehydes containing electron-withdrawing or electron donating substituents converting to desired products in less time. The plausible mechanism for the synthesis of substituted pyrazoles given in Fig. 6. The comparison of catalytic activity and reaction conditions of present catalyst  $P_4Vpy$  for the synthesis of dihydropyrano[3,2-*c*] chromene derivatives and biscoumarin derivatives are summarized in Table 13 and Table 14. This comparison shows disadvantages of the other procedures such as long reaction times, toxic reagents, high temperature, organic solvents, excess reagents and low yields.

An efficient, green and inexpensive synthesis of benzylpyrazolyl coumarin (16) by one-pot multicomponent condensation

of hydrazine hydrate or phenyl hydrazine (15),  $\beta$ -ketoester (6), substituted aldehydes (2) and 4-hydroxycoumarin (8) in the presence of Amberlite IR-120 as a catalyst in an aqueous medium has been reported by Katariya and his coworkers (Scheme 10).<sup>58</sup>

Kaur *et al.* reported the synthesis of 3,3'-(arylmethylene) bis(4-hydroxy-2H-chromen-2-ones) *via* one-pot reaction between substituted aldehydes (2) and 4-hydroxy coumarin (8) catalyzed by camphor sulfonic acid (Scheme 11).<sup>59</sup> Mild reaction conditions, use of metal-free organocatalyst, excellent yields of desired products, high atom economy, eco-friendly, easy isolation of products and no need of column chromatography are some merits of present methodology. To standardize the reaction conditions they conducted a model reaction between 4-methylbenzaldehyde and 4-hydroxycoumarin. Firstly, they explored the reaction in the absence of catalyst as well as solvent at room temperature and they observed that trace amount of yield was obtained after 24 h. Then under catalyst-free conditions, the same reaction was give 22% yield of desired product in ethanol. After getting the poor yields of desired product, they were interested to check the catalytic activity of camphor

Table 11 The effect of different reaction conditions for the synthesis of dihydropyrano[3,2-*c*] chromene derivatives in the presence of  $P_4Vpy$ 

| Entry | Catalyst (mg) | Solvent                         | Temperature (°C) | Time (min) | Yield (%)           |
|-------|---------------|---------------------------------|------------------|------------|---------------------|
| 1     | —             | No solvent                      | RT               | 120        | Nil                 |
| 2     | —             | No solvent                      | 100              | 120        | Nil                 |
| 3     | 20            | CH <sub>3</sub> CN              | RT               | 120        | Nil                 |
| 4     | 20            | CH <sub>3</sub> CN              | Reflux           | 120        | Mixture of products |
| 5     | 20            | CH <sub>2</sub> Cl <sub>2</sub> | RT               | 120        | Nil                 |
| 6     | 20            | CH <sub>2</sub> Cl <sub>2</sub> | Reflux           | 120        | Mixture of products |
| 7     | 20            | H <sub>2</sub> O                | 90               | 180        | 50                  |
| 8     | 24            | H <sub>2</sub> O                | 90               | 120        | 50                  |
| 9     | 24            | EtOH                            | RT               | 150        | 60                  |
| 10    | 20            | EtOH                            | 50               | 120        | 60                  |
| 11    | 20            | EtOH                            | 70               | 120        | 60                  |
| 12    | 20            | H <sub>2</sub> O/<br>EtOH       | 70               | 5          | 95                  |
| 13    | 24            | H <sub>2</sub> O/<br>EtOH       | 70               | 5          | 95                  |

Table 12 Optimization of the reaction conditions for the synthesis of biscoumarin derivatives catalyzed by  $P_4Vpy$ 

| Entry | Catalyst (mg) | Solvent                         | Temperature (°C) | Time (min) | Yield (%)           |
|-------|---------------|---------------------------------|------------------|------------|---------------------|
| 1     | —             | No solvent                      | RT               | 120        | Nil                 |
| 2     | —             | No solvent                      | 100              | 120        | Nil                 |
| 3     | 20            | CH <sub>3</sub> CN              | RT               | 120        | Nil                 |
| 4     | 20            | CH <sub>3</sub> CN              | Reflux           | 120        | Mixture of products |
| 5     | 20            | CH <sub>2</sub> Cl <sub>2</sub> | RT               | 120        | Nil                 |
| 6     | 20            | CH <sub>2</sub> Cl <sub>2</sub> | Reflux           | 120        | Mixture of products |
| 7     | 20            | EtOH                            | RT               | 120        | 30                  |
| 8     | 20            | EtOH                            | Reflux           | 120        | 60                  |
| 9     | 10            | H <sub>2</sub> O                | RT               | 90         | 40                  |
| 10    | 15            | H <sub>2</sub> O                | RT               | 90         | 50                  |
| 11    | 20            | H <sub>2</sub> O                | RT               | 90         | 75                  |
| 12    | 20            | H <sub>2</sub> O                | 90               | 5          | 96                  |
| 13    | 24            | H <sub>2</sub> O/<br>EtOH       | 90               | 5          | 96                  |






Table 13 Comparison of different catalysts for the synthesis of dihydropyrano[3,2-c]chromene derivatives


| Entry | Catalyst (mol%)                     | Reaction conditions             | Time (min) | Yield (%) | References |
|-------|-------------------------------------|---------------------------------|------------|-----------|------------|
| 1     | SDS                                 | Water/60 °C                     | 150        | 88        | 48         |
| 2     | Nano ZnO                            | Ethanol reflux                  | 90         | 49        | 49         |
| 3     | Nano Al(OH) <sub>3</sub>            | Ethanol reflux                  | 120        | 48        | 49         |
| 4     | DAHP                                | Ethanol-H <sub>2</sub> O/25 °C  | 240        | 85        | 50         |
| 5     | (S)-proline                         | Ethanol-H <sub>2</sub> O/100 °C | 180        | 78        | 50         |
| 6     | Nano Al <sub>2</sub> O <sub>3</sub> | Ethanol reflux                  | 120        | 71        | 51         |
| 7     | P <sub>4</sub> VPy                  | Ethanol-H <sub>2</sub> O/70 °C  | 5          | 95        | 47         |

Table 14 Comparison of different catalysts used for the synthesis of biscoumarins

| Entry | Catalyst (mol%)  | Reaction conditions     | Time (min) | Yield (%) | References |
|-------|--|-------------------------|------------|-----------|------------|
| 1     | SDS  | Water/60 °C             | 150        | 93        | 48         |
| 2     | [bmim]BF <sub>4</sub>  | Solvent-free/60–70 °C   | 150        | 91        | 52         |
| 3     | I <sub>2</sub>   | H <sub>2</sub> O/100 °C | 27         | 93        | 53         |
| 4     | CHOH   | Solvent-free/50 °C      | 120        | 99        | 54         |
| 5     | [P <sub>4</sub> VPy-BuSO <sub>3</sub> H]Cl-X(AlCl <sub>3</sub> ) | Toluene/90 °C           | 36         | 93        | 55         |
| 6     | PSA  | Solvent-free/100 °C     | 240        | 96        | 56         |
| 7     | Piperidine   | EtOH/r.t                | 240        | 96        | 57         |
| 8     | P <sub>4</sub> VPy   | H <sub>2</sub> O/90 °C  | 5          | 96        | 47         |



Scheme 10 General scheme for the synthesis of benzylpyrazolyl coumarin.



Scheme 11 Reported protocols for the synthesis of biscoumarin.

the Knoevenagel intermediate. Then second molecule of 4-hydroxycoumarin attack on Knoevenagel intermediate followed by enolisation gives the desired product in excellent yield.

A novel heterogeneous catalytic method was developed for the synthesis of coumarin (7) *via* reaction between  $\beta$ -ketoesters (6) and substituted phenols (5) in presence of Zn<sub>0.925</sub>Ti<sub>0.075</sub>O as catalyst by Jadhav and his coworkers (Scheme 12).<sup>60</sup> They also observed that this shows recycle activity up to seven cycles with very good stability. Firstly, they standardized the reaction

conditions in order to verify the role of catalyst by conducting a model reaction between phloroglucinol and ethylacetoacetate under solvent-free conditions and the results are summarized in Table 16. They observed that Zn<sub>0.925</sub>Ti<sub>0.075</sub>O is best catalyst for optimization studies in the synthesis of coumarin by Pechmann condensation. The various solvents effect *viz.* DCM, ethylacetate, acetonitrile, water, ethanol, toluene and DMF also studied for optimizing the reaction conditions during the synthesis of coumarin and the results are summarized in Table




Table 15 Standardization of reaction conditions for the synthesis of 3,3'-(arylmethylene)bis(4-hydroxy-2H-chromen-2-ones)

| Entry | Catalyst (mol%)                        | Solvent                                  | Time (h) | Yield (%) |
|-------|--|--|----------|-----------|
| 1     | No catalyst                            | Solvent-free                             | 24       | Nil       |
| 2     | No catalyst                            | EtOH                                     | 6        | 22        |
| 3     | Camphor sulfonic acid (20 mol%)        | EtOH                                     | 6        | 78        |
| 4     | Camphor sulfonic acid (20 mol%)        | MeOH                                     | 6        | 72        |
| 5     | Camphor sulfonic acid (20 mol%)        | H <sub>2</sub> O                         | 6        | 61        |
| 6     | <b>Camphor sulfonic acid (20 mol%)</b> | <b>EtOH : H<sub>2</sub>O (1 : 1 v/v)</b> | <b>2</b> | <b>94</b> |
| 7     | Camphor sulfonic acid (15 mol%)        | EtOH : H <sub>2</sub> O (1 : 1 v/v)      | 2        | 86        |
| 8     | Camphor sulfonic acid (20 mol%)        | EtOH : H <sub>2</sub> O (1 : 1 v/v)      | 2        | 94        |



Fig. 7 Proposed mechanism for the synthesis of biscoumarin catalyzed by camphor sulfonic acid.



Scheme 12 Coumarin synthesis by Pechmann condensation.

17. They conclude that solvent-free conditions and temperature 110 °C was suitable for the synthesis of desired products under the optimized reaction conditions. The effect of catalyst concentration was studied on model reaction and the results are presented in Table 18. They found that 10 mol% Zn<sub>0.925</sub>Ti<sub>0.075</sub>O catalyst was the most optimal for Pechmann condensation of ethylacetoacetate and phloroglucinol.

The reaction pathway for the synthesis of coumarin through Pechmann condensation is represented in Fig. 8. Initially,

Table 16 Catalytic screening for synthesis of substituted coumarin by Pechmann condensation reaction

| Entry | Catalyst                                     | Time (h) | Yield (%) |
|-------|--|----------|-----------|
| 1     | No catalyst                                  | 24       | Nil       |
| 2     | ZnO  | 5        | Nil       |
| 3     | Zn <sub>0.975</sub> Ti <sub>0.025</sub> O    | 3        | 37        |
| 4     | Zn <sub>0.950</sub> Ti <sub>0.050</sub> O    | 4        | 60        |
| 5     | <b>Zn<sub>0.925</sub>Ti<sub>0.075</sub>O</b> | <b>3</b> | <b>88</b> |
| 6     | Zn <sub>0.900</sub> Ti <sub>0.100</sub> O    | 3        | 88        |



**Table 17** Solvent screening for synthesis of substituted coumarin by Pechmann condensation reaction

| Entry | Solvent       | Temperature (°C) | Time (h) | Yield (%) |
|-------|---------------|------------------|----------|-----------|
| 1     | DCM           | 40               | 8        | 24        |
| 2     | Ethyl acetate | 78               | 8        | 16        |
| 3     | Acetonitrile  | 80               | 8        | 37        |
| 4     | Water         | 100              | 5        | 41        |
| 5     | Ethanol       | 78               | 5        | 63        |
| 6     | Toluene       | 110              | 10       | Nil       |
| 7     | DMF           | 150              | 10       | Nil       |
| 8     | Solvent-free  | 110              | 3        | 88        |
| 9     | Solvent-free  | 90               | 5        | 61        |
| 10    | Solvent-free  | 130              | 3        | 80        |

**Table 18** Effect of catalyst concentration for synthesis of substituted coumarin by Pechmann condensation reaction


| Entry | Catalyst amount (mol%) | Time (h) | Yield (%) |
|-------|------------------------|----------|-----------|
| 1     | 5                      | 5        | 67        |
| 2     | 10                     | 3        | 88        |
| 3     | 15                     | 3        | 88        |

reaction proceeds with the nucleophilic attack of the hydroxyl group of phloroglucinol on the activated ethylacetoacetate, resulting in the formation of intermediate. The formed intermediate rapidly undergoes cyclization through Lewis acid-catalyzed intramolecular condensation and followed by removal of water molecule give desired products.

A magnetic nanocatalyst of  $\text{Fe}_3\text{O}_4@\text{SiO}_2\text{-ZnCl}_2$  has been used for the synthesis of coumarin derivatives (7) *via* Pechmann condensation reaction of substituted phenols (5) and  $\beta$ -ketoesters (6) in excellent yield under solvent-free conditions by Rahimi and Soleimani (Scheme 13).<sup>61</sup> The advantages of this method are straightforward, easy work-up, catalyst reusability and leading to excellent yields.

Carrillo and his coworkers reported the synthesis of substituted coumarins (7) *via* one-pot reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in presence of propylsulfonic acid supported in FDU-5 (FDU-5-Pr-SO<sub>3</sub>H) as a catalyst (Scheme 14).<sup>62</sup> The catalytic activity of FDU-5-Pr-SO<sub>3</sub>H for the synthesis of substituted coumarins under optimized conditions was compared with other organic and inorganic catalysts summarized in Table 19.

Saffarian *et al.* reported the synthesis of coumarin containing 1,4-dihydropyridines (18) *via* condensation reaction between substituted aldehydes (2), 4-hydroxycoumarin (8) and ammonium acetate (17) under solvent-free conditions


**Fig. 8** Plausible mechanism for Pechmann condensation using EAA and phloroglucinol promoted by  $\text{Zn}_{0.925}\text{Ti}_{0.075}\text{O}$  NPs.**Scheme 13** Direct synthesis of coumarin derivatives.



Scheme 14 Synthesis of coumarin derivatives via Pechmann condensation of phenols with  $\beta$ -keto-ester catalyzed by FDU-5-Pr-SO<sub>3</sub>H.

Table 19 Comparison of different catalysts used for the synthesis of substituted coumarins

| Catalyst   | Catalyst amount (mol%) | Reaction time (min) | Temperature (°C) | Yield (%) | References |
|--|------------------------|---------------------|------------------|-----------|------------|
| FDU-5  | 1.65                   | 120                 | 130              | NR        | 62         |
| FDU-5-Pr-SO <sub>3</sub> H   | 1.65                   | 60                  | 130              | 97        | 62         |
| MCM-41-10SO <sub>3</sub> H   | 3.6                    | 120                 | 120              | 99        | 63         |
| SBA-15-10SO <sub>3</sub> H   | 2.0                    | 120                 | 120              | 88        | 63         |
| C@TiO <sub>2</sub> -SO <sub>3</sub> -SbCl <sub>2</sub>                   | 100.0                  | 35                  | 120              | 94        | 64         |
| <i>m</i> -ZrP  | 2.0                    | 240                 | 160              | 76        | 65         |
| SiO <sub>2</sub> -SnCl <sub>3</sub>                                      | 5.0                    | 35                  | 120              | 64        | 66         |
| FeCl <sub>3</sub> (ultrasound)   | 10.0                   | 20                  | 100              | 97        | 67         |
| Fe <sub>3</sub> O <sub>4</sub> @SiO <sub>2</sub> @Et-PhSO <sub>3</sub> H | 0.3                    | 120                 | 120              | 93        | 68         |
| CMK-5-SO <sub>3</sub> H  | 3.0                    | 130                 | 130              | 95        | 69         |
| SBA-15-Ph-Pr-SO <sub>3</sub> H   | 7.0                    | 130                 | 130              | 90        | 70         |
| <i>p</i> -TsOH   | 7.0                    | 130                 | 130              | 65        | 70         |
| Zr-TMS-BSA-10  | 10 wt%                 | 150                 | 150              | 81.4      | 71         |



Scheme 15 Catalytic synthesis of coumarin containing 1,4-DHPs.

(Scheme 15).<sup>72</sup> Simple protocol, simplicity of product isolation using water, decrease the temperature of reaction, reduce the use of hazardous solvents, excellent yield of products, eco-

Table 20 Optimization of reaction conditions

| Entry | Solvent                         | Temperature (°C) | Catalyst (mg) | Time (min) | Yield (%) |
|-------|---------------------------------|------------------|---------------|------------|-----------|
| 1     | —                               | 90               | —             | 90         | 30        |
| 2     | —                               | 90               | 5             | 30         | 80        |
| 3     | —                               | 90               | 10            | 20         | 86        |
| 4     | —                               | 90               | 15            | 20         | 85        |
| 5     | —                               | 100              | 10            | 20         | 81        |
| 6     | —                               | 80               | 10            | 20         | 85        |
| 7     | —                               | 60               | 10            | 30         | 70        |
| 8     | H <sub>2</sub> O                | Reflux           | 10            | 30         | 85        |
| 9     | EtOH                            | Reflux           | 10            | 45         | 70        |
| 10    | EtOAc                           | Reflux           | 10            | 90         | 20        |
| 11    | CH <sub>2</sub> Cl <sub>2</sub> | Reflux           | 10            | 90         | Nil       |
| 12    | <i>n</i> -Hexane                | Reflux           | 10            | 90         | Nil       |

friendly conditions and less reaction times are some beauties of present methodology. Firstly, to optimize the reaction conditions they conducted a model reaction between 4-methyl benzaldehyde, 4-hydroxycoumarin and ammonium acetate. They observed that 10 mg of the Fe<sub>3</sub>O<sub>4</sub>@SiO<sub>2</sub>@(CH<sub>2</sub>)<sub>3</sub>-urea-quinoline sulfonic acid chloride at 80 °C under solvent free conditions supplied the best results as presented in Table 20. They performed the model reaction also in the presence of

Table 21 Screening the model reaction in the presence of desired catalyst

| Entry | Catalyst   | Yield (%) |
|-------|--|-----------|
| 1     | Fe <sub>3</sub> O <sub>4</sub>   | 40        |
| 2     | Fe <sub>3</sub> O <sub>4</sub> @SiO <sub>2</sub>   | 40        |
| 3     | Fe <sub>3</sub> O <sub>4</sub> @SiO <sub>2</sub> @(CH <sub>2</sub> ) <sub>3</sub> -urea-quinoline                        | 65        |
| 4     | Fe <sub>3</sub> O <sub>4</sub> @SiO <sub>2</sub> @(CH <sub>2</sub> ) <sub>3</sub> -urea-quinoline sulfonic acid chloride | 85        |






Scheme 16 Pechmann reaction of resorcinol with ethylacetoacetate to produce 7-hydroxy-4-methylcoumarin.

Table 22 Condensation reaction of resorcinol with ethyl acetoacetate using various heterogeneous solid acids catalysts

| Catalyst                | Acidity | Yield (%) |
|-------------------------|---------|-----------|
| Amberlyst-15            | 4.30    | 97        |
| H- $\beta$              | 1.01    | 21        |
| TS-OS-SO <sub>3</sub> H | 1.24    | 44        |



Scheme 17 Synthesis of substituted coumarins.

related intermediates of the Fe<sub>3</sub>O<sub>4</sub>@SiO<sub>2</sub>@(CH<sub>2</sub>)<sub>3</sub>-urea-quinoline sulfonic acid chloride at 80 °C under solvent free conditions for 20 min and results are summarized in Table 21.

A suitable protocol for synthesis of coumarins derivatives (7) was reported by Bouasla and his coworkers *via* one-pot reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in presence of heterogeneous solid acid catalyst *viz.* Amberlyst-15 in solvent-free medium under microwave irradiation (Scheme 16).<sup>73</sup> Initially, they conducted a model reaction between resorcinol and ethylacetoacetate as model substrate. They observed that by changing the reaction time from 5 min to 20 min, a maximum yield of 97% was obtained and no reaction was observed in absence of catalyst as summarized in Table 22. The plausible mechanism for the reaction is shown in Fig. 9.

An efficient method for the synthesis of 3-carboxycoumarins (20) was reported *via* Knoevenagel condensation reaction between substituted aldehydes (2) and Meldrum's acid (19) in presence of polymeric magnetic nanocatalyst by Maleki *et al.* (Scheme 17).<sup>74</sup> This method has many advantages such as less reaction time, high yield and easy isolation of catalyst. The plausible mechanism for the reaction is shown in Fig. 10.

Suryawanshi and his coworkers reported the synthesis of coumarins (7) *via* Pechmann condensation reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in presence of reusable polymeric SO<sub>3</sub>H-functionalized cation exchange resins *viz.* Amberlite IR-120, Dowex 50, X-8100 and Tulsion T-42 (Scheme 18).<sup>75</sup> Excellent yield of products, short reaction time,




Fig. 9 A plausible mechanism for the Pechmann condensation of phenol and ethylacetoacetate in presence of Amberlyst-15.










Fig. 10 Proposed mechanism for the synthesis of 3-carboxy coumarins in presence of polymeric magnetic nanocatalyst.



Scheme 18 The Pechmann condensation between resorcinol and ethyl acetoacetate catalyzed by different cation exchange resins.



Scheme 19 Synthesis of substituted coumarins.

easy work-up and use of safe catalyst are some advantages of present methodology.

Rostami and Zare reported the synthesis of substituted coumarins (7) *via* one-pot reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in presence of carbonized sugarcane

bagasse (CSCB) as a new and efficient solid acid catalyst (Scheme 19).<sup>76</sup> Simple preparation of catalyst, safe handling, inexpensive, excellent yield of products, catalyst reusability, solvent-free and easy work-up are some benefits of present methodology. Initially, model reaction was considered between

Table 23 Optimization of reaction conditions for AHS@CSCB catalyzed Pechmann condensation between 1,3 dihydroxy phenol and ethyl acetoacetate

| Entry | Concentration of catalyst (mg) | Solvent          | Temperature (°C) | Time (min) | Yield (%) |
|-------|--------------------------------|------------------|------------------|------------|-----------|
| 1     | None                           | H <sub>2</sub> O | Reflux           | 120        | 0         |
| 2     | 30                             | H <sub>2</sub> O | Reflux           | 60         | 50        |
| 3     | 30                             | EtOH             | Reflux           | 40         | 80        |
| 4     | 30                             | Solvent-free     | 80               | 15         | 91        |
| 5     | 30                             | Solvent-free     | 70               | 30         | 89        |
| 6     | 30                             | Solvent-free     | 120              | 120        | 26        |
| 7     | 20                             | Solvent-free     | 15               | 15         | 92        |
| 8     | 10                             | Solvent-free     | 5                | 5          | 92        |







Fig. 11 Plausible mechanism for the synthesis of coumarins, biscoumarins and benzoxanthenes in the presence of AHS@CSCB.



Scheme 20 Synthesis of substituted coumarins.

3-hydroxyphenol and ethylacetoacetate and the effect of different solvents, temperature and amount of catalyst was investigated and results were summarized in Table 23. The plausible mechanism for the reaction is shown in Fig. 11.

Sun and his coworkers reported the synthesis of substituted coumarins (7) *via* Pechmann condensation reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) catalyzed from Wells–Dawson heteropolyacid ( $H_6P_2W_{18}O_{62}$ ). This work provides a novel, cheaper and safer way to synthesize coumarins unsubstituted on the pyranic nucleus (Scheme 20).<sup>77</sup> Initially, they optimized the reaction conditions by exploring a model

reaction between 2-methyl-3-hydroxyphenol and ethyl 3,3-diethoxypropionate. The effect of the temperature and reaction time were investigated and results were summarized in Table 24. The comparison of efficiencies of various catalysts used in the synthesis of 7-hydroxy-8-methylcoumarin was summarized in Table 25. The plausible mechanism for the reaction is shown in Fig. 12.

An efficient and facile synthesis of coumarins (7) was reported in excellent yields *via* Pechmann condensation reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) under solvent-free medium using both conventional method and microwave irradiation in less reaction times in presence of

Table 24 Optimization of Pechmann condensation reaction for the synthesis of 7-hydroxy-8-methylcoumarin

| Entry | Catalyst concentration | Temperature (°C) | Time (h) | Yield (%) |
|-------|------------------------|------------------|----------|-----------|
| 1     | 0.10                   | 100              | 3        | 75        |
| 2     | 0.25                   | 100              | 3        | 87        |
| 3     | 0.50                   | 100              | 3        | 86        |
| 4     | 1.00                   | 100              | 3        | 84        |
| 5     | 0.25                   | 80               | 3        | 74        |
| 6     | 0.25                   | 90               | 3        | 90        |
| 7     | 0.25                   | 90               | 2        | 72        |
| 8     | 0.25                   | 90               | 4        | 89        |
| 9     | 0.25                   | 90               | 3        | 84        |
| 10    | 0.25                   | 90               | 3        | 90        |
| 11    | 0.25                   | 90               | 3        | 95        |
| 12    | 0.25                   | 90               | 3        | 95        |

Table 25 Synthesis of 7-hydroxy-8-methylcoumarin mediated by different catalysts

| Entry | Catalyst  | Time (h) | Yield (%) |
|-------|---|----------|-----------|
| 1     | MeSO <sub>3</sub> H   | 3        | 20        |
| 2     | MeSO <sub>3</sub> H/basic Al <sub>2</sub> O <sub>3</sub>      | 3        | 30        |
| 3     | MeSO <sub>3</sub> H/neutral Al <sub>2</sub> O <sub>3</sub>    | 3        | 34        |
| 4     | MeSO <sub>3</sub> H/acidic Al <sub>2</sub> O <sub>3</sub>     | 3        | 80        |
| 5     | Acidic Al <sub>2</sub> O <sub>3</sub>                         | 3        | 30        |
| 6     | Al <sub>2</sub> O <sub>3</sub>                                | 2        | 10        |
| 7     | AlCl <sub>3</sub> /MeSO <sub>3</sub> H                        | 2        | 12        |
| 8     | ZnCl <sub>2</sub> /MeSO <sub>3</sub> H                        | 2        | 5         |
| 9     | Cu(CH <sub>3</sub> CN) <sub>4</sub> PF <sub>6</sub>           | 2        | 10        |
| 10    | H <sub>6</sub> P <sub>2</sub> W <sub>18</sub> O <sub>62</sub> | 2        | 82        |
| 11    | FeCl <sub>3</sub>   | 3        | 8         |
| 12    | TiCl <sub>4</sub>   | 3        | 5         |







Fig. 12 Possible mechanism for the synthesis of coumarins catalyzed from Wells–Dawson heteropolyacid ( $H_6P_2W_{18}O_{62}$ ).




Scheme 21 Synthesis of coumarins by using cellulose sulfuric acid as a solid acid catalyst.

Table 26 Comparison of efficiency of cellulose sulfuric acid with reported catalysts

| Entry | Catalyst                        | Yield (%) |
|-------|---------------------------------|-----------|
| 1     | Cellulose sulfuric acid         | 97        |
| 2     | Silica sulfuric acid            | 92        |
| 3     | <i>p</i> -Toluene sulfonic acid | 85        |
| 4     | Sulfuric acid in acetic acid    | 55        |
| 5     | No catalyst                     | 15        |

cellulose sulfuric acid by Kuram *et al.* (Scheme 21).<sup>78</sup> The efficiency of the cellulose sulfuric acid compared with other catalysts is summarized in Table 26. It was found that cellulose sulfuric acid is a more efficient and superior catalyst over other acidic catalysts with respect to reaction time and yield.

Palaniappan and John *et al.* reported the synthesis of substituted coumarins (7) *via* one-pot reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in presence of novel polyaniline–fluoroboric acid–dodecylhydrogensulfate (PANI–HBF<sub>4</sub>–DHS) as reusable catalyst (Scheme 22).<sup>79</sup>



Scheme 22 Synthesis of substituted coumarins.







Scheme 23 Synthesis of substituted coumarins catalyzed by PeSA.

Table 27 Comparison of activity of the PeSA catalyst with some other reported catalysts

| Entry | Catalyst                | Condition           | Yield (%) | Time (min) | References |
|-------|-------------------------|---------------------|-----------|------------|------------|
| 1     | PeSA                    | 110 °C/Solvent-free | 97        | 15         | 80         |
| 2     | ASA                     | 110 °C/Solvent-free | 85        | 30         | 81         |
| 3     | CMK-5-SO <sub>3</sub> H | 110 °C/Solvent-free | 95        | 20         | 82         |

Kolvari and his coworkers reported the synthesis of substituted coumarins (7) *via* one-pot reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in presence of perlite sulfonic acid (perlite-SO<sub>3</sub>H (PeSA)) as heterogeneous reusable solid acid catalysts (Scheme 23).<sup>80</sup> Inexpensive, ease of preparation, more stability and reusability, low toxicity and easy of handling are some advantages of present catalytic systems. To show the advantages of current protocol in comparison with reported results in literature was summarized in Table 27. They

Scheme 24 W/ZrO<sub>2</sub> solid acid catalyzed synthesis of substituted coumarins.

Scheme 25 TfOH-mediated preparation of coumarins.






Fig. 13 Plausible mechanism for the synthesis of coumarins in the presence of TfOH as Brønsted acid catalyst.



Scheme 26 Synthesis of coumarins using  $\text{HClO}_4 \cdot \text{SiO}_2$  under solvent-free conditions.

Scheme 27 Synthesis of coumarin by xanthan sulfuric acid as a solid acid catalyst.

Table 28 Effect of catalysts on yield of synthesis of substituted coumarins

| Entry | Catalyst                     | Quantity | Yield (%) |
|-------|------------------------------|----------|-----------|
| 1     | Xanthan sulfuric acid        | 0.08 g   | 96        |
| 2     | Silica sulfuric acid         | 0.08 g   | 92        |
| 3     | Methane sulfonic acid        | 0.1 mmol | 86        |
| 4     | Sulfuric acid in acetic acid | 0.1 mmol | 56        |
| 5     | No catalyst                  | None     | 10        |

Table 29 Influence of the catalytic amounts of xanthan sulfuric acid for synthesis of substituted coumarins

| Entry | Catalyst (g) | Time (min) | Yield (%) |
|-------|--------------|------------|-----------|
| 1     | None         | 60         | Nil       |
| 2     | 0.01         | 20         | 28        |
| 3     | 0.03         | 20         | 51        |
| 4     | 0.05         | 20         | 79        |
| 5     | 0.08         | 40         | 96        |
| 6     | 0.08         | 20         | 96        |

found that PeSA showed greater activity than some other than some other heterogeneous catalysts.

Reddy *et al.* reported the synthesis of substituted coumarins (7) *via* Pechmann condensation reaction between substituted


phenols (5) and  $\beta$ -ketoesters (6) in presence of  $\text{W/ZrO}_2$  solid acid catalyst (Scheme 24).<sup>83</sup>

Kim *et al.* reported the synthesis of substituted coumarins (22) *via* condensation reaction between substituted phenols (5) and allenes (21) in the presence of TFOH as Bronsted acid catalyst in excellent yield (Scheme 25).<sup>84</sup> The plausible mechanism for the reaction is shown in Fig. 13.

Maheswara and his coworkers synthesized substituted coumarins (7) *via* Pechmann condensation reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in presence of heterogeneous recyclable catalyst ( $\text{HClO}_4 \cdot \text{SiO}_2$ ) under solvent-free medium (Scheme 26).<sup>85</sup> Cost-effective, less reaction time and operational simplicity are some benefits of present methodology.



Kuram *et al.* reported the synthesis of substituted coumarins (7) *via* Pechmann condensation reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in the presence of xanthan sulfuric acid as a solid acid catalyst under solvent-free conditions (Scheme 27).<sup>86</sup> They found that this method is very simple, inexpensive, less reaction time and catalyst could be reused. The effect of catalyst on the yield of products was summarized in Table 28. They also investigated the efficiency of the XSA compared to various sulphur analog acidic catalysts and results are summarized in Table 29.

Singh and his coworkers reported the synthesis of substituted coumarins (7) *via* Pechmann condensation reaction between substituted phenols (6) and  $\beta$ -ketoesters (5) in presence of sulphamic acid (Scheme 28).<sup>87</sup>




Scheme 28 Pechmann condensation using sulphamic acid (SA) as catalyst.



Scheme 29 Synthesis of coumarins via von Pechmann condensation of phenols with  $\beta$ -ketoesters induced by  $\text{In}(\text{III})\text{Cl}_3$ .

Scheme 30 Synthesis of 7-hydroxy-4-methyl coumarin.



Scheme 31 Synthesis of substituted coumarins.

Table 30 Effect of different solvents on Pechmann condensation reaction for synthesis of substituted coumarins

| Solvent      | Time (h) | Temperature ( $^{\circ}\text{C}$ ) | Yield (%) |
|--------------|----------|------------------------------------|-----------|
| Nitrobenzene | 4        | 120                                | 25        |
| Toluene      | 15       | 120                                | 34        |
| Solvent-free | 4        | 120                                | 51        |
| Solvent-free | 4        | 150                                | 76        |

Table 31 Effect of  $\text{SD-SO}_3\text{H}$  catalyst concentration on the yield of product


| Entry | Catalyst (g) | Temperature ( $^{\circ}\text{C}$ ) | Time (min) | Yield (%) |
|-------|--------------|------------------------------------|------------|-----------|
| 1     | —            | 90                                 | 120        | 0         |
| 2     | 0.025        | 90                                 | 120        | 40        |
| 3     | 0.05         | 90                                 | 75         | 72        |
| 4     | 0.075        | 90                                 | 120        | 70        |
| 5     | 0.10         | 90                                 | 120        | 60        |
| 6     | 0.15         | 90                                 | 120        | 60        |

Bose *et al.* reported the synthesis of substituted coumarins (7) via Pechmann condensation reaction between substituted phenols (6) and  $\beta$ -ketoesters (5) in presence of indium(III) chloride as an efficient catalyst (Scheme 29).<sup>88</sup>

An efficient and facile synthesis of substituted coumarins (7) was reported by one-pot reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in presence of new magnetic

nanocomposites of  $\text{ZrO}_2\text{-Al}_2\text{O}_3\text{-Fe}_3\text{O}_4$  as green solid acid catalysts (Scheme 30).<sup>89</sup>

Mesoporous zirconium phosphate (*m*-ZrP) is used as solid acid catalyst for the synthesis of substituted coumarins (7) via Pechmann condensation reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in both conventional heating as well as microwave

Scheme 32 Synthesis of coumarins catalyzed by  $\text{SD-SO}_3\text{H}$  under solvent-free conditions.

**Table 32** Effect of solvents and temperature on the synthesis of substituted coumarins

| Entry | Solvent                         | Temperature (°C) | Time (min) | Yield (%) |
|-------|---------------------------------|------------------|------------|-----------|
| 1     | —                               | 90               | 75         | 72        |
| 2     | CHCl <sub>3</sub>               | Reflux           | 200        | 35        |
| 3     | CH <sub>3</sub> CN              | Reflux           | 200        | 10        |
| 4     | CH <sub>2</sub> Cl <sub>2</sub> | Reflux           | 200        | 20        |
| 5     | THF                             | Reflux           | 200        | 0         |
| 6     | MeOH                            | Reflux           | 200        | 0         |
| 7     | H <sub>2</sub> O                | Reflux           | 200        | 0         |
| 8     | —                               | 70               | 120        | 70        |
| 9     | —                               | 110              | 25         | 91        |
| 10    | —                               | 130              | 25         | 92        |

assisted method by Sinhamahapatra and his coworkers (Scheme 31).<sup>90</sup> The effect of solvent on reaction was summarized in Table 30.


Tahanpesar and Sarami reported the synthesis of substituted coumarins (7) *via* one-pot Pechmann

condensation reaction between substituted phenols (5) and  $\beta$ -ketoesters (6) in presence of sulfonated sawdust (SD-SO<sub>3</sub>H) as solid acid catalyst under solvent-free conditions (Scheme 32).<sup>91</sup> Further, they observed the catalytic efficiency of SD-SO<sub>3</sub>H on the yield of product and results were presented in Table 31. They also observed the effects of different solvents *viz.* CHCl<sub>3</sub>, CH<sub>3</sub>CN, CH<sub>2</sub>Cl<sub>2</sub>, THF, MeOH and H<sub>2</sub>O and temperature on the synthesis of desired products and results were presented in Table 32.

The plausible mechanism for the synthesis of substituted coumarins was presented in Fig. 14. The comparison of catalytic activity of SD-SO<sub>3</sub>H with other catalyst found in literature was presented in Table 33.

## Conclusion and future prospects

This review article summarized the synthesis of substituted coumarins using solid acid catalysts. Benefits of these methods include clean reaction profiles, minimization of side products,



**Fig. 14** The plausible mechanism of formation of 7-hydroxy-4-methylcoumarin in presence of sulfonated sawdust (SD-SO<sub>3</sub>H) as solid acid catalyst.

**Table 33** Comparison of catalytic activity of SD-SO<sub>3</sub>H with some other catalysts

| Entry | Catalyst  | Temperature (°C) | Time (min) | Yield (%) | References |
|-------|---|------------------|------------|-----------|------------|
| 1     | SD-SO <sub>3</sub> H  | 110              | 9          | 98        | 91         |
| 2     | <i>m</i> -ZrP   | 160              | 240        | 94        | 92         |
| 3     | SCZ   | 120              | 143        | 87        | 93         |
| 4     | ASA   | 100              | 30         | 98        | 94         |
| 5     | CMK-5-SO <sub>3</sub> H   | 130              | 20         | 95        | 95         |
| 6     | H <sub>6</sub> P <sub>2</sub> W <sub>18</sub> O <sub>62</sub> ·24H <sub>2</sub> O | 130              | 42         | 87        | 96         |
| 7     | Zeolite-E4a   | 110              | 180        | 97        | 97         |
| 8     | HClO <sub>4</sub> ·SiO <sub>2</sub>   | 130              | 35         | 95        | 98         |



efficient and facile experimental procedures and inexpensive. This review is endeavouring to find potential future directions in the development of more potent and specific analogs of nitrogen and oxygen containing heterocyclic compounds for the biological target by the use of heterogeneous catalysts. The information illustrated in this review also encourage organic chemist for the design of novel molecules to identify many more biologically active heterocycles for the benefit of humanity.

## Author contributions

Conceptualization: Susheel Gulati, Rajvir Singh, Suman Sangwan, formal analysis: Susheel Gulati, Suman Sangwan, investigation: Susheel Gulati, Rajvir Singh, Suman Sangwan, supervision: Susheel Gulati, Rajvir Singh, validation: Susheel Gulati, Suman Sangwan, writing-original draft: Susheel Gulati, writing-review & editing: Susheel Gulati, Rajvir Singh, Suman Sangwan.

## Conflicts of interest

Authors declared that there is no conflict of interest regarding the publication of this paper.

## Abbreviations

|         |   |
|---------|---|
| TSA     | Tungstate sulphuric acid                        |
| MW      | Microwave irradiation                           |
| PVPHS   | Poly(4-vinylpyridinium) hydrogen sulfate        |
| SCCMNPs | Sulfonated carbon-coated magnetic nanoparticles |
| SSA     | Silica sulphuric acid                           |
| XSA     | Xanthan sulphuric acid                          |

## Acknowledgements

The authors are thankful to the Department of Chemistry, Chaudhary Charan Singh Haryana Agricultural University, Hisar for providing the necessary facilities. Financial assistance from Department of Science and Technology (DST), New Delhi, India is gratefully acknowledged.

## References

- X. He, Y. Shang, Y. Zhou, Z. Yu, G. Han, W. Jin and J. Chen, *Tetrahedron*, 2015, **71**, 863–868.
- N. Zhang, S. Ayril-Kaloustian, T. Nguyen, R. Hernandez, J. Lucas, C. Discafani and C. Beyer, *Bioorg. Med. Chem.*, 2009, **17**, 111–118.
- X. He, Y. Chen, J. Shi, W. Tang, Z. Pan, Z. Dong, B. Song, J. Li and X. Liu, *Bioorg. Med. Chem.*, 2014, **22**, 3732–3738.
- L. Huang, X. Yuan, D. Yu, K. Lee and C. Chen, *Virology*, 2005, **332**, 623–628.
- F. Borges, F. Roleira, N. Milhazes, L. Santana and E. Uriarte, *Curr. Med. Chem.*, 2005, **12**, 887–916.
- H. Abd El-Wahab, M. Abd El-Fattah, N. Abd El-Khalik, H. Nassar and M. Abdelall, *Prog. Org. Coat.*, 2014, **77**, 1506–1511.
- G. Zhao, C. Peng, W. Du and S. Wang, *Fitoterapia*, 2013, **89**, 250–256.
- T. Smyth, V. Ramachandran and W. Smyth, *Int. J. Antimicrob. Agents*, 2009, **33**, 421–426.
- M. Razavi, H. Nazemiyeh, A. Delazar, R. Hajiboland, M. Rahman, S. Gibbons, L. Nahar and S. Sarker, *Phytochem. Lett.*, 2008, **1**, 159–162.
- N. Ballin and A. Sørensen, *Food Control*, 2014, **38**, 198–203.
- N. Khaligh, *Catal. Sci. Technol.*, 2012, **2**, 1633–1636.
- A. Fischer, C. Cremer and E. H. K. Stelzer, *Appl. Opt.*, 1995, **34**, 1989–2003.
- G. Cravotto, G. M. Nano, G. Palmisano and S. Tagliapietra, *Tetrahedron: Asymmetry*, 2001, **12**, 707–709.
- I. B. Masesane and Z. Y. Desta, *Beilstein J. Org. Chem.*, 2012, **8**, 2166–2175.
- B. M. Reddy, M. Patil and P. Lakshmanan, *J. Mol. Catal. A: Chem.*, 2006, **256**, 290–294.
- B. C. Raju, T. H. Babu and J. M. Rao, *Indian J. Chem. B*, 2009, **48**, 120–123.
- N. Ahmed and Z. Siddiqui, *J. Mol. Catal. A: Chem.*, 2014, **387**, 45–56.
- S. Sharma, P. Parikh and R. Jasra, *Appl. Catal., A*, 2010, **386**, 34–42.
- E. Kolvari, N. Koukabi and M. Hosseini, *J. Mol. Catal. A: Chem.*, 2015, **397**, 68–75.
- S. Ghodke and U. Chudasama, *Appl. Catal., A*, 2013, **453**, 219–226.
- B. Movassagh, L. Tahershamsi and A. Mobaraki, *Tetrahedron Lett.*, 2015, **56**, 1851–1854.
- D. R. Naikwadi, B. D. Bankar, K. Ravi and A. V. Biradar, *Res. Chem. Intermed.*, 2021, 1–13, DOI: 10.1007/s11164-021-04499-3.
- B. Karami, M. Farahi, N. Farmani and H. M. Tanuraghaj, *New J. Chem.*, 2015, 1–5.
- N. G. Khaligh, *Ultrason. Sonochem.*, 2013, **20**, 1062–1068.
- Z. Z. Akbari, S. Dastmalchi, L. Edijlali, L. Dinparast and M. Es'haghi, *Appl. Organomet. Chem.*, 2019, 1–14.
- A. N. Nadaf and K. Shivashankar, *J. Heterocycl. Chem.*, 2018, **55**, 1375–1381.
- A. Zhu, S. Bai, L. Li, M. Wang and J. Wang, *Catal. Lett.*, 2015, **145**, 1089–1093.
- A. Zhu, M. Wang, L. Li and J. Wang, *RSC Adv.*, 2015, **5**, 73974–73979.
- Z. Samiei, S. S. Amiri and Z. Azizi, *Mol. Diversity*, 2019, 1–20.
- M. A. Nasser and S. M. Sadeghzadeh, *J. Iran. Chem. Soc.*, 2014, **11**, 27–33.
- Z. Abbasi, S. Rezayati, M. Bagheri and R. Hajinasiri, *Chin. Chem. Lett.*, 2017, **28**(1), 75–82.
- F. K. Esfahani, D. Zareyee and R. Yousef, *ChemCatChem*, 2014, **6**(12), 3333–3337.
- E. D. Glendening, *NBO 6.0*. Theoretical Chemistry Institute, University of Wisconsin, Madison, 2013.
- D. Zareyee and M. Serehneh, *J. Mol. Catal. A: Chem.*, 2014, **1**, 88–91.





- 35 M. Samadzadeh, S. Nouri and F. K. Moghadam, *Res. Chem. Intermed.*, 2016, **42**(6), 6089–6103.
- 36 R. Sun, Y. Gao, Y. Ma, G. Yang and Y. Li, *J. Iran. Chem. Soc.*, 2017, **14**, 737–742.
- 37 H. M. Altass and A. E. R. S. Khder, *React. Kinet., Mech. Catal.*, 2018, **125**(1), 227–243.
- 38 A. Mirosanloo, D. Zareyee and M. A. Khalilzadeh, *Appl. Organomet. Chem.*, 2018, **32**(12), e4546.
- 39 N. H. Jadhav, S. S. Sakate, N. K. Rasal and R. A. Pawar, *ACS Omega*, 2019, **4**(5), 8522–8527.
- 40 S. Pakdel, B. Akhlaghinia and A. Mohammadinezhad, *Chem. Afr.*, 2019, **2**(3), 10.
- 41 S. A. Khan, S. B. Khan, A. M. Asiri and I. Ahmad, *Nanoscale Res. Lett.*, 2016, **11**, 345.
- 42 F. K. Esfahani, D. Zareyee and R. Zareyee, *ChemCatChem*, 2014, **6**(12), 3333–3337.
- 43 V. Vahabi and F. Hatamjafari, *Molecules*, 2014, **19**(9), 13093–13103.
- 44 D. Zareyee and M. Serehneh, *J. Mol. Catal. A: Chem.*, 2014, **391**, 88–91.
- 45 S. S. Kumbar, K. M. Hosamani, G. C. Gouripur and S. D. Joshi, *R. Soc. Open Sci.*, 2018, **5**, 172416.
- 46 F. M. Moghaddam, M. Eslami and G. Hoda, *Sci. Rep.*, 2020, **10**, 20968.
- 47 L. N. Nasirmahale, O. G. Jolodar, F. Shirini and H. Tajik, *Polycyclic Aromat. Compd.*, 2021, **41**, 199–210.
- 48 H. Mehrabi and H. Abusaidi, *J. Iran. Chem. Soc.*, 2010, **7**(4), 890–894.
- 49 J. M. Khurana and K. Vij, *J. Chem. Sci.*, 2012, **124**(4), 907–912.
- 50 S. Abdolmohammadi and S. Balalaie, *Tetrahedron Lett.*, 2007, **48**(18), 3299–3303.
- 51 A. R. Montaghani and N. Montazeri, *Orient. J. Chem.*, 2014, **30**(3), 1361–1364.
- 52 J. M. Khurana and S. Kumar, *Monatsh. Chem.*, 2010, **141**(5), 561–564.
- 53 M. Kidwai, V. Bansal, P. Mothra, S. Saxena, R. K. Somvanshi, S. Dey and T. P. Singh, *J. Mol. Catal.*, 2007, **268**(1), 76–81.
- 54 A. Zhu, Sh Bai, L. Li, M. Wang and J. Wang, *Catal. Lett.*, 2015, **145**, 1089–1093.
- 55 K. P. Boroujeni and P. Ghasemi, *Catal. Commun.*, 2014, **37**, 50–54.
- 56 A. Kiasat and L. Hemat-Alian, *Res. Chem. Intermed.*, 2015, **41**(2), 873–880.
- 57 K. M. Khan, S. Iqbal, M. A. Lodhi, G. M. Maharvi, Z. Ullah, M. I. Choudhary, A. ur Rahman and S. Perveen, *Bioorg. Med. Chem.*, 2004, **12**(8), 1963–1968.
- 58 A. P. Katariya, S. U. Deshmukh, S. U. Tekale, M. V. Katariya and R. P. Pawar, *Lett. Appl. NanoBioSci.*, 2021, **10**(3), 2525–2534.
- 59 G. Kaur, D. Singh, A. Singh and B. Banerjee, *Synth. Commun.*, 2020, 1–14, DOI: 10.1080/00397911.2020.1856877.
- 60 N. H. Jadhav, S. S. Sakate, N. K. Rasal, D. R. Shinde and R. A. Pawar, *ACS Omega*, 2019, **4**, 8522–8527.
- 61 S. Rahimi and E. Soleimani, *Results Chem.*, 2020, **2**, 1–7.
- 62 G. G. Carrillo, J. Gonzalez, M. J. E. Legaspi, G. J. L. Lopez, I. A. A. Villarreal, S. G. C. Magana, F. J. M. Martinez and R. M. Valencia, *Microporous Mesoporous Mater.*, 2020, **307**, 1–10.
- 63 A. E. R. S. Khder, S. A. Ahmed, K. S. Khairou and H. M. Altass, *J. Porous Mater.*, 2018, **25**, 1–13.
- 64 M. Kour and S. Paul, *Monatsh. Chem.*, 2017, **148**, 327–337.
- 65 A. Sinhamahapatra, N. Sutradhar, S. Pahari, H. C. Bajaj and A. B. Panda, *Appl. Catal., A*, 2011, **394**, 93–100.
- 66 R. Sun, Y. Gao, Y. Ma, G. Yang and Y. Li, *J. Iran. Chem. Soc.*, 2017, **14**, 737–742.
- 67 K. C. Prousis, N. Avlonitis, G. A. Heropoulos and T. Calogeropoulou, *Ultrason. Sonochem.*, 2014, **21**, 937–942.
- 68 A. Mobaraki, S. Yasham and B. Movassagh, *Synlett*, 2015, **26**, 1263–1268.
- 69 D. Zareyee and M. Serehneh, *J. Mol. Catal. A: Chem.*, 2014, **391**, 88–91.
- 70 B. Karimi and D. Zareyee, *Org. Lett.*, 2008, **10**, 3989–3992.
- 71 S. Selvakumar, M. Chidambaram and A. P. Singh, *Catal. Commun.*, 2007, **8**, 777–783.
- 72 H. Saffarian, F. Karimi, M. Yarie and M. A. Zolfigol, *J. Mol. Struct.*, 2020, **1224**, 1–12.
- 73 S. Bouasla, J. A. Gahete, D. Esquivel, M. I. Lopez, C. J. Sanchidrian, M. Teguche and F. J. R. Salguero, *Molecules*, 2017, **22**, 1–8.
- 74 A. Maleki, P. Ravaghi, M. Aghaie and H. Movahed, <http://sciforum.net/conference/ecsoc-20>.
- 75 V. D. Suryawanshi, A. H. Gore, P. V. Anbhule, S. R. Patil and G. B. Kolekar, *J. Shivaji Univ.*, 2017, **42**(1), 40–45.
- 76 E. Rostami and S. H. Zare, *ChemistrySelect*, 2019, **4**, 13295–13303.
- 77 Y. F. Sun, J. M. Liu, J. Sun, Y. T. Haung, J. Lu, M. M. Li, N. Jin, X. F. Dai and B. Fan, *Molecules*, 2018, DOI: 10.20944/preprints201809.0349.v1.
- 78 B. S. Kuram, J. V. Madhav, S. Vijaya Laxmi, B. Rajitha, Y. Thirupathi Reddy, P. Narsimha Reddy and P. A. Crooks, *Synth. Commun.*, 2010, 1770–1777, <http://www.tandfonline.com/loi/lscy20>.
- 79 S. Palaniappan and A. John, *J. Mol. Catal.*, 2005, **233**, 9–15.
- 80 E. Kolvari, N. Koukabi and M. M. Hosseini, *J. Mol. Catal.*, DOI: 10.1016/j.molcata.2014.10.026.
- 81 A. Amoozadeh, M. Ahmadzadeh and E. Kolvari, *J. Chem.*, 2012, **2013**, 1–7.
- 82 D. Zareyee and M. Serehneh, *J. Mol. Catal. A: Chem.*, 2014, **391**, 88–91.
- 83 B. M. Reddy, V. R. Reddy and D. Giridhar, *Synth. Commun.*, 2006, **31**(23), 3603–3607.
- 84 S. Kim, D. Kang, C. H. Lee and P. H. Lee, *J. Org. Chem.*, 2012, **77**, 6530–6537.
- 85 M. Maheswara, V. Siddaiah, G. Lakishmi, V. Damu, Y. K. Rao and C. V. Rao, *J. Mol. Catal. A: Chem.*, 2006, **255**, 49–52.
- 86 B. S. Kuram, J. V. Madhav and B. Rajitha, *Synth. Commun.*, 2011, **42**(12), 1770–1777.
- 87 P. R. Singh, D. U. Singh and S. D. Samant, *Synlett*, 2004, **11**, 909–1912.
- 88 D. S. Bose, A. P. Rudradas and M. H. Babu, *Tetrahedron Lett.*, 2002, **43**, 9195–9197.
- 89 A. Wang, X. Lu, Z. Su and H. Jing, *Catal. Sci. Technol.*, 2014, **4**, 71–80.



Review

- 90 A. Sinhamahapatra, N. Sutradhar, S. Pahari, H. C. Bajaj and A. B. Panda, *Appl. Catal., A*, 2011, **394**, 93–100.
- 91 E. Tahanpesar and L. Sarami, *Russ. J. Gen. Chem.*, 2015, **85**(9), 2135–2140.
- 92 A. Sinhamahapatra, N. Sutradhar, S. Pahari, H. C. Bajaj and A. B. Panda, *Appl. Catal., A*, 2011, **394**, 93.
- 93 B. M. Reddy, M. K. Patil and P. J. Lakshmanan, *J. Mol. Catal. A: Chem.*, 2006, **256**, 290.
- 94 A. Amoozadeh, M. Ahmadzadeh and E. Kolvari, *J. Chem.*, 2013, **2013**, 6.
- 95 D. Zareyee and M. J. Serehneh, *J. Mol. Catal. A: Chem.*, 2014, **391**, 88.
- 96 G. P. Romanelli, D. Bennardi, D. M. Ruiz, G. Baronetti, H. J. Thomas and J. C. Autino, *Tetrahedron Lett.*, 2004, **45**, 8935.
- 97 A. Hegedüs and Z. Hell, *Catal. Lett.*, 2006, **112**, 105.
- 98 M. Maheswara, V. Siddaiah, G. L. V. Damu, Y. K. Rao and C. V. Rao, *J. Mol. Catal. A: Chem.*, 2006, **255**, 49.

