Analyst



View Article Online

CORRECTION

Check for updates

Cite this: Analyst, 2021, 146, 6665

Correction: Spatiotemporal distribution and speciation of silver nanoparticles in the healing wound

DOI: 10.1039/d1an90083h

rsc.li/analyst

Correction for 'Spatiotemporal distribution and speciation of silver nanoparticles in the healing wound' by Marco Roman *et al.*, *Analyst*, 2020, **145**, 6456–6469, DOI: 10.1039/D0AN00607F.

The authors regret that Fig. 3D, 5F and 7 were shown incorrectly in the original article. The correct versions of Fig. 3D, 5F and 7 are shown below.

^aCa' Foscari University of Venice, Department of Environmental Sciences, Informatics and Statistics (DAIS), Via Torino 155, 30172 Venice Mestre, Italy. E-mail: marco.roman@unive.it; Tel: +0039 041 234 7731

^bEuropean Synchrotron Radiation Facility (ESRF), 71 avenue des Martyrs, 38000 Grenoble, France

^cUniversity of Aberdeen, Trace Element Speciation Laboratory, Aberdeen AB24 3UE, Scotland, UK

^dUniversity of Graz, Institute of Chemistry, Universitätsplatz 1, 8010 Graz, Austria

^eUniversity Hospital of Padua, Burns Centre, Division of Plastic Surgery, Via Giustiniani 2, 35128 Padua, Italy

^fUniversity of Padua, Department of Biomedical Sciences, Via Ugo Bassi 58/B, 35131 Padua, Italy

^gEcamRicert Srl, European Centre for the Sustainable Impact of Nanotechnology (ECSIN), Corso Stati Uniti 4, 35127 Padua, Italy

^hInstitute for Polar Sciences (ISP-CNR), Via Torino 155, 30172 Venice Mestre, Italy



Fig. 3 Full profile biopsy specimen of the wound of patient A, 3 days after a single application of the AgNP-containing dressing, analysed by SR- μ XRF and SR-XANES. (A) Histological image of the tissue (adjacent slice). (B) Picture of the analysed slice. The scanned area locates in the white frame. (C) Maps of signal intensities (cps) of Ag (log scale), P, S and Cl (linear scales); pixel size 2 μ m. (D) Ternary RGB plot of the same elemental maps. The arrows indicate: *stratum corneum; **stratum spinosum; ***stratum basale; ****fragments of epidermis extending in the granulation tissue; [#]granulation tissue; ⁺reticular dermis; °hypodermis. (E) Zoom RGB map of the region located in the frame (independent acquisition, pixel size 2 μ m), the arrow indicates the region selected for SR- μ XANES analysis; (F) average SR- μ XRF spectrum of the whole zoom map area; (G) average SR- μ XANES spectrum in the selected region (dotted line) overlapped to its best LCF function (solid line).



Fig. 5 Full profile biopsy specimen of the wound of patient A, 12 days after a single application of the AgNP-containing dressing (complete healing), analysed by SR-µXRF. (A) Histological image of the tissue (adjacent slice), with zoom images of the regenerated epidermis (B) and an arterial blood vessel (C). (D) Picture of the analysed slice. The scanned area locates in the white frame. (E) Maps of signal intensities (cps) for Ag (logarithmic scale), P, S and Cl (linear scale); pixel size 2 µm. (F) Ternary RGB plot of the same elemental maps. The symbols indicate: *epidermis; **reticular dermis; **arterial vessel. (G) and (H): two consecutive zoom maps of the region located in the white frame, independent acquisitions with pixel size 1 µm and 0.5 µm, respectively. (I) Average µXRF spectrum of the whole zoom map area H.





The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.