

CORRECTION

View Article Online
View Journal | View Issue



Cite this: *Biomater. Sci.*, 2021, **9**, 1048

Correction: *In vivo* migration of Fe₃O₄@polydopamine nanoparticle-labeled mesenchymal stem cells to burn injury sites and their therapeutic effects in a rat model

Xiuying Li,^a Zhenhong Wei,^a Binxi Li,^b Jing Li,^a Huiying Lv,^a Liya Wu,^a Hao Zhang,^b Bai Yang,^b Mingji Zhu^{*a} and Jinlan Jiang^{*a}

DOI: 10.1039/d0bm90110e
rsc.li/biomaterials-science

Correction for '*In vivo* migration of Fe₃O₄@polydopamine nanoparticle-labeled mesenchymal stem cells to burn injury sites and their therapeutic effects in a rat model' by Xiuying Li *et al.*, *Biomater. Sci.*, 2019, **7**, 2861–2872, DOI: 10.1039/C9BM00242A.

The authors regret errors in Fig. 1c, 3a and 5e in the original article. The corrected figures are shown below. In Fig. 1c, the authors wish to use a different TEM image to the one in Fig. 2a in their previous Biomaterials Science paper.¹ In Fig. 3a, the 100, 150 and 200 $\mu\text{g ml}^{-1}$ panels have been replaced as the original panels were incorrect leading to partial overlap with the 50 $\mu\text{g ml}^{-1}$ panel. In Fig. 5e, the MSC and MSC + NPs bands for β -actin have been reversed as the original bands were incorrectly labelled due to being in the wrong orientation.

The raw data for Fig. 3a and 5e were provided by the authors and reviewed by an independent expert.




Fig. 1 Fe₃O₄@PDA nanoparticle preparation and internalization by MSCs. (C) The TEM image of a representative Fe₃O₄@PDA nanoparticle. Scale bar = 20 nm.

^aScientific Research Center, China-Japan Union Hospital of Jilin University, Changchun, Jilin, China. E-mail: jiangjinlan@jlu.edu.cn, zhumingji0822@qq63.com; Tel: +86 18186811856

^bState Key Laboratory of Supramolecular Structure and Materials, College of Chemistry, Jilin University, Changchun, Jilin, China






Fig. 3 Viability and proliferation potential of the $\text{Fe}_3\text{O}_4\text{@PDA}$ -labeled MSCs. (A) Photomicrographs showing the morphology of the MSCs labeled with the NPs at concentrations of 0, 25, 50, 100, 150, and 200 $\mu\text{g mL}^{-1}$. Scale bars = 100 μm .




Fig. 5 Effect of the $\text{Fe}_3\text{O}_4\text{@PDA}$ NPs on MSC migration *in vitro*. The migration of the MSCs labeled with 50 $\mu\text{g mL}^{-1}$ NPs for 16 h. (E) Protein expression of CXCR4 in the MSCs labeled with 50 $\mu\text{g mL}^{-1}$ NPs for 16 h, evaluated by western blotting.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

References

- 1 L. Wu, F. Zhang, Z. Wei, X. Li, H. Zhao, H. Lv, R. Ge, H. Ma, H. Zhang, B. Yang, J. Li and J. Jiang, *Biomater. Sci.*, 2018, **6**, 2714.

