




Showcasing research from Professor Maurizio Taddei's laboratory, Department of Biotechnology, Chemistry and Pharmacy, University of Siena and Dr Giuseppe Giannini's research group, Alfasigma S.p.A., Pomezia, Italy

Antibody drug conjugates with hydroxamic acid cargos for histone deacetylase (HDAC) inhibition

The antibody creeping along the streets of the medieval historical center of Siena is an ADC loaded with a hydroxamic acid. Once the target is reached, the conjugate releases its cargo and the hydroxamate drug can hit Histone Deacetylase. In this way it is possible to overcome the poor physicochemical properties of hydroxamic acids.

As featured in:



See Maurizio Taddei, Giuseppe Giannini *et al.*, *Chem. Commun.*, 2021, **57**, 867.



Cite this: *Chem. Commun.*, 2021, 57, 867

Received 10th September 2020,
Accepted 22nd November 2020

DOI: 10.1039/d0cc06131j

rsc.li/chemcomm

Antibody drug conjugates with hydroxamic acid cargos for histone deacetylase (HDAC) inhibition†

Claudio Cianferotti,^a Valentina Faltoni,^a Elena Cini,^a Elena Ermini,^a Francesca Migliorini,^a Elena Petricci,^a Maurizio Taddei,^a Laura Salvini,^b Gianfranco Battistuzzi,^c Ferdinando Maria Milazzo,^c Anna Maria Anastasi,^c Caterina Chiapparino,^c Rita De Santis^c and Giuseppe Giannini^{*,c}

Antitumor hydroxamates SAHA and Dacinostat have been linked to cetuximab and trastuzumab through a non-cleavable linker based on the *p*-mercaptobenzyl alcohol structure. These antibody drug conjugates (ADCs) were able to inhibit HDAC in several tumour cell lines. The cetuximab based ADCs block human lung adenocarcinoma cell proliferation, demonstrating that bioconjugation with antibodies is a suitable approach for targeted therapy based on hydroxamic acid-containing drugs. This work also shows that ADC-based delivery might be used to overcome the classical pharmacokinetic problems of hydroxamic acids.

Hydroxamic acids are a peculiar type of organic molecules featuring a large variety of biological activities. They are potent enzyme inhibitors binding lipoxygenase, hydroxylase, matrix-metalloproteinase, histone deacetylase, carbonic anhydrase, ribonucleotide reductase and several other enzyme families.¹ Besides this wide range of inhibition activities, hydroxamic acids have found therapeutic applications in cancer,² cardiovascular diseases, including hypertension,³ tuberculosis,⁴ HIV infection,⁵ and Alzheimer's disease.⁶ Most of their biological activity is related to the complexation of metals present in enzymes through interaction with *N*-hydroxy and carbonyl groups.⁷ They are excellent bidentate ligands for metals such as iron, nickel and zinc. However, translation of this exceptional potential to real drugs has been highly limited by several drawbacks such as mutagenicity, poor pharmacokinetics and several off-target interactions, resulting in ventricular repolarization impairment, thrombocytopenia, gastrointestinal toxicity and several other adverse side effects.⁸

Hydroxamic acids are privileged structures to block histone deacetylase (HDAC) enzymes, and HDAC inhibitors (HDACis)

have emerged as one of the most likely classes of multifunctional anticancer drugs.⁹ Cell cycle arrest, apoptosis, angiogenesis regulation, activation of tumour suppressor genes and oncogene suppression are some of the down-stream effects observed upon pharmacological treatment with HDACis.³

Although acting on the pharmacophore shape through the HDACis structure modification might improve the target selectivity,¹⁰ an attractive alternative would be redirecting the inhibitor to a suitable target using selective delivery. Few examples of hydroxamic acid pro-drug delivery systems have been described to date.¹¹

One of the most efficient methods for drug delivery is the antibody–drug conjugate (ADC) strategy, where the antibody drives the drug inside the cell expressing the antibody receptor.¹² Once internalized, the antibody–drug linker is cleaved, and the drug is released inside the malignant cells.¹³ The FDA has approved half a dozen ADCs, all of them carrying highly cytotoxic payloads like vedotin, emtansine and ozogamicin, and many others are under clinical trials.¹⁴ It is worth noting that trastuzumab emtansine (T-DM1, Kadcyla) is the only product approved for the treatment of solid tumours.

The current linker technology allows charging antibodies mostly with drugs carrying amines, alcohols or phenols through carbamates or carbonates, while the bioconjugation of drugs with other functional groups is still an undeveloped issue. It is not surprising that the only example of hydroxamic acid linked to antibodies deals with bifunctional chelates bearing hydroxamate arms for the radiometal labeling of monoclonal antibodies.¹⁵

Following our long-standing interest in HDAC inhibition,¹⁶ we report here the first example of ADC cargos carrying the FDA approved Vorinostat/SAHA (**1**) and Dacinostat/NVP-LAQ824 (**4**),¹⁷ a highly potent HDAC pan-inhibitor with IC₅₀ = 32 nM (Scheme 1). These products were conjugated with cetuximab (Ctx) or trastuzumab (Trast), and a preliminary study on their biological activity confirmed the efficient targeting of HDACs within the tumour cells.

Based on our experience with bioconjugation,¹⁸ we started to explore the possibility to link SAHA with mAbs through non-cleavable linkers. Many non-cleavable linkers are based

^a Dipartimento di Biotecnologie, Chimica e Farmacia, Università di Siena, Via A. Moro 2, 53100 Siena, Italy. E-mail: maurizio.taddei@unisi.it

^b Fondazione Toscana Life Science, Via Fiorentina 1, 53100, Siena, Italy

^c R&D Alfasigma S.p.A., Via Pontina, Km. 30.400, 00071 Pomezia, Roma, Italy. E-mail: Giuseppe.Giannini@alfasigma.com

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0cc06131j




13 X = OH
 $\text{14 X = Br} \xleftarrow{\text{PBr}_3}$

$\xrightarrow{\text{SAHA (1), NaOH, H}_2\text{O}_2, \text{MeOH, rt}}$ 46%

$\text{15 R = -C}_2\text{H}_5$ 16 R = H

The synthesis of a Cathepsin B cleavable linker **16** started with the amide **13** composed by dipeptide Fmoc-Val-Cit-OH and *p*-amino benzyl alcohol (PABA) (Scheme 3). Compound **13** was converted into the corresponding bromide **14** with PBr₃.¹⁸

Results show that all the ADCs are internalized after binding to EGFR or ErbB2 receptors depending on the target receptor expression level. Internalization was equal with respect to the



Chemical structures of three antibody-drug conjugates (ADCs) are shown:

- Cbx:17 Trast:18**: The structure shows an antibody (Y-shaped) linked to a drug molecule via a linker. The linker contains a triazole ring and a thioether-linked drug moiety.
- Cbx:19**: The structure shows an antibody (Y-shaped) linked to a drug molecule via a linker. The linker contains a triazole ring and a thioether-linked drug moiety with a hydroxyl group.
- Trast:20**: The structure shows an antibody (Y-shaped) linked to a drug molecule via a linker. The linker contains a triazole ring and a thioether-linked drug moiety with a hydroxyl group.

This journal is © The Royal Society of Chemistry 2021

unconjugated Ctx or Trast, resulting in accumulation within the internal vesicles of the multivesicular bodies and subsequent translocation to the lysosomal compartment (Fig. S6, ESI†).

Release of hydroxamic acids **1** and **4** occurred upon treatment of ADCs **17–20** with human hepatic microsomes. After 72 h of incubation at 37 °C, quantitative HPLC/HRMS analysis showed the presence of peaks at *m/z* 264.1472 and 349.1792, respectively, attributed exclusively to compounds **1** and **4** in concentrations corresponding to more than 70% of release of the payload present in the starting ADC.²⁰




Fig. 1 Effect of SAHA, Dacinostat, Cet and Trast, compared to ADCs **17–20**, on the acetylation of α -tubulin (a) and histone H3 (b) in A549 cells. Cells cultivated for 3 h with reference drugs (100 nM) or antibodies (5 $\mu\text{g mL}^{-1}$) and stained with anti-acetylated- α -tubulin (red) or anti-acetylated-H3 histone (pink). Insets show the Draq5 dye-stained nucleus and cytoplasm. Fluorescence imaging by HCS Operetta. Data not normalized. Each image is representative of at least 5 fields of duplicate wells. Magnification 60 \times .

Inhibition of the HDAC activity was determined through HCS-imaging analysis, monitoring the levels of acetylated histone H3 and acetylated α -tubulin in A549 (human lung adenocarcinoma) cells, as the indicators of specific inhibition of nuclear class I HDACs and of cytoplasmic HDAC6, respectively (Fig. 1). Results showed that ADCs **17**, **19** and **20** induced a relevant increase in the acetylation level of α -tubulin and histone H3, as a consequence of direct enzymatic inhibition, while no significant effect was noted in cells treated with Ctx or Trast alone (Fig. 1). Neither cetuximab nor trastuzumab is endowed with HDAC inhibitor activity. So, an increase in the acetylation degree of α -tubulin- or histone H3 is not expected in the cells treated with these drugs. In fact, the extent of acetylation observed in A549 cells following treatment with the above drugs was observed to be similar or apparently a little lower (depending on the fields observed) with respect to the basal level of untreated cells (as shown in Fig. 1). ADC **18** induced an increased acetylation of both α -tubulin and histone H3, although with lower potency with respect to the others. In these cells, the cetuximab-based ADCs **17** and **19** would seem to act with higher potency, probably due to the highest expression of EGFR as compared to ErbB2 receptors on the cell surface. Increased acetylation of histone H3 and α -tubulin was detected through western blot analysis of the total protein lysate. The results of the densitometric analysis of the specific band intensity, after normalization to β -actin signals, showed that the acetylation enhanced upon treatment of tumour cells with ADCs as compared to the vehicle-treated cells and cells treated with Ctx and Trast alone (Fig. S7, ESI†). These results demonstrate that the ADCs are efficiently internalized by the tumour cells releasing the hydroxamic acids required for epigenetic modulation. Similar results were observed on other human tumor (pancreas, melanoma, and breast) cell lines (data not shown).

The anti-proliferative activity of the conjugates **17** and **19** was evaluated in A549 cells upon the treatment, for 6 days, with several doses of ADCs and with equivalent doses of Ctx compared to the unconjugated SAHA and Dacinostat (Fig. 2). ADC **19** inhibited cell proliferation very efficiently in a dose-dependent manner and with quite a similar potency ($\text{IC}_{50} = 90 \pm 3.4 \text{ nM}$) to




Fig. 2 Antiproliferative activity of ADCs **17** and **19** on A549 cells upon 6 days of treatment. Data are the mean (\pm SD) of percentage inhibition ($n = 5$).



that of the reference Dacinostat ($IC_{50} = 33 \pm 1.7$ nM).²¹ ADC 17 ($IC_{50} = 1440 \pm 17$ nM) was less efficacious than ADC 19.²² Interestingly, ADC 17 retains a certain percentage of activity in a range of concentrations (100–500 nM) in which SAHA resulted utterly inactive.

In conclusion, we have developed a new class of ADCs for the targeted delivery of hydroxamic acids. The new linkers based on PMBA can release the hydroxamic acid after metabolic degradation demonstrating that bioconjugation with mAbs can be employed to develop more effective targeted therapy based on hydroxamate HDACis.²³

This work was supported in part by the AIRC (under IG 2017 – ID. 20758 project – P.I. Taddei Maurizio) and by the project “Development and application of QM/MM technologies for the design of light responsive proteins or protein-mimics based on rhodopsin architecture” within the program “Dipartimenti di Eccellenza – 2018–2022 MIUR (Rome, Italy).

Conflicts of interest

There are no conflicts to declare.

Notes and references

- (a) S. P. Gupta, *Chem. Rev.*, 2015, **115**, 6427–6490; (b) E. M. F. Muri, M. J. Nieto, R. D. Sindelar and J. S. Williamson, *Curr. Med. Chem.*, 2002, **9**, 1631–1653.
- M. Manal, M. J. N. Chandrasekar, J. Gomathi Priya and M. J. Nanjan, *Bioorg. Chem.*, 2016, **67**, 18–42.
- R. Sangwan, R. Rajan and P. K. Mandal, *Eur. J. Med. Chem.*, 2018, **158**, 620–706.
- T. van den Bosch, M. Kwiatkowski, R. Bischoff and F. J. Dekker, *Epigenomics*, 2017, **9**, 1013–1028.
- R. Badia, J. Grau, E. Riveira-Muñoz, E. Ballana, G. Giannini and J. A. Esté, *Antiviral Res.*, 2015, **123**, 62–69.
- P. Narayan and M. Dragunow, *Adv. Exp. Med. Biol.*, 2017, **978**, 321–336.
- J. M. Roosenberg, Y. M. Lin, Y. Lu and M. J. Miller, *Curr. Med. Chem.*, 2000, **7**, 159–197.
- (a) A. Rambaldi, C. M. Dellacasa, G. Finazzi, A. Carobbio, M. L. Ferrari, P. Guglielmelli, E. Gattoni, S. Salmoiraghi, M. C. Finazzi, S. Di Tollo, C. D'Urzo, A. M. Vannucchi, G. Barosi and T. Barbui, *Br. J. Haematol.*, 2010, **150**, 446–455; (b) S. Subramanian, S. E. Bates, J. J. Wright, I. Espinoza-Delgado and R. L. Piekarz, *Pharmaceuticals*, 2010, **3**, 2751–2767.
- (a) Y. Sun, Y. Sun, S. Yue, Y. Wang and F. Lu, *Curr. Top. Med. Chem.*, 2019, **18**, 2420–2428; (b) C. M. Grunewald, W. A. Schulz, M. A. Skowron, M. J. Hoffmann and G. Niegisch, *Transl. Cancer Res.*, 2018, **7**, 1151–1160.
- A. V. Bieliauskas and M. K. H. Pflum, *Chem. Soc. Rev.*, 2008, **37**, 1402.
- (a) W. Fan, L. Zhang, Q. Jiang, W. Song, F. Yan and L. Zhang, *Eur. J. Med. Chem.*, 2020, **203**, 112628; (b) K. Zuwala, A. A. A. Smith, M. Tolstrup and A. N. Zelikin, *Chem. Sci.*, 2016, **7**, 2353–2358; (c) W. Szymanski, M. E. Ourailidou, W. A. Velema, F. J. Dekker and B. L. Feringa, *Chem. – Eur. J.*, 2015, **21**, 16517–16524; (d) S. A. Senevirathne, K. E. Washington, J. B. Miller, M. C. Biewer, D. Oupicky, D. J. Siegwart and M. C. Stefan, *J. Mater. Chem. B*, 2017, **5**, 2106–2114; (e) P. Šilhár, L. M. Eubanks, H. Seki, S. Pellett, S. Javor, W. H. Tepp, E. A. Johnson and K. D. Janda, *J. Med. Chem.*, 2013, **56**, 7870–7879.
- V. Chudasama, A. Maruani and S. Caddick, *Nat. Chem.*, 2016, **8**, 114–119.
- (a) M. R. Gordon, M. Canakci, L. Li, J. Zhuang, B. Osborne and S. Thayumanavan, *Bioconjugate Chem.*, 2015, **26**, 2198–2215; (b) R. V. J. Chari, M. L. Miller and W. C. Widdison, *Angew. Chem., Int. Ed.*, 2014, **53**, 3796–3827.
- K. Tsuchikama and Z. An, *Protein Cell*, 2018, **9**, 33–46.
- (a) S. Ait-Mohand, C. Denis, G. Tremblay, M. Paquette and B. Guérin, *Org. Lett.*, 2014, **16**, 4512–4515; (b) A. Safavy, M. B. Khazaeli, M. Kirk, L. Coward and D. J. Buchsbaum, *Bioconjugate Chem.*, 1999, **10**, 18–23.
- (a) G. Giannini, L. Vesci, G. Battistuzzi, D. Vignola, F. M. Milazzo, M. B. Guglielmi, M. Barbarino, M. Santaniello, N. Fanto, M. Mor, S. Rivara, D. Pala, M. Taddei, C. Pisano and W. Cabri, *J. Med. Chem.*, 2014, **57**, 8358–8377; (b) M. Taddei, S. Ferrini, L. Giannotti, M. Corsi, F. Manetti, G. Giannini, L. Vesci, F. M. Milazzo, D. Alloatti, M. B. Guglielmi, M. Castorina, M. L. Cervoni, M. Barbarino, R. Foderà, V. Carollo, C. Pisano, S. Armaroli and W. Cabri, *J. Med. Chem.*, 2014, **57**, 2258–2274; (c) C. B. Botta, W. Cabri, E. Cini, L. De Cesare, C. Fattorusso, G. Giannini, M. Persico, A. Petrella, F. Rondinelli, M. Rodriguez, A. Russo and M. Taddei, *J. Med. Chem.*, 2011, **54**, 2165–2182; (d) M. Rodriguez, S. Terracciano, E. Cini, G. Settembrini, I. Bruno, G. Bifulco, M. Taddei and L. Gomez-Paloma, *Angew. Chem., Int. Ed.*, 2006, **45**, 423–427.
- S. Remiszewski, *Curr. Med. Chem.*, 2003, **10**, 2393–2402.
- E. Cini, V. Faltoni, E. Petricci, M. Taddei, L. Salvini, G. Giannini, L. Vesci, F. M. Milazzo, A. M. Anastasi, G. Battistuzzi and R. De Santis, *Chem. Sci.*, 2018, **9**, 6490–6496.
- (a) H. Wu, P. J. Levalley, T. Luo, A. M. Kloxin and K. L. Kiick, *Bioconjugate Chem.*, 2018, **29**, 3595–3605; (b) A. D. Baldwin and K. L. Kiick, *Bioconjugate Chem.*, 2011, **22**, 1946–1953.
- A. J. Bessire, T. E. Ballard, M. Charati, J. Cohen, M. Green, M. H. Lam, F. Loganzo, B. Nolting, B. Pierce, S. Puthenveetil, L. Roberts, K. Schildknecht and C. Subramanyam, *Bioconjugate Chem.*, 2016, **27**, 1645–1654.
- T. Beckers, C. Burkhardt, H. Wieland, P. Gimmnich, T. Ciossek, T. Maier and K. Sanders, *Int. J. Cancer*, 2007, **121**, 1138–1148.
- P. Agarwal and C. R. Bertozzi, *Bioconjugate Chem.*, 2015, **26**, 176–192.
- S. Shen and A. P. Kozikowski, *ChemMedChem*, 2016, **11**, 15–21.

