


 Cite this: *Chem. Commun.*, 2021, 57, 5694

 Received 20th April 2021,  
 Accepted 5th May 2021

DOI: 10.1039/d1cc02097h

[rsc.li/chemcomm](http://rsc.li/chemcomm)

## A new mode of cyclobutenedione ring opening for the synthesis of 2-oxobut-3-enamides and tetrasubstituted furans†

 Ryan M. Bennett, Wei Sun, Dharyl C. Wilson, Mark E. Light and David C. Harrowven \*

**A dichotomy between the additions of organolithiums and lithium amides to cyclobutenediones is described wherein the former give carbonyl addition products while the latter induce ring opening by enone cleavage via O- to C-lithium transfer. This distinct mode of ring scission gives access to 2-oxobut-3-enamides and tetrasubstituted furans.**

Cyclobutenediones and squarates, *e.g.* **1a**, are used extensively as precursors to carbocyclic and heterocyclic ring systems through ring expansion (Scheme 1).<sup>1–10</sup> Typically, a carbon nucleophile is first added to give an adduct **2** that rearranges to a product on thermolysis or photolysis. Rearrangements triggered by light usually give rise to 5*H*-furanones **6** *via* vinylketene (*E*)-**4**,<sup>9,10</sup> while those triggered by heat typically proceed *via* the isomeric vinylketene (*Z*)-**4** and give products determined by the nature of the residue introduced at C4. The versatility and reliability of the chemistry is evident from the frequent deployment of such rearrangements in natural products total synthesis.<sup>5,8</sup>

Herein we describe a dichotomy between the additions of organo-lithiums and lithium amides to cyclobutenediones (Scheme 1). Thus, while dimethyl squarate **1a** gives cyclobutenones **2** on treatment with organolithium reagents its reactions with lithium amides lead to vinylolithium intermediates **5** *via* the corresponding adduct **3**. Herein we show how this unprecedented mode of ring opening, involving scission of the C3–C4 bond with concomitant O- to C-lithium transfer, provides access to an array of 2-oxo-but-3-enamides **7** and tetrasubstituted furans **8**.

The discovery was made while preparing a series of 3-amino-4-methoxycyclobutenediones by the addition of amines to dimethyl squarate **1a** (Scheme 2).<sup>9</sup> Though the method worked

well for many substrates, *e.g.* **1a** to **9a–c**, it returned starting materials when applied to 2°-amines with a high steric burden, *e.g.* <sup>1</sup>Pr<sub>2</sub>NH.<sup>11</sup> This prompted a switch to using LDA as the nucleophile, but instead of delivering the anticipated 3-aminocyclobutenedione **9d**, 2-oxo-but-3-enamide **7a** was isolated as the major product in 65% yield. Intrigued by this finding, we decided to explore the generality of the reaction, beginning with extensions to diisopropyl and di-*tert*-butyl squarates, **1b** and **1c**. Pleasingly, both gave the corresponding 2-oxo-but-3-enamides, **7b** and **7c**, albeit more slowly due to the increase in steric demand.

The reaction was next extended to 2-alkoxy-3-aminocyclobuten-ones **9**. All of the cases examined (Table 1) showed excellent regioselectivity leading to the corresponding vinyllogous amides **11** exclusively. This outcome can be attributed to a preferred addition of LDA to the vinyllogous ester carbonyl in **9**, leading to adduct **10**, over its addition to the vinyllogous amide carbonyl.

Steric influences on the reaction were next examined with respect to the lithium amide (Scheme 3). Notably, while LDA and lithium piperidide each induced ring opening of cyclobutenedione **9a** to 2-oxo-but-3-enamides **11a** and **12a** respectively, starting material was returned when bulkier lithium amides were employed, *e.g.* LiHMDS, lithium 2-methylpiperidide and LiTMP. By way of contrast, lowering steric demand with lithium dimethylamide promoted substitution of the alkoxide leading to the bis-aminocyclobutenedione **13b**.

Lithium amide additions to dimethyl squarate **1a** could also be sequenced with substitution of the terminal alkoxide for NH<sub>2</sub> through employment of an ammoniacal work up (Scheme 4).<sup>12</sup>


Reactions of 3-alkoxy-4-alkylcyclobutenediones **16** with LDA and lithium piperidide were next examined and exposed further subtleties (Scheme 5). Thus, while treatment of the *tert*-butyl derivative **16a** with LDA gave the anticipated oxobutenamide **18a**, substitution dominated with lithium piperidide leading to amino-cyclobutenedione **17**. Reducing steric demand with the methyl analogue **16b** led to cleaner reactions and higher yields, with lithium piperidide giving oxobutenamide **18b** in 79% yield.


Chemistry, University of Southampton, Highfield, Southampton, SO17 1BJ, UK.

E-mail: dch2@soton.ac.uk


† Electronic supplementary information (ESI) available: Experimental accounts with spectral details and copies of NMR spectra are available as supplementary information. CCDC 2059384, 2042398 and 1500038. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1cc02097h








**Scheme 5** Reactions of 3-alkoxy-4-alkylcyclobutenediones with lithium piperide and LDA.




**Fig. 1** Summary of DFT calculations on the opening of dimethyl squarate **1a** with lithium dimethylamide.




**Fig. 2** Summary of DFT calculations on the opening of cyclobutenedione **9a** with lithium dimethylamide.

was treated with benzylamine, where 2-oxo-but-3-enamide **27** (Fig. 3) was as a significant byproduct formed in 25% yield alongside with the anticipated aminocyclobutenedione **26**




**Fig. 3** X-ray crystal structure of 2-oxo-but-3-enamide **27** [CCDC# 1500038].



**Scheme 6** A related fragmentation of cyclobutenedione **25** triggered with benzylamine.

**Table 2** Three-component coupling reactions leading to tetrasubstituted furans



(40% yield). To the best of our knowledge, this side reaction has not been reported previously (Scheme 6).

In conclusion, we have uncovered a new mode of cyclobutenedione ring opening that is triggered by lithium amide addition. The reaction proceeds *via* sequential *N*- to *O*- to *C*-lithium transfer, as detailed in Scheme 1, and gives access to an array of 2-oxo-but-3-enamides (*via* protonation) and tetrasubstituted furans (*via* aldehyde addition). We are currently investigating further extensions of the methodology in heterocyclic synthesis.

Dr Wei Sun and Ryan Bennett contributed equally in respect of the experimental work with Dr Mark Light performing the X-ray analyses and Prof. David Harrowven supervising the work.

We gratefully acknowledge financial support from the European Regional Development Fund [ERDF Interreg Va programme (Project 121)] and EPSRC [EP/P013341/1, EP/L003325/1 and EP/K039466/1].




## Conflicts of interest

There are no conflicts of interest to declare.

## Notes and references

- H. W. Moore and O. H. W. Decker, *Chem. Rev.*, 1986, **86**, 821; L. S. Liebeskind, *Tetrahedron*, 1989, **45**, 3053.
- H. W. Moore and S. T. Perri, *J. Org. Chem.*, 1988, **53**, 996; J. M. Heerding and H. W. Moore, *J. Org. Chem.*, 1991, **56**, 4048; L. S. Liebeskind and J. Zhang, *J. Org. Chem.*, 1991, **56**, 6379; L. M. Mayo, M. P. Winters and H. W. Moore, *J. Org. Chem.*, 1992, **57**, 6896; D. J. Krysan, A. Gurski and L. S. Liebeskind, *J. Am. Chem. Soc.*, 1992, **114**, 1412; P. Turnbull, M. J. Heileman and H. W. Moore, *J. Org. Chem.*, 1996, **61**, 2584; B. Yucel, B. Sanli, H. Soylemez and I. Yilmaz, *Tetrahedron*, 2011, **67**, 1406; E. Packard, D. D. Pascoe, J. Maddaluno, T. P. Goncalves and D. C. Harrowven, *Angew. Chem., Int. Ed.*, 2013, **52**, 13076; Y. Yamamoto, T. Kurohara and M. Shibuya, *Chem. Commun.*, 2015, **51**, 16357; J. T. Hill-Cousins, I.-A. Pop, G. Pileio, G. Stevanato, P. Håkansson, S. S. Roy, M. H. Levitt, L. J. Brown and R. C. D. Brown, *Org. Lett.*, 2015, **17**, 2150.
- L. D. Foland, J. O. Karlsson, S. T. Perri, R. Schwabe, S. L. Xu, S. Patil and H. W. Moore, *J. Am. Chem. Soc.*, 1989, **111**, 975; K. Karabelas and H. W. Moore, *J. Am. Chem. Soc.*, 1990, **112**, 5372; C. Boullais, C. Rannou, E. Reveillere and C. Mioskowski, *Eur. J. Org. Chem.*, 2000, 723; T. P. Goncalves, M. Mohamed, R. J. Whitby, H. F. Sneddon and D. C. Harrowven, *Angew. Chem., Int. Ed.*, 2015, **54**, 4531; W. Sun, D. C. Wilson, M. E. Light and D. C. Harrowven, *Org. Lett.*, 2018, **20**, 4346.
- L. S. Liebeskind, S. Iyer and C. F. Jewell, *J. Org. Chem.*, 1986, **51**, 3065; L. S. Liebeskind and J. Zhang, *J. Org. Chem.*, 1991, **56**, 6379; L. S. Liebeskind, K. L. Granberg and J. Zhang, *J. Org. Chem.*, 1998, **63**, 2835; T. Kurohara, J. Jiyue, M. Shibuya and Y. Yamamoto, *Synthesis*, 2018, 1687.
- A. G. Birchler, F. Liu and L. S. Liebeskind, *J. Org. Chem.*, 1994, **59**, 7737; M. Mohamed, T. P. Goncalves, R. J. Whitby, H. F. Sneddon and D. C. Harrowven, *Chem. – Eur. J.*, 2011, **17**, 13698.
- S. L. Xu and H. W. Moore, *J. Org. Chem.*, 1989, **54**, 6018; S. L. Xu, H. Xia and H. W. Moore, *J. Org. Chem.*, 1991, **56**, 6094; Y. Yamamoto, M. Ohno and S. Eguchi, *Bull. Chem. Soc. Jpn.*, 1996, **69**, 1353; J. M. MacDougall and H. W. Moore, *J. Org. Chem.*, 1999, **64**, 7445; S. K. Verma, Q. H. Nguyen, J. M. MacDougall, E. B. Fleischer and H. W. Moore, *J. Org. Chem.*, 2000, **65**, 3379; T. Kurohara, J. Jiyue, M. Shibuya and Y. Yamamoto, *Synthesis*, 2018, 1687; M. A. Manning, W. Sun, M. E. Light and D. C. Harrowven, *Chem. Commun.*, 2021, 57, 4556.
- K. H. Lee and H. W. Moore, *J. Org. Chem.*, 1995, **60**, 735; F. Liu and L. S. Liebeskind, *J. Org. Chem.*, 1998, **63**, 2835.
- S. T. Perri, H. J. Dyke and H. W. Moore, *J. Org. Chem.*, 1989, **54**, 2032; L. D. Foland, O. H. W. Decker and H. W. Moore, *J. Am. Chem. Soc.*, 1989, **111**, 989; M. J. Heileman, R. Tiedemann and H. W. Moore, *J. Am. Chem. Soc.*, 1998, **120**, 3801; A. Enhsen, K. Karabelas, J. M. Heerding and H. W. Moore, *J. Org. Chem.*, 1990, **55**, 1177; S. T. Perri and H. W. Moore, *J. Am. Chem. Soc.*, 1990, **112**, 1897; V. J. Santora and H. W. Moore, *J. Am. Chem. Soc.*, 1995, **117**, 8486; C. S. Tomooka, H. Liu and H. W. Moore, *J. Org. Chem.*, 1996, **61**, 6009; Y. Xiong and H. W. Moore, *J. Org. Chem.*, 1996, **61**, 9168; J. M. MacDougall, V. J. Santora, S. K. Verma, P. Turnbull, C. R. Hernandez and H. W. Moore, *J. Org. Chem.*, 1998, **63**, 6905; D. C. Harrowven, D. D. Pascoe, D. Demurtas and H. O. Bourne, *Angew. Chem., Int. Ed.*, 2005, **44**, 1221; D. Knueppel and S. F. Martin, *Angew. Chem., Int. Ed.*, 2009, **48**, 2569; X. Yan and X. Hu, *J. Org. Chem.*, 2014, **79**, 5282; D. Knueppel, J. Yang, B. Cheng, D. Mans and S. F. Martin, *Tetrahedron*, 2015, **71**, 5741; D. J. Asby, M. G. Radigois, D. C. Wilson, F. Cuda, C. L. L. Chai, A. Chen, A. S. Bienemann, M. E. Light, D. C. Harrowven and A. Tavassoli, *Org. Biomol. Chem.*, 2016, **4**, 9330; J. Feng, X. Lei, Z. Guo and Y. Tang, *Angew. Chem., Int. Ed.*, 2017, **56**, 7895.
- W. Sun, D. C. Wilson and D. C. Harrowven, *Synthesis*, 2017, 3091.
- S. T. Perri, L. D. Foland and H. W. Moore, *Tetrahedron Lett.*, 1988, **29**, 3529; Y. Yamamoto, M. Ohno and S. Eguchi, *Tetrahedron*, 1994, **50**, 7783; D. C. Harrowven, M. Mohamed, T. P. Goncalves, R. J. Whitby, D. Bolien and H. F. Sneddon, *Angew. Chem., Int. Ed.*, 2012, **51**, 4405. See also D. E. Collin, E. H. Jackman, N. Jouandon, W. Sun, M. E. Light, D. C. Harrowven and B. Linclau, *Synthesis*, 2021, 1307.
- C. P. Krimmel, *US Pat.*, 4036828, 1977; N. C. Lim, M. D. Morton, H. A. Jenkins and C. Brueckner, *J. Org. Chem.*, 2003, **68**, 9233; A. Crew, A.-H. Li, L. Qiu, A. L. Castelhan, H. Dong, A. Smith, L. Tardibono and T. Zhang, *WO Pat.*, 34111, 2006; J. J. Baldwin, J. Yuan, B. Lawhorn, C. A. Leach, L. S. Kallander, P. T. Flaherty, S. B. Singh, R. D. Simpson, W. Zhao, G. Mcgeehan, Z. Xu, S. Semus, X. Hou, D. Ghirlanda, J. Zhang, B. Ghavini-Alagha, L. R. Terrell, Q. Lu, A. V. Ishchenko, D. A. Claremon, C. M. Tice, S. Cacatian and L. W. Dillard, *WO Pat.*, 117482, 2007; J. Chen, Y. Chen, A. Regueiro-Ren, S.-Y. Sit and J. Swidorski, *WO Pat.*, 2847, 2018; K. E. Benenato, E. S. Kumarasinghe and M. Cornebise, *WO Pat.*, 170306, 2018; Z. Wang, N. Li, J. Ma and Y. Shao, *WO Pat.*, 120296, 2019; N. Yu, *WO Pat.*, 160054, 2020; Z. Wang, J. Ma, Y. Shao and N. Li, *WO Pat.*, 160707, 2020.
- W. A. Kinney, N. E. Lee, D. T. Garrison, E. J. Podlesny, Jr., J. T. Simmonds, D. Bramlett, R. R. Notvest, D. M. Kowal and R. P. Tasse, *J. Med. Chem.*, 1992, **35**, 4720; A. P. Davis, S. M. Draper, G. Dunne and P. Ashton, *Chem. Commun.*, 1999, 2265; Y. Brandy, N. Brandy, E. Akinboye, M. Lewis, C. Mouamba, S. Mack, R. J. Butcher, A. J. Anderson and O. Bakare, *Molecules*, 2013, **18**, 1973; N. Sim, S. Gottschalk, R. Pal, J. Engelmann, D. Parker and A. Mishra, *Chem. Sci.*, 2013, **4**, 3148; M. Lu, Q.-B. Lu and J. F. Honek, *Bioorg. Med. Chem. Lett.*, 2017, **27**, 282.
- Reduction is presumed to occur by a mechanism akin to the Meerwein-Ponndorf-Verley reduction, *i.e.*



- R. N. McDonald, R. N. Steppe and R. C. Cousins, *J. Org. Chem.*, 1975, **40**, 1694; C. Kowalski, X. Creary, A. J. Rollin and M. C. Burke, *J. Org. Chem.*, 1978, **43**, 2601; M. Majewski and D. M. Gleave, *J. Organomet. Chem.*, 1994, **470**, 1.
- M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, N. J. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and J. J. Fox, *Gaussian 09, Revision C.01*, Gaussian, Inc., Wallingford CT, 2009.
  - P. X. Iten, A. A. Hofmann and C. H. Eugster, *Helv. Chim. Acta*, 1978, **61**, 430; C. H. Eugster, M. Balmer, R. Prewo and J. H. Bieri, *Helv. Chim. Acta*, 1981, **64**, 2636; M. E. Jung and L. J. Street, *J. Am. Chem. Soc.*, 1984, **106**, 8327; L. L. Klein and M. S. Shanklin, *J. Org. Chem.*, 1988, **53**, 5202; M. E. Jung and C. S. Siedem, *J. Am. Chem. Soc.*, 1993, **115**, 3822; G. H. Hakimelahi, M. L. Jain, T. W. Ly, I.-C. Chen, K. S. Ethiraj, J. R. Hwu and A. A. Moshfegh, *J. Org. Chem.*, 2001, **66**, 7067; M. Romero, P. Renard, D.-H. Caignard, G. Atassi, X. Solans, P. Constans, C. Bailly and M. D. Pujol, *J. Med. Chem.*, 2007, **50**, 294; M.-A. Légaré, E. Rochette, J. L. Lavergne, N. Bouchard and F.-G. Fontaine, *Chem. Commun.*, 2016, **52**, 5387; L. Grau, M. Romero, C. Privat-Contreras, D. Presa, M. Viñas, J. Morral, K. Pors, J. Rubio-Martinez and M. D. Pujol, *Eur. J. Med. Chem.*, 2020, **185**, 111807.

