

# ChemComm

Chemical Communications

rsc.li/chemcomm



ISSN 1359-7345



# Synthesis and reactivity of heteroleptic zinc(I) complexes toward heteroallenes†

 Bin Li,  Kevin Huse, Christoph Wölper and Stephan Schulz \*

 Cite this: *Chem. Commun.*, 2021, 57, 13692

 Received 5th October 2021,  
 Accepted 23rd November 2021

DOI: 10.1039/d1cc05617d

rsc.li/chemcomm

**Heteroleptic zinc(I) complexes**  $L^1Zn-ZnCp^*$  ( $L^1 = HC[C(CF_3)NC_6F_5]_2$  **1**;  $L^2 = HC[C(Me)NDipp]_2$ ; Dipp = 2,6-*i*-Pr<sub>2</sub>C<sub>6</sub>H<sub>3</sub> **2**) are synthesized by reactions of  $Cp^*_2Zn_2$  with  $L^1H$  and  $L^2ZnH$ . **2** reacts with *t*-BuNCO to give unprecedented carbamate complex (**4**), while reactions with  $RN_3$  gave bis-hexazene, triazenide, and trimeric azide complexes (**5–7**).

Since the discovery of  $Cp^*_2Zn_2$  ( $Cp^* = C_5Me_5$ ) by Carmona *et al.*,<sup>1</sup> Zn(I) complexes have received increasing interest.<sup>2</sup> In contrast to  $Cp^*_2Zn_2$ , which is accessible by reactions of  $Cp^*_2Zn$  with  $(C_2H_5)_2Zn$  and by reduction of  $Cp^*_2Zn$  and  $ZnCl_2$  with  $KH$ ,<sup>3</sup> the majority of Zn(I) complexes are synthesized by Wurtz-type coupling of organozinc halides  $RZnX$ .<sup>4</sup> Such reactions also yield homotrimeric  $Zn_3$ <sup>5</sup> and heterotetranuclear  $Re_2Zn_2$  complexes,<sup>6</sup> but this pathway typically suffers from low yields. In contrast, ligand exchange reactions of  $Cp^*_2Zn_2$  with H-acidic ligands or potassium salts are more efficient routes for the synthesis of heteroleptic Zn(I) complexes.<sup>7–9</sup> Apart from their bonding nature, Zn(I) complexes are of interest due to their widespread reactivity, *i.e.* disproportionation,<sup>1</sup> acid–base,<sup>10</sup> protonation,<sup>11</sup> redox,<sup>12</sup> and cluster formation reactions,<sup>13</sup> respectively. Recently, we reported an isocyanate insertion reaction into one Zn–Cp\* bond of  $Cp^*_2Zn_2$ ,<sup>14</sup> revealing a new reaction pattern for Zn(I) complexes. Encouraged by this finding, we became interested in reactions of heteroleptic Zn(I) complexes  $Cp^*ZnZnL$  with heteroallenes, and report herein on the synthesis of  $L^1ZnZnCp^*$  **1** and  $L^2ZnZnCp^*$  **2**, and reactions of **2** with *t*-BuNCO and three organoazides  $RN_3$ .

The reaction of  $L^3H$  ( $L^3 = HC[C(Me)NMe]_2$ , Mes = 2,4,6-Me<sub>3</sub>C<sub>6</sub>H<sub>2</sub>) with  $Cp^*_2Zn_2$  in 1 : 1 and 1 : 2 molar ratio yielded the


homoleptic Zn(I) complex  $L^3_2Zn_2$ .<sup>7</sup> In contrast, the reaction of  $Cp^*_2Zn_2$  with one or two equiv. of  $L^1H$  ( $L^1 = HC[C(CF_3)NC_6F_5]_2$ ) in toluene at 6 °C gave the heteroleptic complex  $L^1ZnZnCp^*$  **1** (Scheme 1), whereas  $L^2H$  ( $L^2 = HC[C(Me)NDipp]_2$ , Dipp = 2,6-*i*-Pr<sub>2</sub>C<sub>6</sub>H<sub>3</sub>) failed to react. Inspired by a Pd(II)-induced homo-coupling reaction of  $RZnH$ ,<sup>15</sup>  $L^2ZnH$  was reacted with  $Cp^*_2Zn$  at 4 °C for 3 days, yielding  $L^2Zn-ZnCp^*$  (**2**) and other by-products (Fig. S37, ESI†). **2** also formed in the equimolar reaction of  $L^2ZnH$  with  $Cp^*_2Zn_2$  (Scheme 1), whereas substitution of the second Cp\* group by reacting **2** or  $Cp^*_2Zn_2$  with  $L^2ZnH$  in 1 : 1 or 1 : 2 molar ratio failed, although the desired homoleptic Zn(I) complex  $L^2_2Zn_2$  is known.<sup>4</sup> In contrast,  $Cp^*_2Zn_2$  reacted with the stronger reductant  $L^2MgH$  to give zinc metal and  $L^2MgCp^*$  (**3**, ESI†).

The <sup>1</sup>H and <sup>13</sup>C NMR spectra of **1** and **2** show characteristic resonances of the Cp\* group and  $L^1$  (**1**) and  $L^2$  (**2**). The <sup>19</sup>F NMR spectrum of **1** shows a resonance at –66.40 ppm (CF<sub>3</sub>) and three resonances of the C<sub>6</sub>F<sub>5</sub> groups in a 2 : 1 : 2 intensity ratio.

Single crystals were grown from toluene solutions at –30 °C (**1**, Fig. S42, ESI†) and 4 °C (**2**, Fig. S43, ESI†). The complexes crystallize in the triclinic space group  $P\bar{1}$  (**1**) and the monoclinic space group  $P2_1$  (**2**). The Zn1–Zn2 bond of **1** (2.2883(5) Å) is shorter than that of **2** (2.3008(2) Å) and other Zn(I) complexes, but comparable to those in  $[Zn]_8$  clusters (2.27–2.29 Å).<sup>16</sup> The Cp\* groups are η<sup>5</sup>-bonded with Zn2–Cp\*(<sub>centre</sub>) distances of 1.8858(4) Å (**1**) and 1.9215(3) Å (**2**) and Cp\*(<sub>centre</sub>)–Zn1–Zn2 bond angles of 177.2° (**1**) and 177.5° (**2**), which are close to linearity as observed in  $Zn_2Cp^*_2$ <sup>1</sup> and  $Cp^*ZnZnL$ .<sup>8b,14</sup> The Zn1 atoms are three-coordinated, and the Zn1–N1/2 bonds of **1** (1.9917(13),


*Institute of Inorganic Chemistry and Center for Nanointegration Duisburg-Essen (CENIDE), University of Duisburg-Essen, 45117 Essen, Germany.*  
 E-mail: stephan.schulz@uni-due.de

† Electronic supplementary information (ESI) available: Experimental, analytical (NMR, IR spectra, elemental analysis) and crystallographic data of **1–7**. CCDC 2111218 (**1**), 2111219 (**2**), 2111220 (**3**), 2111221 (**4**), 2111225 (**4S**), 2111222 (**5**), 2111223 (**6**) and 2111224 (**7**). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1cc05617d



Scheme 1 Synthesis of heteroleptic Zn(I) complexes **1** and **2**.





Scheme 2 Synthesis of complex 4.

2.0140(13) Å) are longer than those of **2** (1.9580(10), 1.9598(10) Å), in accordance with the reduced electron donor capacity of the fluorinated  $L^1$  ligand,<sup>17</sup> but comparable to those of homoleptic complexes  $L^2_2Zn_2$  and  $L^3_2Zn_2$ .<sup>4,7</sup>

With heteroleptic complex **2** in hand, we explored its reactivity toward heteroallenes. Homoleptic  $Cp^*_2Zn_2$  reacted with RNCO (R = *t*-Bu, Dipp) at ambient temperature with insertion into one Zn–Cp\* bond.<sup>14</sup> In contrast, the reaction of heteroleptic Zn(i) complex **2** with *t*-BuNCO at 70 °C for 4 days gave complex **4** in 41% yield (Scheme 2), whereas no reaction occurred with DippNCO even at 100 °C. The formation of **4** results from insertion of *t*-BuNCO into the Zn–Zn and Zn–Cp\* bonds and cleavage of a C=O bond accompanied by the formation of *t*-BuNC as was confirmed by *in situ* <sup>1</sup>H NMR spectroscopy (Fig. S39, ESI†). Any attempts to isolate reaction intermediates by varying the temperature and the molar ratio of the reagents failed. However, an excess of *t*-BuNCO promotes the reaction, as **4** was not formed in a 1 : 1 molar ratio reaction at 70 °C. The analogous reaction of  $L^3_2Zn_2$ <sup>7</sup> with *t*-BuNCO also quantitatively gave a zinc carbamate complex **4S** (Fig. S40 and S46, ESI†) and *t*-BuNC.

Complex **4** is thermally stable and decomposes at 210 °C. The <sup>1</sup>H NMR spectrum shows two singlets of the *t*-Bu groups and five singlets of the Cp\* group, indicating an asymmetric nature of the complex in solution. The <sup>13</sup>C NMR spectrum also shows five singlets of the Me groups of the Cp\* ligand as well as two singlets of the tertiary C atom of the *t*-Bu groups and resonances of the NCO units at 170.1 and 172.8 ppm, respectively.

Single crystals of **4**, which crystallize in the monoclinic space group  $P2_1/c$  (Fig. 1), were grown from a saturated toluene solution at 4 °C. Both Zn atoms adopt distorted tetrahedral coordination spheres and are bridged by a carbamate unit.

The Zn1–O2/3 (2.0697(8), 2.0309(8) Å) bond lengths are comparable to those of the zinc carbamate complex  $L^2ZnO_2CN(i-Pr)_2$  (2.028(2), 2.041(1) Å),<sup>18</sup> but longer compared to those of the carboxylate complexes  $[L^2Zn(\mu, \eta^2-O_2CR)]_2$  (R = H, Me, Ph, Oi-Pr), which range from 1.936 to 2.027 Å.<sup>19</sup> The Zn2–N3/4 (1.9000(9), 1.9169(9) Å) bond lengths are virtually identical to that of  $Cp^*Zn-Zn(N(t-Bu)C(Cp^*)O)$  (1.9148(9) Å),<sup>14</sup> while the Zn2–O3 distance (2.2780(8) Å) is rather long, indicating a rather weak coordinative interaction. The C–O (1.2819(12), 1.3273(12) Å) and C–N bond lengths (1.3270(13) Å) indicate a delocalized  $\pi$ -electron system within the carbamate unit. The Cp\* ligand is  $\eta^2$ -coordinated to Zn2, and the Zn–C bonds (2.4906(11), 2.5135(11) Å) are elongated compared to Zn– $\pi$  complexes with  $\eta^2$  interactions, *i.e.* alkyne-coordinated  $ZnBr_2$  (2.217(5), 2.393(5) Å)<sup>20</sup>





Fig. 1 Molecular structure of **4**. Thermal ellipsoids are drawn at 30% probability level. Parts of the ligands are drawn in wire/stick model, while H atoms are omitted for clarity.

and  $[PhC(Nt-Bu)_2(Cp^*)Si-Zn(Cp^*)Cl]$  (2.2519(26), 2.1224(29) Å),<sup>21</sup> but shorter than those in  $Zn(C_6F_5)_2(tol.)$  (2.784(2), 2.6847(15) Å)<sup>22</sup> and arylacetylene-substituted calix[4]arene zinc complexes (2.7695(37), 3.0667(37) Å).<sup>23</sup>

Complex **2** was then reacted with organoazides  $RN_3$  (R = Ph, Ad, SiMe<sub>3</sub>). In contrast to homoleptic  $L^3_2Zn_2$ , which was found to react with  $RN_3$  with formation of zinc hexazene  $[(L^3Zn)_2(\mu-\eta^2:\eta^2-N_6R_2)]$  (R = Ph, 2,6-*i*-Pr<sub>2</sub>C<sub>6</sub>H<sub>3</sub>) or dimeric zinc azide complexes  $[(L^3Zn)(\mu-N_3)]_2$  (R = Me<sub>3</sub>Si, Me<sub>3</sub>Sn),<sup>12c</sup> the reaction of heteroleptic complex **2** with 2 equiv. of AdN<sub>3</sub> at 70 °C for 2 days yielded the first bis-hexazene complex **5** (Scheme 3), which is thermally stable in solution up to 100 °C and in the solid state (decomposition temperature > 300 °C), respectively. Complex **5** is likely formed *via* the  $Cp^*Zn(\mu-\eta^2:\eta^2-N_6Ad_2)ZnL^2$  intermediate, followed by intramolecular elimination of  $Cp^*_2Zn$ .

In contrast, the reaction of **2** with PhN<sub>3</sub> gave the zinc triazenide complex **6** in 48% yield (Scheme 3). Alkaline or alkaline earth metal triazenides are typically formed in reactions of aryl azides and organolithium and magnesium

Scheme 3 Synthesis of complexes **5**–**7**.




Fig. 2 Molecular structure of **5–7**. Thermal ellipsoids drawn at 30% probability level, parts of the ligands are in a wire/stick model, and hydrogen atoms are omitted for clarity. **7** contains two molecules in the asymmetric unit (only the Zn(1)-containing molecule is discussed).

complexes,<sup>24</sup> hence the formation of **6** likely results from a nucleophilic attack of the Cp\* ligand. Since no reaction was observed in a control experiment of L<sup>2</sup>ZnCp\* with PhN<sub>3</sub>, we assume that the first reaction step is an insertion reaction of PhN<sub>3</sub> into the Zn–Zn bond of **2**. Low-valent metal complexes are known to react with RN<sub>3</sub> with formation of metal triazenides as was shown for homo- (Al, Cr) and heterobimetallic (In–Zn) complexes,<sup>25</sup> while a dinuclear iron complex was formed by the reaction of an Fe–N<sub>2</sub> complex with AdN<sub>3</sub>.<sup>26</sup> The reaction of **2** with Me<sub>3</sub>SiN<sub>3</sub> occurred with reductive elimination of Si<sub>2</sub>Me<sub>6</sub> as was reported for the analogous reaction of L<sup>3</sup>Zn<sub>2</sub><sup>12c</sup> and formation of complex **7** featuring a pseudo triangular Zn<sub>3</sub>N<sub>9</sub> moiety (Scheme 3). **7** also formed in 78% yield in the reaction of L<sup>2</sup>ZnH and Me<sub>3</sub>SiN<sub>3</sub>.

<sup>1</sup>H and <sup>13</sup>C NMR spectra of **5** and **6** show resonances of L<sup>2</sup> and Ad (**5**) and Cp\* and Ph (**6**), while **7** shows two sets of resonances of L<sup>2</sup> due to two conformers in solution, which form a temperature-dependent equilibrium as confirmed by VT-<sup>1</sup>H NMR (Fig. S41, ESI<sup>†</sup>). IR spectra show absorption bands of the hexazene (1265, 1218 cm<sup>-1</sup>, **5**), triazene (1313, 1255 cm<sup>-1</sup>, **6**) and azide groups (2157, 2124 cm<sup>-1</sup>, **7**).

Single crystals of **5–7** were grown from toluene solutions. Complexes **5** and **6** crystallize in the monoclinic space groups *P2<sub>1</sub>/n* and *P2<sub>1</sub>/m* and complex **7** in the triclinic space group *P1̄* (Fig. 2). Complex **5** contains two bridging hexazene ligands. The Zn1–N1/2 bonds within the C<sub>3</sub>N<sub>2</sub>Zn ring are slightly shorter than the Zn1–N3/6 bonds (2.0024(19), 2.0378(18) Å) in the neighbouring N<sub>4</sub>Zn ring, but comparable to Zn–N3/6 bonds (2.0079(18), 1.9817(19) Å) in the nonadjacent N<sub>4</sub>Zn ring. The N5–N6 distance (1.400(3) Å) is typical for a single bond, while the other N–N bond lengths (1.297(3)–1.301(3) Å) of the hexazene unit indicate an allyl-like nature as was previously reported for metal hexazene complexes.<sup>12c,27</sup> The Zn atom in triazene complex **6** is tetrahedrally coordinated by four N atoms of the L<sup>2</sup> ligand and the triazene group. The N–N bond lengths within the ZnN<sub>3</sub> metallacycle (1.297(6), 1.307(5) Å) indicate a delocalized π-electron system within the N<sub>3</sub> moiety. The Zn–N2 bonds (1.9592(10) Å) are shorter than the Zn–N1/3 bonds (2.0524(13), 2.0869(14) Å). The only structurally characterized zinc triazene complex [Dipp<sub>2</sub>N<sub>3</sub>]<sub>2</sub>Zn, which was prepared by an ethane elimination reaction of ZnEt<sub>2</sub> with Dipp<sub>2</sub>N<sub>3</sub>H,<sup>28</sup> shows comparable structure parameters.

In contrast to dimeric [(L<sup>3</sup>Zn)(μ-N<sub>3</sub>)<sub>2</sub>]<sub>2</sub>,<sup>29</sup> complex **7** forms a pseudo-triangular Zn<sub>3</sub>N<sub>9</sub> moiety with bridging N<sub>3</sub> units, resulting in an almost planar Zn<sub>3</sub>N<sub>9</sub> metallacycle (r.m.s. deviation from the least-squares plane 0.0655 Å), and each Zn atom is further coordinated by one L<sup>2</sup> ligand. The Zn–N bonds (1.952(5)–2.027(5) Å) within the Zn<sub>3</sub>N<sub>9</sub> moiety are slightly longer than those in the C<sub>3</sub>N<sub>2</sub>Zn rings (1.945(4)–1.959(5) Å). The N–N–N angles of 178.0(7)°, 179.1(7)° and 178.9(7)° are almost linear, and the N–N bond lengths range from 1.147(7) to 1.192(7) Å.

To summarize, heteroleptic Zn(I) complexes L<sup>1/2</sup>ZnZnCp\* (**1**, **2**) were synthesized and reactions of L<sup>2</sup>ZnZnCp\* **2** with heteroallenes are reported. The reaction with *t*-BuNCO proceeded with insertion into both the Zn–Zn and the Zn–Cp\* bonds and formation of carbamate complex **4**, whereas reactions with RN<sub>3</sub> yielded unprecedented bis-hexazene, triazene, and trimeric azide complexes **5–7**, respectively.

S. S. acknowledges support by the University of Duisburg-Essen.

## Conflicts of interest

There are no conflicts to declare.

## Notes and references

- I. Resa, E. Carmona, E. Gutierrez-Puebla and A. Monge, *Science*, 2004, **305**, 1136.
- (a) C.-S. Cao, Y. Shi, H. Xu and B. Zhao, *Coord. Chem. Rev.*, 2018, **365**, 122; (b) R. H. Duncan Lyngdoh, H. F. Schaefer and R. B. King, *Chem. Rev.*, 2018, **118**, 11626; (c) T. Li, S. Schulz and P. W. Roesky, *Chem. Soc. Rev.*, 2012, **41**, 3759; (d) E. Carmona and A. Galindo, *Angew. Chem., Int. Ed.*, 2008, **47**, 6526; (e) A. Grirrane, I. Resa, A. Rodríguez and E. Carmona, *Coord. Chem. Rev.*, 2008, **252**, 1532; (f) R. Ayala and A. Galindo, *J. Organomet. Chem.*, 2019, **898**, 120878.
- A. Grirrane, I. Resa, A. Rodríguez, E. Carmona, E. Alvarez, E. Gutierrez-Puebla, A. Monge, A. Galindo, D. del Río and R. A. Andersen, *J. Am. Chem. Soc.*, 2007, **129**, 693.
- (a) M. Ma, L. Shen, H. Wang, Y. Zhao, B. Wu and X.-J. Yang, *Organometallics*, 2020, **39**, 1440; (b) M. Juckel, D. Dange, C. de Bruin-Dickason and C. Jones, *Z. Anorg. Allg. Chem.*, 2020, **646**, 603; (c) A. Stasch, *Chem. – Eur. J.*, 2012, **18**, 15105; (d) Y. Liu, S. Li, X.-J. Yang, P. Yang, J. Gao, Y. Xia and B. Wu, *Organometallics*, 2009, **28**, 5270; (e) Z. Zhu, M. Brynda, R. J. Wright, R. C. Fischer, W. A. Merrill, E. Rivard, R. R. Wolf, J. C. Fetting, M. M. Olmstead and P. P. Power, *J. Am. Chem. Soc.*, 2007, **129**, 10847; (f) I. L. Fedushkin, A. A. Skatova, S. Y. Ketkov, O. V. Eremenko, A. V. Piskunov and



- G. K. Fukin, *Angew. Chem., Int. Ed.*, 2007, **46**, 4302; (g) Y.-C. Tsai, D.-Y. Lu, Y.-M. Lin, J.-K. Hwang and J.-S. K. Yu, *Chem. Commun.*, 2007, 4125; (h) X.-J. Yang, J. Yu, Y. Liu, Y. Xie, H. F. Schaefer, Y. Liang and B. Wu, *Chem. Commun.*, 2007, 2363; (i) Z. Zhu, R. J. Wright, M. M. Olmstead, E. Rivard, M. Brynda and P. P. Power, *Angew. Chem., Int. Ed.*, 2006, **45**, 5807; (j) Y. Z. Wang, B. Quillian, P. R. Wei, H. Y. Wang, X. J. Yang, Y. M. Xie, R. B. King, P. V. Schleyer, H. F. Schaefer and G. H. Robinson, *J. Am. Chem. Soc.*, 2005, **127**, 11944.
- 5 J. Hicks, E. J. Underhill, C. E. Kefalidis, L. Maron and C. Jones, *Angew. Chem., Int. Ed.*, 2015, **54**, 10000.
- 6 T. D. Lohrey, L. Maron, R. G. Bergman and J. Arnold, *J. Am. Chem. Soc.*, 2019, **141**, 800.
- 7 S. Schulz, D. Schuchmann, U. Westphal and M. Bolte, *Organometallics*, 2009, **28**, 1590.
- 8 (a) H. P. Nayek, A. Lühl, S. Schulz, R. Köppe and P. W. Roesky, *Chem. – Eur. J.*, 1773, **2011**, 17; (b) S. Schulz, S. Gondzik, D. Schuchmann, U. Westphal, L. Dobrzycki, R. Boese and S. Harder, *Chem. Commun.*, 2010, **46**, 7757.
- 9 (a) K. Mayer, L. Jantke, S. Schulz and T. F. Fässler, *Angew. Chem., Int. Ed.*, 2017, **56**, 2350; (b) S. Gondzik, D. Bläser, C. Wölper and S. Schulz, *Chem. – Eur. J.*, 2010, **16**, 13599.
- 10 (a) D. Schuchmann, U. Westphal, S. Schulz, U. Flörke, D. Bläser and R. Boese, *Angew. Chem., Int. Ed.*, 2009, **48**, 807; (b) M. Carrasco, R. Peloso, I. Resa, A. Rodríguez, L. Sánchez, E. Álvarez, C. Maya, R. Andreu, J. J. Calvente, A. Galindo and E. Carmona, *Inorg. Chem.*, 2011, **50**, 6361; (c) P. Jochmann and D. W. Stephan, *Chem. – Eur. J.*, 2014, **20**, 8370.
- 11 (a) S. Schulz, D. Schuchmann, I. Krossing, D. Himmel, D. Bläser and R. Boese, *Angew. Chem., Int. Ed.*, 2009, **48**, 5748; (b) M. Carrasco, R. Peloso, A. Rodríguez, E. Álvarez, C. Maya and E. Carmona, *Chem. – Eur. J.*, 2010, **16**, 9754; (c) K. Freitag, H. Banh, C. Ganesamoorthy, C. Gemel, R. W. Seidel and R. A. Fischer, *Dalton Trans.*, 2013, **42**, 10540; (d) H. Banh, C. Gemel, R. W. Seidel and R. A. Fischer, *Chem. Commun.*, 2015, **51**, 2170.
- 12 (a) S. Gondzik, D. Bläser, C. Wölper and S. Schulz, *J. Organomet. Chem.*, 2015, **783**, 92; (b) S. Gondzik, S. Schulz, D. Bläser and C. Wölper, *Chem. Commun.*, 2014, **50**, 1189; (c) S. Gondzik, S. Schulz, D. Bläser, C. Wölper, R. Haack and G. Jansen, *Chem. Commun.*, 2014, **50**, 927; (d) J. Gao, S. Li, Y. Zhao, B. Wu and X.-J. Yang, *Organometallics*, 2012, **31**, 2978.
- 13 (a) H. Banh, J. Hornung, T. Kratz, C. Gemel, A. Pöthig, F. Gam, S. Kahlal, J.-Y. Saillard and R. A. Fischer, *Chem. Sci.*, 2018, **9**, 8906; (b) K. Freitag, C. Gemel, P. Jerabek, I. M. Oppel, R. W. Seidel, G. Frenking, H. Banh, K. Dilchert and R. A. Fischer, *Angew. Chem., Int. Ed.*, 2015, **54**, 4370; (c) T. Bollermann, K. Freitag, C. Gemel, R. W. Seidel, M. von Hopffgarten, G. Frenking and R. A. Fischer, *Angew. Chem., Int. Ed.*, 2011, **50**, 772; (d) H. Banh, K. Dilchert, C. Schulz, C. Gemel, R. W. Seidel, R. Gautier, S. Kahlal, J.-Y. Saillard and R. A. Fischer, *Angew. Chem., Int. Ed.*, 2016, **55**, 3285.
- 14 B. Li, C. Wölper, K. Huse and S. Schulz, *Chem. Commun.*, 2020, **56**, 8643.
- 15 (a) M. Chen, S. Jiang, L. Maron and X. Xu, *Dalton Trans.*, 2019, **48**, 1931; (b) S. Jiang, M. Chen and X. Xu, *Inorg. Chem.*, 2019, **58**, 13213.
- 16 P. Cui, H.-S. Hu, B. Zhao, J. T. Miller, P. Cheng and J. Li, *Nat. Commun.*, 2015, **6**, 6331.
- 17 (a) K. Huse, C. Wölper and S. Schulz, *Eur. J. Inorg. Chem.*, 2018, 3472; (b) K. Huse, H. Weinert, C. Wölper and S. Schulz, *Dalton Trans.*, 2020, **49**, 9773.
- 18 M. H. Chisholm, J. Gallucci and K. Phomphrai, *Inorg. Chem.*, 2002, **41**, 2785.
- 19 (a) G. Feng, C. Du, L. Xiang, I. del Rosal, G. Li, X. Leng, E. Y.-X. Chen, L. Maron and Y. Chen, *ACS Catal.*, 2018, **8**, 4710; (b) C. Scheiper, C. Wölper, D. Bläser, J. Roll and S. Schulz, *Z. Naturforsch., B: J. Chem. Sci.*, 2014, **69**, 1365; (c) D. R. Moore, M. Cheng, E. B. Lobkovsky and G. W. Coates, *J. Am. Chem. Soc.*, 2003, **125**, 11911; (d) M. Cheng, E. B. Lobkovsky and G. W. Coates, *J. Am. Chem. Soc.*, 1998, **120**, 11018.
- 20 H. Lang, N. Mansilla and G. Rheinwald, *Organometallics*, 2001, **20**, 1592.
- 21 S. Schäfer, R. Köppe, M. T. Gamer and P. W. Roesky, *Chem. Commun.*, 2014, **50**, 11401.
- 22 A. Guerrero, E. Martin, D. L. Hughes, N. Kaltsoyannis and M. Bochmann, *Organometallics*, 2006, **25**, 3311.
- 23 E. Bukhaltsev, I. Goldberg, R. Cohen and A. Vignalok, *Organometallics*, 2007, **26**, 4015.
- 24 (a) S. G. Alexander, M. L. Cole, C. M. Forsyth, S. K. Furfari and K. Konstas, *Dalton Trans.*, 2009, 2326; (b) S.-O. Hauber, F. Lissner, G. B. Deacon and M. Niemeyer, *Angew. Chem., Int. Ed.*, 2005, **44**, 5871.
- 25 (a) W. Uhl, R. Gerding, S. Pohl and W. Saak, *Chem. Ber.*, 1995, **128**, 81; (b) C. Ni, B. D. Ellis, G. J. Long and P. P. Power, *Chem. Commun.*, 2009, 2332; (c) M. D. Anker, Y. Altaf, M. Lein and M. P. Coles, *Dalton Trans.*, 2019, **48**, 16588.
- 26 R. E. Cowley, N. J. DeYonker, N. A. Eckert, T. R. Cundari, S. DeBeer, E. Bill, X. Ottenwaelder, C. Flaschenriem and P. L. Holland, *Inorg. Chem.*, 2010, **49**, 6172.
- 27 (a) R. E. Cowley, J. Elhaik, N. A. Eckert, W. W. Brennessel, E. Bill and P. L. Holland, *J. Am. Chem. Soc.*, 2008, **130**, 6074; (b) S. J. Bonyhady, S. P. Green, C. Jones, S. Nembenna and A. Stasch, *Angew. Chem., Int. Ed.*, 2009, **48**, 2973; (c) S. J. Bonyhady, C. Jones, S. Nembenna, A. Stasch, A. J. Edwards and G. J. McIntyre, *Chem. – Eur. J.*, 2010, **16**, 938; (d) J. A. Bellow, P. D. Martin, R. L. Lord and S. Groysman, *Inorg. Chem.*, 2013, **52**, 12335; (e) L. Fohlmeister and C. Jones, *Aust. J. Chem.*, 2014, **67**, 1011; (f) S. Gondzik, C. Wölper, R. Haack, G. Jansen and S. Schulz, *Dalton Trans.*, 2015, **44**, 15703; (g) C. Stienen, S. Gondzik, A. Gehlhaar, R. Haack, C. Wölper, G. Jansen and S. Schulz, *Organometallics*, 2016, **35**, 1022.
- 28 N. Nimitsiriwat, V. C. Gibson, E. L. Marshall, P. Takolpuckdee, A. K. Tomov, A. J. P. White, D. J. Williams, M. R. J. Elsegood and S. H. Dale, *Inorg. Chem.*, 2007, **46**, 9988.
- 29 G. Bendt, S. Schulz, J. Spielmann, S. Schmidt, D. Bläser and C. Wölper, *Eur. J. Inorg. Chem.*, 2012, 3725.

