



Cite this: *Chem. Commun.*, 2022, 58, 64

Received 29th October 2021,  
Accepted 29th November 2021

DOI: 10.1039/d1cc06119d

rsc.li/chemcomm

# C,C'-Ru to C,B'-Ru isomerisation in bis(phosphine)Ru complexes of [1,1'-bis(ortho-carborane)]†‡

Rebekah J. Jeans, Georgina M. Rosair and Alan J. Welch\*

**We report herein the first example of the controlled isomerisation of a C,C'-bound (to metal) bis(ortho-carborane) ligand to C,B'-bound with no other change in the molecule. Since the C and B vertices of carboranes have different electron-donating properties this transformation allows the reactivity of the metal centre to be fine-tuned.**

Carboranes are exceedingly versatile ligands to transition-metals.<sup>1</sup> Deboronation of the neutral carborane [*closo*-1,2-C<sub>2</sub>B<sub>10</sub>H<sub>12</sub>] to the anion [*nido*-7,8-C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>2-</sup> affords a ligand which is isolobal with the ubiquitous Cp<sup>-</sup>, able to bind to metals in full η<sup>5</sup> fashion,<sup>2</sup> or η<sup><5</sup> in slipped metallocarboranes.<sup>3</sup> Alternatively carboranes, particularly anionic carboranes, are able to co-ordinate metals through one,<sup>4</sup> two<sup>5</sup> or three<sup>6</sup> B-H→M *B*-agostic interactions, taking advantage of the hydridic nature of H atoms bonded to B. Finally direct C-M<sup>7</sup> and B-M<sup>8</sup> sigma bonding is well established and is sometimes accompanied by B-H→M *B*-agostic bonding from adjacent B atoms.<sup>9</sup>

Whether a carborane binds directly to a metal, or to a substituent which is subsequently linked to a metal, through a C or B vertex is particularly important in that, everything else being equal, a *B*-bound carborane is more electron-donating than a *C*-bound carborane.<sup>10</sup> This affords two isomeric forms of the same ligand which are isosteric but not isoelectronic, and recently this has been exploited to fine-tune the properties of metal-carborane complexes.<sup>11</sup>

Bis(carboranes) are molecules composed of two carborane moieties connected by a direct C-C, C-B or B-B bond and, of the various possible bis(carboranes), [1-(1'-*closo*-1',2'-C<sub>2</sub>B<sub>10</sub>H<sub>11</sub>)-*closo*-1,2-C<sub>2</sub>B<sub>10</sub>H<sub>11</sub>] or more simply [1,1'-bis(*ortho*-carborane)], is the most

studied and has undergone a resurgence of interest in recent years.<sup>12</sup> Following double deprotonation at the protonic C2H and C2'H sites, [1,1'-bis(*ortho*-carborane)] can be used as a κ<sup>2</sup> chelating ligands in both homoleptic<sup>13</sup> and heteroleptic<sup>9b,11c,d,12f,14</sup> transition-metal complexes.

In 2016 we reported catalytically-active (arene)Ru complexes of doubly-deprotonated [1,1'-bis(*ortho*-carborane)] in which the metal coordination was completed by a B3'-H→Ru *B*-agostic interaction.<sup>9b</sup> Reaction of these compounds with phosphine (2 × PPh<sub>3</sub> or dppe) resulted in displacement of the arene, coordination of the phosphine and a change in the ligating mode of the bis(carborane) from X<sub>2</sub>(C,C')L (L = agostic interaction) to X<sub>2</sub>(C,B')L, the first time C,B' ligation of [1,1'-bis(*ortho*-carborane)] had been observed.<sup>9b</sup> Subsequently, Spokoiny and co-workers reported the synthesis of an isomeric mixture of Pt complexes of [1,1'-bis(*ortho*-carborane)] with bipyridyl coligands;<sup>11c</sup> in one isomer the bis(carborane) was C,C'-bound and in the other it was C,B'-bound (subsequently he was able to prepare exclusively the C,C'-bound isomer by using a different synthetic strategy).<sup>11d</sup> Heating the mixture 'under forcing conditions' did not drive it to one isomer suggesting that the two isomers were formed *via* different pathways.

Thus, although it is potentially of great interest to be able to isomerise bis(carborane) from C,C'-bound to a metal centre to C,B'-bound under controlled conditions, no system has so far achieved this. We now describe the controlled isomerisation of a C,C'-bound bis(phosphine) ruthenium complex to its C,B'-bound isomer.

The room temperature reaction of [Ru(κ<sup>3</sup>-2,2',3'-{1-(1'-*closo*-1',2'-C<sub>2</sub>B<sub>10</sub>H<sub>10</sub>)-*closo*-1,2-C<sub>2</sub>B<sub>10</sub>H<sub>10</sub>})(*p*-cymene)] (I) with 5 equivalents of PMePh<sub>2</sub> in THF produced a deep-red solution from which both red and yellow components were isolated by preparative thin-layer chromatography (TLC). Although the yellow product appeared stable to work-up, repeated chromatography of the red product (at room temperature) always afforded a small amount of the yellow species, implying that the red and yellow species were related as kinetically- and thermodynamically-stable isomers.

Institute of Chemical Sciences, Heriot-Watt University, Edinburgh EH14 4AS, UK.

E-mail: a.j.welch@hw.ac.uk

† Dedicated with very best wishes to Professors Francesc Teixidor and Clara Viñas, colleagues and good friends for more than 30 years, on the occasion of their 70th birthdays.

‡ Electronic supplementary information (ESI) available. CCDC 2117898 and 2117899. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1cc06119d



Repeating the reaction at 0 °C, eliminating the chromatographic work-up and crystallising the product at –20 °C allowed the red species (**1**) to be isolated in good yield (80%) in pure form. § Mass spectrometry of **1** gave a molecular ion consistent with displacement of the *p*-cymene ligand of **I** by two PMePh<sub>2</sub> ligands. Although the <sup>11</sup>B{<sup>1</sup>H} NMR spectrum at –50 °C was largely uninformative the <sup>1</sup>H spectrum revealed, in addition to multiplet resonances associated with the Ph groups, two doublets arising from the two PMePh<sub>2</sub> units, implying the two phosphine ligands are inequivalent and confirmed by the presence of two mutual doublets, *J*<sub>PP</sub> = 28.3 Hz, in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum. Importantly, the <sup>1</sup>H{<sup>11</sup>B} NMR spectrum showed, in addition to resonances between 3 and –1 ppm associated with cage BH<sub>exo</sub> atoms, a doublet resonance at –3.27 ppm integrating for 1H and indicative of B–H→Ru (showing coupling to only one P atom). Notably absent from the <sup>1</sup>H and <sup>1</sup>H{<sup>11</sup>B} spectra of **1** was a resonance arising from cage CH.

Collectively these data suggest that in **1** the bis(*ortho*-carborane) unit is bound to the Ru atom in X<sub>2</sub>(C,C')L mode, *i.e.* via both cage C atoms, unlike the situation in the previously isolated PPh<sub>3</sub> and dppe analogues,<sup>9b</sup> and this was subsequently confirmed by crystallographic analysis (Fig. 1). ¶


The bis(carborane) unit is indeed bonded to the metal atom via σ bonds from C2 and C2' and a B3'–H3'→Ru *B*-agostic interaction; thus compound **1** is formulated as [Ru(κ<sup>3</sup>-2,2',3'–{1-(1'-*closo*-1',2'-C<sub>2</sub>B<sub>10</sub>H<sub>10</sub>)-*closo*-1,2-C<sub>2</sub>B<sub>10</sub>H<sub>10</sub>})(PMePh<sub>2</sub>)<sub>2</sub>]. The geometry at Ru is approximately square-pyramidal (C2 apex). The Ru–C2' σ bond is particularly distorted, as evidenced by the angle Ru1–C2'–P *ca.* 134° *cf.* Ru1–C2–Q *ca.* 164° (*P* and *Q* are the centroids of the primed and unprimed icosahedra, respectively), presumably as a result of the need to accommodate C2' and the B3'H3' unit in two *cis* ligand positions. The Ru–C bond lengths are significantly different (shorter to C2), as are the

Ru–P bond lengths (shorter to P1), in both cases reflecting the relative trans influences of the ligands (or vacant site) opposite.


Solutions of **1** slowly change from deep red to yellow in colour as the compound isomerises to a new species **2**, a process easily followed by <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy with two new higher-frequency doublets growing in at the expense of the original ones. At room temperature in CD<sub>2</sub>Cl<sub>2</sub> the conversion is typically 15% after 6 h but is accelerated on heating (*ca.* 75% conversion after 2 h at 40 °C) and retarded on addition of excess phosphine (*ca.* 10% conversion after 24 h).

Compound **2** can be conveniently prepared in good yield (64%) by repeating the original synthesis at room temperature and then stirring for *ca.* 2 h at 40 °C followed by work-up involving column chromatography. || Mass spectrometry is fully consistent with **2** being an isomer of compound **1**. The <sup>11</sup>B{<sup>1</sup>H} NMR spectrum of **2** is again relatively uninformative save that, as for **1**, the chemical shifts imply a *closo* cage. The <sup>1</sup>H NMR spectrum of **2** again reveals two sets of doublets for the methyl protons of the PMePh<sub>2</sub> ligands and, additionally, an integral-1 resonance assigned to C<sub>cage</sub>H which unfortunately overlaps with the high-frequency component of one of the CH<sub>3</sub> doublets (δ 1.85 ppm). However, further evidence for a cage {CH} unit derives from a resonance in the <sup>13</sup>C NMR spectrum at δ 67.5 ppm assigned as CH by DEPT spectroscopy. The presence of a B–H→Ru interaction in **2** was established by the observation of an integral-1 doublet at –3.99 ppm in the <sup>1</sup>H{<sup>11</sup>B} NMR spectrum.

Thus the NMR data imply an X<sub>2</sub>(C,B')L bonding mode for the bis(*ortho*-carborane) unit in **2** as has previously been established for the PPh<sub>3</sub> and dppe analogues,<sup>9b</sup> and this was subsequently confirmed by an X-ray diffraction study (Fig. 2). \*\* Crystals of **1** and **2** (both studied as their 0.5CH<sub>2</sub>Cl<sub>2</sub> solvates) are isomorphous and at a molecular level the two species differ only in the relative positions of a C and B atom in one cage.




**Fig. 1** Structure of compound **1** (solvent omitted) with thermal ellipsoids drawn at the 50% probability level, except for H atoms. Both phenyl groups bound to P2 are partially disordered. Ru1–C2 2.0452(15), Ru1–C2' 2.1472(14), Ru1–H3' 1.872(17), Ru1–B3' 2.3388(15), Ru1–P1 2.2769(3), Ru1–P2 2.3341(4), C1–C1' 1.5113(19) Å.



**Fig. 2** Structure of compound **2** (solvent omitted) with thermal ellipsoids drawn as in Fig. 1. The view is chosen to reflect similarity with the structure of compound **1**. Ru1–C2 2.1375(13), Ru1–B3' 2.0273(15), Ru1–H6 1.867(18), Ru1–B6 2.3265(14), Ru1–P3 2.2555(3), Ru1–P4 2.3153(3), C1–C1' 1.5116(18) Å.





**Scheme 1** Suggested transformation of compound **1** (left) into compound **2** (right).

It was therefore imperative that cage vertices were correctly identified as either C or B in both crystallographic studies and this was established unambiguously by use of the VCD and BHD methods.<sup>15</sup>

The atomic numbering chosen for **2** reflects the likelihood that the B atom now  $\sigma$ -bonded to Ru (B3') is the same atom which was involved in the B–agostic bond in **1**, chemically-sensible in that the B3'–H3' bond of **1** would be activated by such an interaction. In compound **2** the B–H→Ru interaction is now from the unprimed cage and involves B6H6 by conventional numbering. Thus, as compound **1** isomerises to compound **2** the essential changes may be summarised as; breaking of the B3'–H3' bond and formation of a direct Ru–B3' bond; breaking of the Ru–C2' bond; formation of a new B6–H6→Ru bond; and net transfer of H from B3' to C2'. These changes are summarised in Scheme 1, but at this stage we do not have any detailed mechanistic information. Compound **2** is therefore established as [Ru( $\kappa^3$ -2,3',6-{1-(1'-*closo*-1',2'-C<sub>2</sub>B<sub>10</sub>H<sub>10</sub>)-*closo*-1,2-C<sub>2</sub>B<sub>10</sub>H<sub>10</sub>})(PMePh<sub>2</sub>)<sub>2</sub>]. In compound **2** the two phosphine ligands again lie opposite the C–Ru  $\sigma$  bond and the B–agostic bond and have been labelled P3 and P4 to avoid any implied direct relationship to the phosphines in **1**. Fully consistent with the structure of **1** there is considerable distortion of the Ru1–C2 bonding relative to the Ru1–B3' bonding (Ru1–C2–R *ca.* 134° *cf.* Ru1–B3'–S *ca.* 158° where R and S are the centroids of the primed and unprimed icosahedra, respectively) and the relative lengths of the Ru–P bonds (shorter to P3) reflect the nature of the trans unit.

In conclusion we have demonstrated the first controlled isomerisation of a bis(carborane) ligand from C,C'-bound to a metal centre to C,B'-bound, with no other change in the molecule. In the C,B'-bound isomer the metal centre will be relatively electron rich, and so this kind of isomerisation has the potential to allow the properties of the molecule, including catalytic properties, to be tuned in a controlled way.

We thank the Engineering & Physical Sciences Research Council and the CRITICAT Centre for Doctoral Training for a PhD studentship awarded to R. J. J. (grant no. EP/L016419/1). We also thank Prof. S. A. Macgregor for many helpful discussions.

## Conflicts of interest

There are no conflicts to declare.

## Notes and references

§ NMR data for **1** (CD<sub>2</sub>Cl<sub>2</sub>, –50 °C): <sup>1</sup>H{<sup>1</sup>H};  $\delta$  14 to –22 overlapping resonances with prominent maxima at –3.7, –7.9 (assume 20B). <sup>1</sup>H;  $\delta$  7.51–7.45 [m, 2H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>], 7.44–7.38 [m, 1H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>], 7.36–7.23 [m, 7H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>], 7.21–7.09 [m, 6H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>], 7.06–6.99 [m, 2H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>], 6.96–6.89 [m, 2H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>], 2.35, [d, *J*<sub>HP</sub> = 7.9 Hz, 3H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>], 2.16 [d, *J*<sub>HP</sub> = 10.0 Hz, 3H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>]. <sup>1</sup>H{<sup>1</sup>B} as for 1H plus  $\delta$  2.67 to 2.08 considerable overlap of BH and CH<sub>3</sub> resonances (total integral 9BH + 6CH) with prominent BH maxima at 2.61, 2.46 and 2.24, 1.89 (4H), 1.75 to 1.47 overlapping resonances with prominent maxima at 1.70, 1.67, 1.60 and 1.55 (total integral 5H, BH), 0.95 (1H, BH), –3.27 (d, *J*<sub>HP</sub> = 31.0 Hz, 1H, BHRu). <sup>31</sup>P{<sup>1</sup>H};  $\delta$  34.3 (d, *J*<sub>PP</sub> = 28.3 Hz, 1P), 22.6 (d, *J*<sub>PP</sub> = 28.3 Hz, 1P). EIMS: envelope centred on *m/z* 786.3 (M<sup>+</sup>).

¶ Crystal data for 1.0.5CH<sub>2</sub>Cl<sub>2</sub>: C<sub>30.5</sub>H<sub>47</sub>B<sub>20</sub>ClP<sub>2</sub>Ru, *M* = 828.34, monoclinic, *C*2/*c*, *a* = 38.4205(9), *b* = 10.5927(2), *c* = 21.1877(5) Å,  $\beta$  = 111.3870(10)°, *V* = 8029.1(3) Å<sup>3</sup>, *F*(000) = 3368.0 e, *D*<sub>calc</sub> = 1.371 Mg m<sup>–3</sup>, Cu–K $\alpha$  X-radiation,  $\lambda$  = 1.54178 Å,  $\mu$  = 4.693 mm<sup>–1</sup>, of 54551 data measured to  $\theta_{\max}$  = 74.78° on a Bruker D8 Venture diffractometer, 8168 were unique and were used to solve and refine the structure to *R* = 0.0213, *wR*<sub>2</sub> = 0.0529, CCDC 2117898‡.

|| NMR data for **2** (CD<sub>2</sub>Cl<sub>2</sub>, room temperature): <sup>1</sup>H{<sup>1</sup>H};  $\delta$  0.2 (2B), –1.9 to –15.9 (overlapping resonances with maxima at –4.6, –8.7, and –14.6, total integral 17B), –17.0 (1B). <sup>1</sup>H;  $\delta$  7.60–7.55 [m, 2H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>], 7.43–7.20 [m, 18H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>], 2.34 [d, *J*<sub>PH</sub> = 7.9 Hz, 3H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>], 1.85 (s, 1H, C<sub>cage</sub>H) overlapping with 1.84 [d, *J*<sub>PH</sub> = 9.5 Hz, 3H, PCH<sub>3</sub>(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>]. <sup>1</sup>H{<sup>1</sup>B} as for <sup>1</sup>H plus  $\delta$  2.53 (2H, BH), 2.50 (1H, BH), 2.38 (1H, BH), 2.26 (3H, BH), 2.19 (1H, BH), 2.17 (2H, BH), 2.10 (2H, BH), 2.00 (1H, BH), 1.96 (1H, BH), 1.76 (1H, BH), 1.58 (3H, BH), –3.99 (d, *J*<sub>HP</sub> = 27.6 Hz, 1H, BHRu). <sup>13</sup>C;  $\delta$  132.4 (d, *J*<sub>CP</sub> = 11.3 Hz, C<sub>arom</sub>,H), 132.1 (d, *J*<sub>CP</sub> = 10.5 Hz, C<sub>arom</sub>,H), 131.8 (d, *J*<sub>CP</sub> = 11.2 Hz, C<sub>arom</sub>,H), 131.5 (d, *J*<sub>CP</sub> = 9.7 Hz, C<sub>arom</sub>,H), 130.8 (d, *J*<sub>CP</sub> = 2.2 Hz, C<sub>arom</sub>,H), 130.4 (d, *J*<sub>CP</sub> = 2.0 Hz, C<sub>arom</sub>,H), 130.3 (d, *J*<sub>CP</sub> = 2.5 Hz, C<sub>arom</sub>,H), 130.2 (d, *J*<sub>CP</sub> = 1.9 Hz, C<sub>arom</sub>,H), 129.3 (d, *J*<sub>CP</sub> = 9.7 Hz, C<sub>arom</sub>,H), 129.1 (d, *J*<sub>CP</sub> = 9.3 Hz, C<sub>arom</sub>,H), 128.9 (d, *J*<sub>CP</sub> = 9.3 Hz, C<sub>arom</sub>,H), 128.7 (d, *J*<sub>CP</sub> = 9.6 Hz, C<sub>arom</sub>,H), 91.8 (C), 77.8 (C), 67.5 (C<sub>cage</sub>H), 18.8 (d, *J*<sub>CP</sub> = 28.2 Hz, CH<sub>3</sub>), 14.2 (d, *J*<sub>CP</sub> = 34.6 Hz, CH<sub>3</sub>). <sup>31</sup>P{<sup>1</sup>H};  $\delta$  41.5 (d, *J*<sub>PP</sub> = 28.0 Hz, 1P), 28.0 (d, *J*<sub>PP</sub> = 28.0 Hz, 1P). EIMS: envelope centred on *m/z* 786.3 (M<sup>+</sup>), 586.2 (M<sup>+</sup> – PMePh<sub>2</sub>).

\*\* Crystal data for 2.0.5CH<sub>2</sub>Cl<sub>2</sub>: C<sub>30.5</sub>H<sub>47</sub>B<sub>20</sub>ClP<sub>2</sub>Ru, *M* = 828.34, monoclinic, *C*2/*c*, *a* = 38.0931(9), *b* = 10.6717(3), *c* = 21.3413(5) Å,  $\beta$  = 111.8580(10)°, *V* = 8051.9(4) Å<sup>3</sup>, *F*(000) = 3368.0 e, *D*<sub>calc</sub> = 1.367 Mg m<sup>–3</sup>, Cu–K $\alpha$  X-radiation,  $\lambda$  = 1.54178 Å,  $\mu$  = 4.680 mm<sup>–1</sup>, 64154 data to  $\theta_{\max}$  = 74.73°, 8221 unique, *R* = 0.0208, *wR*<sub>2</sub> = 0.0512, CCDC 2117899‡.

- 1 R. N. Grimes, *Carboranes*, Elsevier, Amsterdam, The Netherlands, 3rd edn, 2016.
- 2 M. F. Hawthorne, D. C. Young, T. D. Andrews, D. V. Howe, R. L. Pilling, A. D. Pitts, M. Reintjes, L. F. Warren Jnr and P. A. Wegner, *J. Am. Chem. Soc.*, 1968, **90**, 879–896.
- 3 D. M. P. Mingos, M. I. Forsyth and A. J. Welch, *J. Chem. Soc., Dalton Trans.*, 1978, 1363–1374.
- 4 G. G. Hlatky, H. W. Turner and R. R. Eckman, *J. Am. Chem. Soc.*, 1989, **111**, 2728–2729.
- 5 S. A. Brew, J. C. Jeffery, M. D. Mortimer and F. G. A. Stone, *J. Chem. Soc., Dalton Trans.*, 1993, 1365–1374.
- 6 C. B. Knobler, T. B. Marder, E. A. Mizusawa, R. G. Teller, J. A. Long, P. E. Behnken and M. F. Hawthorne, *J. Am. Chem. Soc.*, 1984, **106**, 2990–3004.
- 7 N. Bresciani, M. Calligaris, P. Delise, G. Nardin and L. Randaccio, *J. Am. Chem. Soc.*, 1974, **96**, 5642–5643.
- 8 I. A. Lobanova, V. I. Bregadze, S. V. Timofeev, P. V. Petrovskii, Z. A. Starikova and F. M. Dolgushin, *J. Organomet. Chem.*, 2000, **597**, 48–53.
- 9 (a) R. A. Love and R. Bau, *J. Am. Chem. Soc.*, 1972, **94**, 8274–8276; (b) L. E. Riley, A. P. Y. Chan, J. Taylor, W. Y. Man, D. Ellis, G. M. Rosair, A. J. Welch and I. B. Sivaev, *Dalton Trans.*, 2016, **45**, 1127–1137.
- 10 (a) Z. Zheng, M. Diaz, C. B. Knobler and M. F. Hawthorne, *J. Am. Chem. Soc.*, 1995, **117**, 12338–12339; (b) F. Teixidor, R. Núñez, C. Viñas, R. Sillanpää and R. Kivekäs, *Angew. Chem., Int. Ed.*, 2000, **39**, 4290–4292; (c) F. Teixidor, G. Barberà, A. Vaca, R. Kivekäs, R. Sillanpää, J. Oliva and C. Viñas, *J. Am. Chem. Soc.*, 2005, **127**, 10158–10159.



- 11 (a) A. M. Spokoyny, C. W. Machan, D. J. Clingerman, M. S. Rosen, M. J. Wiester, R. D. Kennedy, C. L. Stern, A. A. Sarjeant and C. A. Mirkin, *Nat. Chem.*, 2011, **3**, 590–596; (b) A. M. Spokoyny, C. D. Lewis, G. Teverovskiy and S. L. Buchwald, *Organometallics*, 2012, **31**, 8478–8481; (c) K. O. Kirlikovali, J. C. Axtell, A. Gonzalez, A. C. Phung, S. I. Khan and A. M. Spokoyny, *Chem. Sci.*, 2016, **7**, 5132–5138; (d) K. O. Kirlikovali, J. C. Axtell, K. Anderson, P. I. Djurovich, A. L. Rheingold and A. M. Spokony, *Organometallics*, 2018, **37**, 3122–3131.
- 12 Recent publications: (a) I. B. Sivaev and V. I. Bregadze, *Coord. Chem. Rev.*, 2019, **392**, 146–176 and references therein; (b) S. Yruegas, J. C. Axtell, K. O. Kirlikovali, A. M. Spokoyny and C. D. Martin, *Chem. Commun.*, 2019, **55**, 2892–2895; (c) J. Wu, K. Cao, C.-Y. Zhang, T.-T. Xu, L.-F. Ding, B. Li and J. Yang, *Org. Lett.*, 2019, **21**, 5986–5989; (d) R. J. Jeans, A. P. Y. Chan, L. E. Riley, J. Taylor, G. M. Rosair, A. J. Welch and I. B. Sivaev, *Inorg. Chem.*, 2019, **58**, 11751–11761; (e) A. Benton, D. J. Durand, Z. Copeland, J. D. Watson, N. Fey, S. M. Mansell, G. M. Rosair and A. J. Welch, *Inorg. Chem.*, 2019, **58**, 14818–14829; (f) I. Chambrier, D. L. Hughes, R. J. Jeans, A. J. Welch, P. H. M. Budzelaar and M. Bochmann, *Chem. – Eur. J.*, 2020, **26**, 939–947; (g) A. P. Y. Chan, J. A. Parkinson, G. M. Rosair and A. J. Welch, *Inorg. Chem.*, 2020, **59**, 2011–2023; (h) A. P. Y. Chan, G. M. Rosair and A. J. Welch, *Molecules*, 2020, **25**, 519; (i) A. J. Welch, *Structure and Bonding*, Springer, Berlin, Heidelberg, 2021, DOI: 10.1007/430\_2020\_80.
- 13 (a) M. F. Hawthorne and D. A. Owen, *J. Am. Chem. Soc.*, 1971, **93**, 873–880; (b) D. E. Harwell, J. McMillan, C. B. Knobler and M. F. Hawthorne, *Inorg. Chem.*, 1997, **36**, 5951–5955.
- 14 (a) M. J. Martin, W. Y. Man, G. M. Rosair and A. J. Welch, *J. Organomet. Chem.*, 2015, **798**, 36–40; (b) Z.-J. Yao, Y.-Y. Zhang and G.-X. Jin, *J. Organomet. Chem.*, 2015, **798**, 274–277.
- 15 (a) A. McAnaw, G. Scott, L. Elrick, G. M. Rosair and A. J. Welch, *Dalton Trans.*, 2013, **42**, 645–664; (b) A. McAnaw, M. E. Lopez, D. Ellis, G. M. Rosair and A. J. Welch, *Dalton Trans.*, 2014, **43**, 5095–5105; (c) A. J. Welch, *Crystals*, 2017, **7**, 234.

