



Cite this: *Phys. Chem. Chem. Phys.*, 2021, 23, 12494

Correction: Histidine protonation controls structural heterogeneity in the cyanobacteriochrome AnPixJg2

Aditya G. Rao,^a Christian Wiebeler,^a Saumik Sen,^a David S. Cerutti^b and Igor Schapiro*^a

DOI: 10.1039/d1cp90071d

rsc.li/pccp

Correction for 'Histidine protonation controls structural heterogeneity in the cyanobacteriochrome AnPixJg2' by Aditya G. Rao *et al.*, *Phys. Chem. Chem. Phys.*, 2021, DOI: 10.1039/d0cp05314g.

The authors would like to update Fig. 6 in the published article with the correct version shown below.




Fig. 6 Absorption and CD spectra computed using RI-ADC(2). (A) Absorption spectra for substate $D-\alpha_f$ (solid) and substate $D-\beta_f$ (dotted) of the DPH model. (B) Absorption spectra for substate $D-\alpha_f$ (solid) of the SPH model. (C) CD spectra for substate $D-\alpha_f$ (solid) and substate $D-\beta_f$ (dotted) of the DPH model. (D) CD spectra for substate $D-\alpha_f$ (solid) of the SPH model. The spectra are averages over 100 snapshots for each substate.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Fritz Haber Center for Molecular Dynamics Research, Institute of Chemistry, The Hebrew University of Jerusalem, Jerusalem 91904, Israel.
 E-mail: igor.schapiro@mail.huji.ac.il

^b Department of Chemistry and Chemical Biology, Rutgers University, USA

