



Cite this: *Phys. Chem. Chem. Phys.*,
2021, 23, 21350

Correction: Self-diffusion, velocity cross-correlation, distinct diffusion and resistance coefficients of the ionic liquid [BMIM][Tf₂N] at high pressure

Kenneth R. Harris ^a and Mitsuhiro Kanakubo ^b

DOI: 10.1039/d1cp90181h

rsc.li/pccp

Correction for 'Self-diffusion, velocity cross-correlation, distinct diffusion and resistance coefficients of the ionic liquid [BMIM][Tf₂N] at high pressure' by Kenneth R. Harris et al., *Phys. Chem. Chem. Phys.*, 2015, 17, 23977–23993, DOI: 10.1039/C5CP04277A.

The published version of this manuscript contained errors in Table 5 on page 23983. Values for the coefficients for the VFT equation (17), for the ionic self-diffusion coefficients of [HMIM][Tf₂N] at 0.1 MPa were entered in error for [OMIM][Tf₂N]. The correct [OMIM][Tf₂N] values are given below:

| [OMIM][Tf ₂ N] | | |
|---------------------------|---------------|---------------|
| | D_{S+} | D_{S-} |
| A^a | 10.266 (0.27) | 11.671 (0.76) |
| $-B/K$ | 1211.5 (101) | 1802.1 (348) |
| T_0/K | 142.20 (7.6) | 102.74 (22) |
| $u_r^b/\%$ | 1.3 | 2.3 |
| T range/ $^{\circ}C$ | 25–89 | 30–90 |

^aThe standard errors for the fitted coefficients are given in parentheses. D_{Si} units: $10^{-12} \text{ m}^2 \text{ s}^{-1}$. ^b u_r , relative standard uncertainty of the fit.

In Fig. 9 and 12, the units on the ordinate labels should read $\text{m}^2 \cdot \text{s}^{-1} \cdot \text{K}^{-1}$.

These changes have no effect on the discussion or the conclusions of this work.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a School of Physical, Environmental and Mathematical Sciences, University College, University of New South Wales, P. O. Box 7916, Canberra BC, ACT 2610, Australia.
E-mail: k.harris@adfa.edu.au

^b National Institute of Advanced Industrial Science and Technology (AIST), 4-2-1 Nigatake, Miyagino-ku, Sendai 983-8551, Japan. E-mail: m-kanakubo@aist.go.jp

