

CORRECTION

View Article Online
View Journal | View Issue



Cite this: *Environ. Sci.: Nano*, 2021, 8, 1481

DOI: 10.1039/d1en90014e

rscl.li/es-nano

Correction: Potentiation of polycyclic aromatic hydrocarbon uptake in zebrafish embryos by nanoplastics

Yueyang Zhang^a and Greg G. Goss^{*abc}

Correction for 'Potentiation of polycyclic aromatic hydrocarbon uptake in zebrafish embryos by nanoplastics' by Yueyang Zhang *et al.*, *Environ. Sci.: Nano*, 2020, 7, 1730–1741, DOI: 10.1039/D0EN00163E.

In the Fig. 2A image, the colour of ¹⁴C-phenanthrene + 20 nm and ¹⁴C-phenanthrene + 500 nm were switched in error: ¹⁴C-phenanthrene + 20 nm group is labelled as the red triangle but it is supposed to be green, and the green dot (¹⁴C-phenanthrene + 500 nm) is supposed to be red. The corrected image, caption and sentence are shown below.




Fig. 2 Percentage of initial ¹⁴C-phenanthrene that either passed through the chorionic membrane of zebrafish embryos (A), or accumulated in zebrafish larvae (B) after incubation for various periods in the control (¹⁴C-phenanthrene alone no PS-NPs), in the presence of 20 nm PS-NPs at 10 mg L⁻¹, or 500 nm PS-NPs at 10 mg L⁻¹. The rate ($\mu\text{g kg}^{-1} \text{h}^{-1}$) of ¹⁴C-phenanthrene crossing the zebrafish embryo chorion over 4 h (C) and uptake by larvae over 24 h (D) with or without 20 nm or 500 nm PS-NPs. Means sharing the same letter are not significantly different from each other ($p > 0.05$). Values are mean \pm SEM. $n = 4$.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Department of Biological Sciences, University of Alberta, 11455 Saskatchewan Drive, Edmonton, Alberta, T6G 2E9, Canada. E-mail: greg.goss@ualberta.ca; Tel: +1 (780) 492 2381

^b National Institute for Nanotechnology, 11421 Saskatchewan Drive, Edmonton, Alberta, T6G 2M9, Canada

^c Director of Office of Environmental Nanosafety, University of Alberta, Canada

