

CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#)



Cite this: *Food Funct.*, 2021, **12**, 12159

Correction: Hepatoprotective effect of piceatannol against carbon tetrachloride-induced liver fibrosis in mice

Wei-Lun Hung,^a Yi-Ting Hsiao,^b Yi-Shiou Chiou,^b Kalyanam Nagabhushanam,^c Chi-Tang Ho^d and Min-Hsiung Pan^{*b,e,f}

DOI: 10.1039/d1fo90103f

rsc.li/food-function

Correction for 'Hepatoprotective effect of piceatannol against carbon tetrachloride-induced liver fibrosis in mice' by Wei-Lun Hung *et al.*, *Food Funct.*, 2021, DOI: 10.1039/D1FO02545G.

The authors regret that an incorrect version of Fig. 3 was included in the original article. The correct version of Fig. 3 is presented below.

^aSchool of Food Safety, Taipei Medical University, Taipei, 11031, Taiwan

^bInstitute of Food Science and Technology, National Taiwan University, Taipei 10617, Taiwan. E-mail: mhpan@ntu.edu.tw; Fax: +(886)-2-33661771; Tel: +(886)-2-33664133

^cSabinsa Corporation, East Windsor, New Jersey 08520, USA

^dDepartment of Food Science, Rutgers University, New Brunswick, New Jersey 08901, USA

^eDepartment of Medical Research, China Medical University Hospital, China Medical University, Taichung 40402, Taiwan

^fDepartment of Health and Nutrition Biotechnology, Asia University, Taichung, 41354, Taiwan






Fig. 3 Effects of resveratrol and piceatannol on CCl₄-induced histopathological alterations and collagen deposition in the preventive model. Representative haematoxylin and eosin-stained sections are shown in (A) the control group, (B) CCl₄-alone treatment group, (C) the CCl₄ + silymarin (200 mg per kg bw) treatment group, (D) CCl₄ + resveratrol (30 mg per kg bw) treatment group, (E) CCl₄ + piceatannol (30 mg per kg bw) treatment group and (F) CCl₄ + piceatannol (60 mg per kg bw) treatment group. The length of the scale bar of (A) to (F) is 60 μ m. Arrowheads (\downarrow and \downarrow) in (A) to (F) point to leukocyte infiltration and hepatocyte ballooning degeneration area in liver sections, respectively. Representative Picrosirius red-stained liver sections are shown in (G) the control group, (H) CCl₄-alone treatment group, (I) CCl₄ + silymarin (200 mg per kg bw) treatment group, (J) CCl₄ + resveratrol (30 mg per kg bw) treatment group, (K) CCl₄ + piceatannol (30 mg per kg bw) treatment group and (L) CCl₄ + piceatannol (60 mg per kg bw) treatment group. (M) Quantification of collagen deposition. The length of scale bar of (G) to (L) is 200 μ m. Data are expressed as mean \pm SEM ($n = 3$). [#] $P < 0.05$ denotes a statistical difference as compared to the vehicle group. ^{*} $P < 0.05$ denotes a statistical difference as compared to the CCl₄ alone-treated group. SIL, silymarin; R, resveratrol; and P, piceatannol.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

