Green Chemistry



CORRECTION

View Article Online



Cite this: Green Chem., 2022, 24.

Correction: Tuning the selectivity of electrochemical levulinic acid reduction to 4-hydroxyvaleric acid: a monomer for biocompatible and biodegradable plastics

Francisco W. S. Lucas. (1) **a,b Yuval Fishler** and Adam Holewinski (1) **a,b

DOI: 10.1039/d1gc90130c

rsc.li/greenchem

Correction for Tuning the selectivity of electrochemical levulinic acid reduction to 4-hydroxyvaleric acid: a monomer for biocompatible and biodegradable plastics' by Francisco W. S. Lucas et al., Green Chem., 2021. 23. 9154-9164. DOI: 10.1039/D1GC02826J.

An incorrect version of Fig. 2(b) was included in the original publication. This figure had the x-axis shifted erroneously by 100 wavenumbers. The following version of Fig. 2(b) shows the correct x-axis and replaces Fig. 2(b) within the original manuscript.

The peak at 1520 cm⁻¹ was also mis-referenced as 1620 cm⁻¹ in the original text, section 3.2. Potential effects, discussing the figure. The corrected sentence is as follows:

"This experiment shows the formation of (detectable) lead hydride starting around -1.60 V vs. RHE, evidenced by the appearance of a peak at about 1520 cm⁻¹ associated with a Pb-H stretch."




Fig. 2 (b) In situ ATR-FTIR spectra for Pb film polarized at different potentials from -0.05 V to -1.90 V vs. RHE, with steps of 150 mV, in 0.1 M KHCO₃ + 0.1 M KClO₄ at 20 °C.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aDepartment of Chemical and Biological Engineering, University of Colorado, Boulder, CO 80303, USA. E-mail: adam.holewinski@colorado.edu, willlucas@yahoo.com.br

^bRenewable and Sustainable Energy Institute, University of Colorado, Boulder, CO 80303, USA