

Cite this: *Nanoscale*, 2021, **13**, 1365

Correction: Electrostatically modulated magnetophoretic transport of functionalised iron-oxide nanoparticles through hydrated networks

 Stephen Lyons,^a Eoin P. Mc Kiernan,^b Garret Dee,^c Dermot F. Brougham*^b and Aoife Morrin*^a

DOI: 10.1039/d0nr90262d

rsc.li/nanoscale

 Correction for 'Electrostatically modulated magnetophoretic transport of functionalised iron-oxide nanoparticles through hydrated networks' by Stephen Lyons *et al.*, *Nanoscale*, 2020, **12**, 10550–10558, DOI: 10.1039/D0NR01602K.

This erratum relates to Fig. 3 and 4 in the published manuscript, where the labels of both y-axes as given are incorrect. The label given in the published manuscript for both figures is $v_{\text{exp}}/d_{\text{hyd}}$. This should be corrected to $v_{\text{exp}} \cdot d_{\text{hyd}}$ and the units should be $\text{mm}^2 \text{h}^{-1}$. The numerical values for the data are correct and remain unchanged (hence, it is a mis-labelling). The same labelling error ($v_{\text{exp}}/d_{\text{hyd}}$ in place of $v_{\text{exp}} \cdot d_{\text{hyd}}$) was made in two places in the text (page 10555 in the "Electrostatic effects on magnetophoretic mobility" section). The scaling of the v_{exp} values in this way (as a product with d_{hyd}) is consistent with expectation, as formulated by eqn (1). Hence, there is no change to the interpretation of the results or any of the conclusions.




Fig. 3 Normalised magnetophoretic velocities, $v_{\text{exp}} \cdot d_{\text{hyd}}$, for PEG1000, citrate-, and arginine-MNP suspensions through the different classes of agarose–H₂O (Aga/Low, Aga/Med, Aga/High) (0.3% w/v). Error bars are included for all functionalised MNPs. The $v_{\text{th}} \cdot d_{\text{hyd}}$ values are represented as black bars.

^aInsight SFI Research Centre For Data Analytics, National Centre for Sensor Research, School of Chemical Sciences, Dublin City University, Ireland.

E-mail: aoife.morrin@dcu.ie

^bSchool of Chemistry, University College Dublin, Ireland

^cSchool of Chemistry, Trinity College Dublin, Ireland






Fig. 4 Normalised magnetophoretic velocities, $v_{exp} * d_{hyd}$, for PEG1000-, arginine- and citrate-MNP suspensions through agarose-PBS (Aga/High, 0.3% w/v). MNP suspensions and agarose gels were prepared in PBS buffer to give IS of 0, 0.0014, 0.007 and 0.014 at pH 7.0. The $v_{th} * d_{hyd}$ values are represented as black bars.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

