



Cite this: *Nanoscale*, 2021, **13**, 9040

## Correction: *In situ* TEM measurement of activation volume in ultrafine grained gold

Saurabh Gupta,<sup>a</sup> Sandra Stangebye,<sup>a</sup> Katherine Jungjohann,<sup>b</sup> Brad Boyce,<sup>b</sup> Ting Zhu,<sup>a</sup> Josh Kacher<sup>a</sup> and Olivier N. Pierron<sup>\*a</sup>

DOI: 10.1039/d1nr90099d


rsc.li/nanoscale

Correction for '*In situ* TEM measurement of activation volume in ultrafine grained gold' by Saurabh Gupta *et al.*, *Nanoscale*, 2020, **12**, 7146–7158, DOI: 10.1039/D0NR01874K.

The authors have found an error in calculating the true activation volume using eqn (8):

$$V^* = \sqrt{3}kT \frac{\ln(\dot{\epsilon}_{i2}/\dot{\epsilon}_{f1})}{\Delta\sigma_{12}} = \sqrt{3}kT \frac{\ln(\dot{\sigma}_{i2}/\dot{\sigma}_{f1})}{\Delta\sigma_{12}} \quad (8)$$

A base 10 logarithm was mistakenly employed in that equation, instead of the natural logarithm, resulting in the  $V^*$  values shown in the original paper being lower by a factor of 2.3 compared to the correct values. The two figures below (Fig. 7(c) and (d)) show the corrected values. The logarithmic fit for eqn (5) was done correctly in the original paper, hence the apparent activation volumes,  $V_{a3}$ , are correct.



**Fig. 7** *In situ* TEM repeated stress relaxations of a 100 nm-thick Au microspecimen. (c) and (d) corresponding  $V^*$  versus stress and plastic strain, respectively.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup>G.W. Woodruff School of Mechanical Engineering, Georgia Institute of Technology, Atlanta, GA 30332-0405, USA

<sup>b</sup>Sandia National Laboratories, P.O. Box 5800, Albuquerque, NM 87185, USA. E-mail: [olivier.pierron@me.gatech.edu](mailto:olivier.pierron@me.gatech.edu)

