


Organic & Biomolecular Chemistry

Volume 19
Number 2
14 January 2021
Pages 287-488

rsc.li/obc



ISSN 1477-0520

REVIEW ARTICLE

Oleg N. Chupakhin, Bert U. W. Maes *et al.*
C(sp²)-H functionalization in non-aromatic azomethine-
based heterocycles



Cite this: *Org. Biomol. Chem.*, 2021, **19**, 297

Received 30th July 2020,
Accepted 11th September 2020
DOI: 10.1039/d0ob01580f
rsc.li/obc

C(sp²)-H functionalization in non-aromatic azomethine-based heterocycles

Alexey A. Akulov,^a Mikhail V. Varaksin,^{a,b} Pieter Mampuy,^c Valery N. Charushin,^{a,b} Oleg N. Chupakhin^{b,*a,b} and Bert U. W. Maes^{b,*c}

Direct C(sp²)-H functionalization of the endocyclic azomethine and aldonitrone moieties in non-aromatic aza-heterocycles has established itself as a promising methodology over the last decade. Transition metal-catalyzed cross-coupling reactions, α -metalation-electrophile quenching protocols, and (metal-free) nucleophilic substitution of hydrogen reactions (S_NH) are the major routes applied on cyclic imines and their derivatives. In this overview, we show the tangible progress made in this area during the period from 2008 to 2020.

1. Introduction

Nitrogen-containing heterocycles certainly belong to one of the most important classes of organic compounds, and their significance for the existence of any living system can hardly be overestimated. The accumulation of knowledge on these structures, which are widely spread in nature, has enhanced the

progress in organic chemistry, as well as in other scientific fields, particularly in biomolecular chemistry, biology, medicinal chemistry, and materials science.¹ Over the past century, a large number of synthetic methods to modify azaheterocyclic scaffolds have been elaborated, allowing one to obtain novel, often quite complicated molecules, useful for the direct synthesis of fine chemicals (e.g., pharmaceuticals or agrochemicals), dyes, and other industrially important compounds.² Moreover, since the earliest stage of heterocyclic chemistry, azaaromatic compounds have been considered as very attractive structural motifs, based on their versatile practical applications. The synthesis of these compounds is generally realized through lengthy multi-step approaches.³ With the increasing awareness and importance of green chemistry,⁴ che-

^aDepartment of Organic & Biomolecular Chemistry, Ural Federal University, 19 Mira Str., 620002 Ekaterinburg, Russia. E-mail: chupakhin@ios.uran.ru

^bInstitute of Organic Synthesis, Ural Branch of the Russian Academy of Sciences, 22 S. Kovalevskaya Str., 620990 Ekaterinburg, Russia

^cDivision of Organic Synthesis, Department of Chemistry, University of Antwerp, Groenenborgerlaan 171, B-2020 Antwerp, Belgium. E-mail: bert.maes@uantwerpen.be



Alexey A. Akulov

transition metal-catalyzed and metal-free cross-dehydrogenative coupling approaches.

Alexey A. Akulov obtained his MSc degree in Chemistry at Ural Federal University in 2020. In 2016, he joined the Department of Organic & Biomolecular Chemistry at the same university, where he is currently working as a research engineer. His scientific activities are focused on the C-H functionalization methodology development with regard to azaheterocyclic compounds and, in particular, on the elaboration of both novel



Mikhail V. Varaksin

(hetero)aromatic and non-aromatic substrates in the design of novel heterocyclic derivatives of nitroxide radicals, calixarenes, carboranes, and also advanced materials for molecular electronics.

Mikhail V. Varaksin graduated from the Ural State Technical University in 2008. In 2010, he earned his PhD in Organic Chemistry. In 2011, he joined the Department of Organic & Biomolecular Chemistry at Ural Federal University as an Associate Professor. His current research interests include the development of novel pot, atom, and step economical approaches based on the methodology of C-H bond functionalization in both



mists need to develop alternative efficient synthetic sequences towards azaheterocyclic compounds in accordance with the principles of green chemistry encompassing the pot, atom, and step-economical (PASE) philosophy.⁵ In particular, the direct functionalization of C(sp²)-H bonds has gained significant interest of the scientific community as a general approach to construct C-C bonds between two pre-designed structural units.⁶ Such a methodology has the potential to perform better with respect to the principles of green chemistry in comparison to the very powerful classical cross-coupling reactions, which revolutionized the fine chemicals and specialty chemicals synthesis at both discovery and development levels.⁷ However, regioselectivity problems, directing groups, and requirements of specific [(super)stoichiometric] additives or solvents can hamper its green potential, so a holistic green metrics analysis objectively comparing the different processes, *i.e.* direct C(sp²)-H bond functionalization *versus* cross-coupling, will always be required for every specific case.^{4,8} This does not only involve the C-C bond forming reaction itself, but also how the reactants are synthesized.

The progress in exploiting this methodology for the direct C(sp²)-H functionalization of non-aromatic azaheterocycles proved to be not so straightforward and well-studied in comparison with the field of (hetero)aromatic chemistry. However, in the last decade, significant headway in C(sp²)-H functionalization of non-aromatic azaheterocycles has been made. Interestingly, most compounds in those studies contain an endocyclic azomethine (*i.e.* -C=N-) moiety. Cyclic azomethine derivatives of both natural and synthetic origin are known to exhibit profound biological activities or other practically useful properties.⁹⁻¹³ For instance, the cyclic aldimine *nectrisine* (Fig. 1a) isolated from the fungus *Nectria lucida* was found to induce the expression of antigen Ia, thus enabling the restor-

ation of the immune response depressed by immunosuppressive factors of tumors. *Nectrisine* could also be used for the treatment of type 2 diabetes, since it exhibits α -glucosidase inhibitory activity.⁹ *Koranimine* (Fig. 1b), another natural product with a cyclic imine moiety, belongs to the family of peptides obtained by non-ribosomal biosynthesis. Compounds of this type are likely to exhibit a wide spectrum of biological activities ranging from antibiotic to antitumor effects.¹⁰ Besides that, a number of cycloimines of natural origin are known for their fast-acting toxic effects; the latter include spiricyclic compounds secreted by marine shellfish, (*e.g.*, gymnodimine A, Fig. 1c).¹¹

As for compounds obtained synthetically, one has to highlight the family of psychoactive benzodiazepines used broadly in medicinal practice for the treatment of anxiety disorders, epilepsy, insomnia, and withdrawal syndrome (Fig. 1d shows diazepam, one of the most famous benzodiazepine tranquilizers). Another class of compounds that deserves special attention is the cyclic nitrones, many of which proved to be effective spin traps, used for detection and identification of free radicals,¹² and also potential therapeutic agents, particularly nitric oxide donors.¹³ A classic example of a cyclic nitron is 5,5-dimethylpyrroline *N*-oxide (DMPO, Fig. 1e).

Taking into account the versatile applications of cyclic imine derivatives, there is no doubt that the development of efficient synthetic approaches towards such compounds is of importance for modern organic chemistry. In this overview, we discuss in detail the novel approaches based on C(sp²)-H bond functionalization in non-aromatic azomethine-containing azaheterocycles that have been developed over the last decade. The reported methods can be classified into three general categories:



Pieter Mampuy

Pieter Mampuy obtained his PhD in Organic Chemistry from UAntwerpen and the Vrije Universiteit Amsterdam under the supervision of Profs Bert Maes and Romano Orru in 2017. He also worked at Likat in Germany with Prof. Matthias Beller. In 2018, he was appointed as a post-doctoral researcher at UAntwerpen. His second post-doc is in collaboration with the company Covestro. His current research

interests include base metal-catalyzed multicomponent chemistry, organometallics, sulfur chemistry, photoredox catalysis and green metrics analysis.



Valery N. Charushin

Valery N. Charushin is a full member of the Russian Academy of Sciences (RAS), Professor at the Department of Organic and Biomolecular Chemistry at Ural Federal University, and Head of Postovsky Institute of Organic Synthesis of the Ural Branch of RAS. Prof. Charushin graduated with honors from Ural Polytechnical Institute in 1973. In 1976, he earned his PhD in Organic Chemistry. In 1981-1982, he joined the

research group of Prof. Henk van der Plas (Wageningen University, Wageningen, Netherlands), where he was working on the ring transformations of azines. In 1987, he received a Doctor of Science degree in Organic Chemistry. His current research interests include novel methods in organic synthesis, heterocyclic and medicinal chemistry.






Fig. 1 Selected examples of cyclic azomethine derivatives of specific interest for their application.

- transition metal-catalyzed C(sp²)-H bond functionalization of azomethine moieties;
- C(sp²)-H metalation of azomethine moieties (followed by treatment with electrophiles);
- nucleophilic substitution of hydrogen (S_N^H) at the sp²-carbon in azomethine moieties.

We should note that this survey does not embrace the C(sp²)-H activation in planar azomethine-based heterocycles, containing an aromatic resonance structure or aromatic tautomer, which can at least partly be regarded as aromatic ones

(e.g., quinoxalin-2(1*H*)-ones, quinazolin-4(3*H*)-ones, pyrazin-2(1*H*)-ones, etc.). The vigorously developing chemistry of some of these valuable compounds has recently been highlighted in other review articles,¹⁴ and it keeps updating dynamically with new functionalization examples.¹⁵

1.1. Transition metal-catalyzed C(sp²)-H bond functionalization of azomethine moieties

The discovery of transition metal-catalyzed C-C bond forming cross-coupling reactions in the sixties and seventies has had a



Oleg N. Chupakhin

Oleg N. Chupakhin is a full member of the Russian Academy of Sciences, Professor and ex-Head of the Department of Organic & Biomolecular Chemistry at Ural Federal University, and Chief Researcher at Postovsky Institute of Organic Synthesis of Ural Branch of RAS. Prof. Chupakhin graduated with honors from Ural Polytechnical Institute in 1957. He earned his PhD in Organic Chemistry in 1962 and Doctor of Science

degree in 1976. Current research interests: new methodologies in organic synthesis, structural analysis of organic compounds, mechanisms of organic reactions, heterocyclic chemistry, green chemistry, industrial chemistry, environmental chemistry, medicinal chemistry, chemosensors, molecular recognition and supramolecular chemistry.




Bert U. W. Maes

Bert U. W. Maes obtained his PhD in Organic Chemistry at UAntwerp and subsequently received a Post-Doctoral Fellowship of the Science Foundation (FWO-Flanders) in Belgium. He worked at the École Normale Supérieure in Paris (mechanisms in catalysis) with Prof. Anny Jutand (CNRS). Bert Maes was appointed Assistant Professor (Docent) in the Department of Chemistry at UAntwerp in 2003 and currently

holds a Full Professorship (Gewoon Hoogleraar) of Organic Chemistry at UAntwerp; since 2009 he has been a Research Professor. In 2019 he was appointed as Collen-Francqui Research Professor by the Francqui Foundation. In 2015–2016 he acted as chairman of the Department of Chemistry. His research interests cover the fields of heterocyclic chemistry, organometallic chemistry and homogeneous catalysis with a special focus on the development of sustainable synthetic methodology. The research in his group involves base metal catalysis, aerobic oxidation, strong bond activation, renewables, and green metrics analysis. His research group partakes in one of the Excellence Centers of UAntwerp, i.e. CASCH. He is an editor of Topics in Heterocyclic Chemistry and an editorial board member of SynOpen and Advances in Heterocyclic Chemistry.





Scheme 1 Two possible modes for the TM-catalyzed C(sp²)-H bond functionalization of azomethines.

large impact on the disconnections currently made in organic synthesis. The outstanding achievements of R. Heck, A. Suzuki, E. Negishi, J. Stille, K. Sonogashira, and other distinguished scientists have contributed greatly to the progress in the chemistry of the functionalization of (hetero)aromatics.¹⁶ The first results in the ‘conventional’ cross-coupling chemistry involving AG-substituted cyclic azomethines (AG = halogen, SnAlk₃, or other auxiliary groups) were reported in the early 1980s,¹⁷ whereas the analogous metal-catalyzed transformations exploiting the reactivity of the endocyclic azomethine C(sp²)-H bond remained unexplored until the late 2000s (*vide infra*).¹⁸ Notably, the C-H functionalization methodology can be realized here in two modes.


The first approach involves a new carbon-carbon bond formation through the reaction of the imine C(sp²)-H with a (pseudo)halogenated coupling partner (Scheme 1a), while the second one encompasses the formation of a new bond *via* double C-H bond activation. The latter is also known as cross-dehydrogenative coupling (CDC), requiring an oxidant to obtain a catalytic process (Scheme 1b).¹⁹

1.2. C(sp²)-H metalation of azomethine moieties

The inherent insufficient nucleophilicity of the carbon center prevents direct C-H functionalization of azomethine (R-N=CH-R') moieties *via* electrophilic agents. Therefore, C-C coupling reactions can only be performed by activating the azomethine C(sp²)-H bond first *via* metalation through deprotonation, *e.g.*, using organolithium or organomagnesium reagents,²⁰ followed by treatment with electrophiles (Scheme 2). The pioneering research studies in the field of electrophilic transformations of cyclic azomethine derivatives



Scheme 2 Metalation of the C(sp²)-H bond of a cyclic azomethine derivative (*i.e.* nitron **1**) followed by electrophilic substitution.



Scheme 3 Selected examples of the coupling reaction of α-lithiated aldonitrones with electrophilic reactants.


were carried out by Voinov & Grigor'ev *et al.* in the late 1990s.²¹ They succeeded in developing α-lithiation of cyclic aldonitrones **3** and **7a** generating the corresponding lithium salts (*e.g.*, **3-Li**), which were reacted with electrophilic reactants, such as aldonitrones **3** as well as ketones/aldehydes **5** (Scheme 3).

1.3. Nucleophilic substitution of hydrogen (S_N^H) in azomethine moieties

Nucleophilic substitution of hydrogen (S_N^H) has proved to be an efficient synthetic tool to modify (hetero)aromatic compounds with nucleophilic reactants since the seventies, and it was successfully exploited during the next four decades by many researchers.^{6a,22} Although a few scattered examples employing this strategy for non-aromatic unsaturated heterocycles date back to the early eighties,²³ the majority of such S_N^H reactions have only been reported in the 21st century.

A cyclic aldimine derivative is polarized and contains a partial positive charge at the sp²-hybridized carbon, and this center is therefore vulnerable to reaction with nucleophiles (Scheme 4). Nucleophilic substitution of hydrogen proceeds either *via* an ‘Addition-Elimination’ (S_N^H(AE), Scheme 4a) or an ‘Addition-Oxidation’ mechanism (S_N^H(AO), Scheme 4b).^{22h}






Scheme 4 'Addition–Elimination' (a) and 'Addition–Oxidation' (b) mechanisms for the nucleophilic substitution of hydrogen (S_N^H).

They both start with the addition of a nucleophilic reactant (Nu^-) to the azomethine carbon of **11** to form an anionic σ^H -adduct **12**. According to an $S_N^H(AE)$ mechanism, subsequent spontaneous or externally-promoted elimination of the leaving group regenerates the azomethine π -bond. Selection of the elimination promoters determines the specific features of the S_N^H reaction used. On the other hand, *via* an oxidizing reagent, the equilibrium can also be shifted in favor of the S_N^H product **14**, *i.e.* the oxidative $S_N^H(AO)$ mechanism.


Functionalization of azomethine derivatives involving an $S_N^H(AE)$ pathway was not reported until the last decade.²⁴ Notably, this 'Addition–Elimination' strategy has been frequently applied for the modification of (hetero)aromatic substrates, but not for saturated systems.^{22c} In contrast, the alternative oxidative $S_N^H(AO)$ protocols are well documented here, as illustrated by transformations of aldonitrones **15**, **16a–b** (Scheme 5).²⁵ Transition metal oxides derived from manganese (Scheme 5b and c), nickel (Scheme 5a), lead,^{23a} and mercury^{23b} are commonly used as oxidizing agents. Oxidations employing air^{23c} or hydrogen peroxide,^{25a} supplemented with catalytic additives, which are more sustainable,²⁶ were also exploited.

2. $C(sp^2)$ –H functionalization in cyclic aldimines and related compounds

The first report describing the direct arylation of a cyclic imine was published in 2008.¹⁸ A research team from the University of California (Berkeley) introduced a phenyl moiety at the 2-position of 4,4-dimethyloxazoline (**17a**), a non-aromatic compound (Scheme 6), in the presence of a rhodium source and




Scheme 6 Rh^I -Catalyzed $C(sp^2)$ –H arylation of 4,4-dimethyloxazoline (**17a**).



Scheme 5 Selected examples of $S_N^H(AO)$ reactions on cyclic azomethines.







Scheme 7 Proposed mechanism for the Rh^I-catalyzed C(sp²)-H arylation of 4,4-dimethyloxazoline (**17a**).

(*Z*)-1-*tert*-butyl-2,3,6,7-tetrahydrophosphepine (**20**) as a ligand. The catalytic cycle proposed by the authors (Scheme 7) involves the initial formation of dimeric complex **21** from dimeric [RhCl(coe)₂]₂ and bidentate phosphepine ligand **20**. Dissociation of this complex and coordination with the imidate nitrogen of 4,4-dimethyloxazoline (**17a**) are likely to promote the formation of intermediate **22** in a similar manner as described for heteroaromatic substrates with rhodium complexes, such as RhCl(PCy₃)₂.²⁷ A low-valent electron-rich carbene-rhodium complex **23** is then generated *via* a C-H activation/tautomerization process.²⁷ Subsequent oxidative addition with phenyl bromide **18a** delivers a (phenyl)(carbene) rhodium complex **24**. Elimination of HBr could then either occur *via* the amine base or *via* an intramolecular process, affording intermediate **25**. Finally, reductive elimination results in the formation of **19a** and regeneration of the rhodium catalyst.


In 2011, the Ackermann group reported a similar C(sp²)-H arylation of oxazolines **17** involving Pd rather than Rh catalysis (Scheme 8).²⁸ A palladium complex **26** formed from Pd(OAc)₂ and secondary phosphine oxide 'pre-ligands' were found to be suitable (Scheme 9). A series of arylated oxazolines (**19**) were



Scheme 8 Pd-Catalyzed C(sp²)-H arylation of oxazolines (**17**).



Scheme 9 Preparation of Pd catalyst (**26**) for C(sp²)-H arylation of oxazolines (**17**).




Scheme 10 Cu-Catalyzed C(sp²)-H arylation of benzotriazepinones (**29**).

successfully obtained, varying substituents in both the aryl and oxazoline moieties.

Another report on the successful C(sp²)-H arylation of non-aromatic heterocycles was published by the Ellman group.²⁹ Herein, benzotriazepinones **29**, which can be considered as cyclic hydrazones, were subjected to Cu-catalyzed C-H functionalization with a range of aryl iodides (**30**, Scheme 10). Being close analogues of pharmaceutically privileged benzodiazepines, these azaheterocyclic derivatives are of interest as biologically active substances. The reaction conditions were initially similar to those proposed by Daugulis for the arylation of aromatic C-H bonds (polyfluoroarene or heteroaromatic) with aryl iodides and bromides.³⁰ However, under those conditions, arylated benzotriazepinone **31c** was only obtained in 40% yield. To reach sufficient conversion, a longer reaction time (12 h) was required than that initially reported (10–30 min). Further optimization of the reactant ratio (**29** : **30**) afforded a range of 5-aryl-substituted benzotriazepinones **31** in rather good yields (Scheme 10).


Palladium in combination with copper catalysis was explored by the scientists from the University of Rouen (France). In 2015, they reported the direct arylation and olefination of imidazolones **32** (Scheme 11).³¹ Heterocycles of this class are present in nature and exhibit therapeutic and fluorescence properties.³¹ Aryl or alkenyl halides (**33** and **35**) were used as coupling partners with 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) as a base. It is noteworthy that such arylations and olefinations required different solvents (DMF or toluene) and phosphine ligands (PPh₃ or P(*o*-Tol)₃) to obtain appropriate yields of products (**34** and **36**, see Scheme 11). The potential of this strategy was illustrated for the synthesis of imidazolone-containing fatty acid synthase (FAS) inhibitors.³¹ This group also extended this approach for the C(sp²)-H functionalization of 4-arylidene imidazolones, enabling a library of GFP and Kaede protein-type fluorophores.³²






Scheme 11 Pd-Catalyzed direct arylation and olefination of 4,4-dialkyl-1*H*-imidazol-5(4*H*)-ones (**32**).

In 2012, researchers from the Zelinsky Institute of Organic Chemistry in Russia reported an atypical *C*-alkylation of 2-pyrazolines **38** with 2-(hetero)arylcyclopropane-1,1-dicarboxylates **37** in the presence of superstoichiometric gallium trichloride.³³ A *C,N*-bis-alkylation product **39** was obtained as a major compound in refluxing dichloromethane (Scheme 12). However, performing this reaction at room temperature led to a greater proportion of byproducts (**40** and **41**). Interestingly, other Lewis acids, such as scandium and ytterbium triflates, predominantly resulted in the formation of cycloaddition product **41**.³⁴ This cycloadduct **41** is generated *via* an interaction of the *N*(2)-atom in 2-pyrazoline **38** with the electrophilic center of cyclopropane **37** after initial activation with a Lewis acid (Scheme 13). In contrast, activation of both reactants (**37** and **38**) seems to occur at higher temperature, which redistributes the electron density within the pyrazoline ring. This enhances the nucleophilicity at the *C*(3)-atom and results in the formation of 1-pyrazoline *C*(3)-adduct **42**. Subsequent tautomerization *via* migration of the hydrogen from *C*(3)-




Scheme 12 GaCl₃-mediated coupling of 5-substituted 2-pyrazolines (**38**) with 2-arylcyclopropane-1,1-dicarboxylates (**37**).



Scheme 13 Proposed mechanism for a GaCl₃-mediated coupling of 5-substituted 2-pyrazolines (**38**) with 2-arylcyclopropane-1,1-dicarboxylates (**37**).


N(1)-position, followed by interaction with another equivalent of cyclopropane (**37**) generates the bis-alkylation product **39**.

While analyzing the data of various studies focused on the *C*(sp²)-*H* functionalization of cyclic azomethines, the tendency to apply compounds of the aldonitrone family (featuring an additional *N*-oxide on the azomethine) as *C*-*H* substrates for these transformations became apparent.³⁵ They are easily accessible *via* a multitude of procedures (*e.g.*, oxidation reactions (Scheme 14a), condensation reactions of *N*-monosubstituted hydroxylamines (Scheme 14b), or reactions of oximes with electrophiles (Scheme 14c)).³⁵ The *N*-oxide group of nitrones has been exploited by various research



Scheme 14 General routes towards nitrones.³⁵






Scheme 15 Formal C(sp²)-H alkenylation of imidazolones (**43**) via the N-oxide.



groups as an elegant directing group for C(sp²)-H functionalization of azomethines.^{6c} In particular, significant attention has been paid to imidazolone N-oxides.

In 2010, the Merck Research Laboratory in the USA achieved the synthesis of 2-alkenylimidazolones (**45**) via a 1,3-dipolar cycloaddition (1,3-DC) of alkenes (**44**) to cyclic nitrones (**43**, Scheme 15).³⁶ The procedure involves a one-pot three-step process (Scheme 16). Initially, the cycloadduct (**46**) is formed from imidazolone N-oxide (**43**) and alkene (**44**) in refluxing ethanol. After cooling to ambient temperature, subsequent treatment with aqueous NaOH (1 N) (base-induced N-O bond cleavage) and HCl (1 N) (acid-induced water elimination) delivered the target compound **45** in good to excellent yields. Various aromatic and aliphatic alkenes **44** were compatible with the developed reaction conditions.

Two years later, researchers from Albany Molecular Research Inc. reported the first Pd-catalyzed arylation of aldonitrones **47** with aryl bromides (**18**, Scheme 17).³⁷ The reaction conditions were similar to those previously reported by Fagnou for the arylation of heteroaromatic N-oxides.³⁸ A cross-coupling between **47** and phenyl bromide (**18a**) successfully delivered **48a**; however, only a moderate yield of 49% was obtained (Scheme 17a). A significant improvement in yield was achieved via protection of the imidazolone moiety with either the *tert*-butoxycarbonyl (Boc) or 2,4-dimethoxybenzyl (DMB) protecting group (**48b-c**). With the optimal DMB protecting group in hand, the authors were able to arylate aldonitronone **47a** with various aryl bromides (**18**) in good to excellent yields (Scheme 17b). This approach was also successfully applied for the synthesis of GSK2137305 **56** (Scheme 18), a known selective inhibitor of glycine transporter type 1 (GlyT1), exhibiting effectiveness for neurological and neuropsychiatric dis-



Scheme 16 A proposed mechanism for alkenylated imidazolone (**45**) synthesis from the corresponding imidazolone N-oxide (**43**).





Scheme 17 (a) Effects of the N-substituent in lactam **47** on the direct arylation reaction yield; (b) Pd-catalyzed coupling of 4-DMB-protected imidazol-3-one 1-oxide (**47a**) with aryl bromides (**18**).

orders.³⁹ The procedure nicely illustrates the N-oxide directing group, which can be easily introduced via oxidation of cyclic amine **53** with *m*-CPBA (*vide supra*) and removed by employing PBr₃, yielding cyclic azomethine target compound **56**. Other reagents to cleave N-oxides are also available (e.g., CuI/Zn system, PⁿBu₃, Mo(CO)₆, InCl₃, etc.).³⁵

In 2011, the synthesis of novel pseudouridine-like compounds (e.g., **62**), bearing 1-pyrroline-1-oxide moieties instead of sugar residues, was reported by Koszytkowska-Stawińska (Scheme 19).⁴⁰ Pseudouridine (ψ -uridine) is a natural C-nucleoside found in various types of RNA. This compound is supposed to play a key role in fine-tuning the functions of RNA, and it contributes likely to the preservation of the RNA structure during translation.⁴¹ Moreover, ψ -uridine is also active against ionizing radiation and chemical mutagens.⁴² These protective effects are associated with the ability of the nucleoside to trap free radicals formed by mutagens. The derivatives of 1-pyrroline 1-oxide are efficient spin-traps for EPR analysis (the most prominent example is 5,5-dimethyl-1-pyrroline N-oxide, DMPO).⁴³ Many azaheterocycles, which are structurally similar to the pyrroline scaffold (pyrrolidines, oxazoles, isoxazoles, etc.), are able to mimic a furanose residue. The replacement of the ribose residue in pseudonucleosides with a 5,5-bis(hydroxymethyl)-1-pyrroline-1-oxide was therefore envisioned to improve the antimutagenic activity. To obtain such modified ψ -uridines, the authors first synthesized *O,O'*-cyclohexylidene protected 5,5-bis(hydroxymethyl)-1-pyrroline 1-oxide (**58**) as an aldonitronone precursor. Subsequent addition of either C(5)-lithiated 2,4-dimethoxypyrimidine (**57-Li**, Scheme 19a) or the analogous Grignard reactant (**57-MgBr**, Scheme 19b), followed by copper-catalyzed aerobic oxidation of the adduct and acetal deprotection of the resulting nitronone **59** with NaOAc in AcOH (Scheme 19c), afforded the desired pseudouridine aza-analogues **62**. The coupling with the



Scheme 18 The synthesis of GSK2137305 (**56**).Scheme 19 Synthesis of pseudouridine aza-analogue **62**.

lithium derivative (**57-Li**) was less successful, as nitron dimerization was observed as the major process, probably due to deprotonation at the C(2)-H of 1-pyrroline 1-oxide (**58**).

In 2012, Chavant & Blandin *et al.* reported an efficient approach for Pd- or Pd/Cu-catalyzed direct arylation of cyclic aldonitrones (**63–65**),⁴⁴ which is related to the C–C coupling of







Fig. 2 Cyclic and acyclic nitrones studied: MiPNO (**63**); DMPO (**64**); 3,3-dimethyl-3,4-dihydroisoquinoline 2-oxide (**65**); PBN (**66**); glyoxylic nitrone (**67**).

imidazolone *N*-oxides (**47**) with aryl bromides (**18**) reported by the AMRI research group (Scheme 17). The cyclic aldonitrone 2-isopropyl-1,2-dimethyl-5-oxo-2,5-dihydro-1*H*-imidazole 3-oxide (**63**, MiPNO, Fig. 2) is a versatile precursor for the synthesis of α -amino acids (via 1,3-dipolar cycloaddition reactions with alkenes followed by cleavage of the *N*-O bond)⁴⁵ and was one of the key studied substrates. Pivalic acid (20 mol%) was an essential additive for a successful cross-coupling with aryl bromides (**18**), enhancing the yield dramatically (Scheme 20a).

Alternatively, CuBr·DMS (5 mol%) with 1,10-phenanthroline (Phen, 5 mol%) as a ligand proved also to be an effective additive to achieve quantitative conversion of **63**. It is worth noting that no synergistic effect was observed, when these two additives (PivOH and CuBr·DMS) were used simultaneously. Interestingly, in the absence of the palladium catalyst, a catalytic amount of CuBr·DMS enabled the conversion of aldonitrone **63** into the corresponding homocoupling product (i.e. a nitrone 2,2'-dimer without one of two *N*-oxide functions). Employing dibromo(hetero)arenes (**69**) with two equivalents of aldonitrone (*S*-**63**) afforded successfully bis-nitronyl products **70** in good yields (Scheme 20b). The reaction is not specific to MiPNO (**63**), since a range of other heterocyclic non-aromatic aldonitrones can be employed in this cross-coupling reaction, as illustrated for DMPO (**64**, Scheme 21a) and 3,3-dimethyl-3,4-dihydroisoquinoline 2-oxide (**65**, Scheme 21b). The reaction with DMPO is interesting due to the presence of potentially




Scheme 21 Pd-Catalyzed direct arylation of DMPO (**64**, a) and 3,3-dimethyl-3,4-dihydroisoquinoline 2-oxide (**65**, b).

labile allylic H-atoms relative to the imine double bond (Scheme 21a). In contrast, acyclic nitrones, such as *N*-*tert*-butyl- α -phenylnitrone (**66**, PBN) or glyoxylic nitrone (**67**), were not compatible with this kind of arylation (Fig. 2). It has been suggested that these cross-coupling processes are likely to proceed via the so-called concerted metalation–deprotonation (CMD) mechanism,⁴⁶ explaining the beneficial effect of PivOH, described previously for Pd-catalyzed direct arylations of (hetero)aromatic substrates (i.e. for azine *N*-oxides). It is worth noting that this method has been used for the synthesis of chiral imidazolidin-4-one nitroxides starting from enantiopure **63**, and they are regarded as potentially active catalysts for aerobic oxidation of alcohols.⁴⁷ In particular, catalytic application of such compounds in enantiopure form allows one to achieve atroposelective desymmetrization during a diol oxidation process.⁴⁷


Special attention has recently been paid to C–H functionalization of related non-aromatic 2*H*-imidazole 1-oxides (**7**) featuring an extra unsaturation. In 2012, our research group reported the first successful cross-coupling reactions of 2,2-dialkylated 4-phenyl-2*H*-imidazole 1-oxide (**7b**) with 1,2,4-triazine **73** (Scheme 22) or its *N*-oxide **77** (Scheme 23).⁴⁸ The reaction mechanism includes the initial lithiation of the azomethine C–H bond in aldonitrone **7b**, which is assisted by the *N*-oxide (Directed 'ortho' Metalation (DoM)⁴⁹), followed by reaction with 1,2,4-triazine **73** or its *N*-oxide **77** to afford NLi- or OLi-adducts (**74** or **78**). These adducts are subsequently protonated with water into the corresponding NH- or OH-intermediates (**75** or **79**). Finally, rearomatization of the dihydrotriazine ring in **75/79** occurs either through spontaneous elimination of water, affording the coupling product without the *N*-oxide function (**76/80**), or by oxidation with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) to give the corresponding *N*-oxide **81** (Scheme 23). It is worth noting that cleavage of the *N*-oxide moiety in the triazine fragment of **77** occurs via the 'Addition–Elimination (AE)' protocol, while the *N*-oxide moiety remains untouched via the 'Addition–Oxidation (AO)' protocol.

Our group also reported the coupling of 2*H*-imidazole 1-oxide **7b** with 1,10-phenanthroline (**82a**), quinoxaline (**82b**),




Scheme 20 Pd- and Pd-/Cu-Catalyzed direct arylations of MiPNO (**63**).






Scheme 22 Proposed mechanism for the coupling of 2*H*-imidazole 1-oxide **7b** with 1,2,4-triazines (**73**) via DoM lithiation.




Scheme 23 Proposed mechanism for the coupling of 2*H*-imidazole 1-oxide **7b** with 1,2,4-triazine 4-oxide **77** via DoM lithiation.

and their *N*-oxides (**85a–b**) as electron π -deficient azaheterocyclic substrates (Schemes 24 and 25).⁵⁰ Interestingly, the *N*-oxide moiety is eliminated from the imidazole ring in the (AE)-reac-



Scheme 24 Proposed mechanism for the coupling of 2*H*-imidazole 1-oxide **7b** with azines (**82**) via DoM lithiation.




Scheme 25 Proposed mechanism for the coupling of 2*H*-imidazole 1-oxide **7b** with azine *N*-oxides (**85**) via DoM lithiation.


tion of aldonitrone **7b** with quinoline *N*-oxide **85c**, whereas the *N*-oxide fragment of the pyridine ring remains intact (Scheme 25), in contrast to the reaction of nitron **7b** with triazine *N*-oxide **77** (Scheme 23). Clearly, retention of the specific *N*-oxide function in products is controlled by the stability of the tautomeric intermediates (**86a** and **86b**, respectively).




The same substrate **7** can be used as an electrophile rather than as a nucleophile. Our group reported the cross-coupling reaction of *2H*-imidazole 1-oxides (**7a–c**) with electron-rich heteroarenes, such as indoles (**90**),²⁴ pyrroles (**92**)^{51,52} and thiophenes (**93**) (Schemes 26–28).⁵² The coupling of these partners



Scheme 26 Pd^{II}-Catalyzed oxidative coupling of *2H*-imidazole 1-oxides (**7a–c**) with indoles (**90**, a), pyrroles (**92**), and thiophenes (**93**, b).



Scheme 27 Proposed catalytic cycle for the oxidative coupling of *2H*-imidazole 1-oxides (**7a–c**) with pyrroles (**92**) or thiophenes (**93**).




Scheme 28 The S_N^H reaction of *2H*-imidazole 1-oxides **7a–c** with indoles (**90**, a) and pyrroles (**92**, b).

can be done in two different ways. The first approach is based on a Pd^{II}-catalyzed cross-dehydrogenative coupling (Scheme 26). During the first step of the proposed catalytic cycle (Scheme 27), organopalladium intermediate **96/97** is formed *via* the reaction of Pd(OAc)₂ with a heteroaromatic substrate (**92/93**). Palladium in the intermediate **96/97** is hypothesized to be coordinated with the oxygen atom of nitrene **7a–c** *via* a ligand exchange mechanism, leading to the formation of intermediate **98/99**. Subsequent carboxylate-assisted functionalization of the nitrene C(1)–H bond takes place (CMD-process, **TS1/TS2**), generating intermediate **100/101**. Finally, reductive elimination affords the target products **94/95** and Pd⁰. The latter is reoxidized to Pd^{II} with the superstoichiometric oxidant Cu(OAc)₂. Interestingly, pyridine was found to be an essential additive and presumed to play a role of ligand (L_n) stabilizing organometallic intermediates.


The second approach is a *cine*-type reaction based on the S_N^H(AE) process between 2,2-dialkylated 4-phenyl-*2H*-imidazole 1-oxides (**7a–c**) and indoles or pyrroles (**90** or **92**) as coupling partners (Scheme 28). For these transformations, the acylating reagent plays a crucial role, since it reacts with the *N*-oxide moiety of **7a–c** to generate a more electrophilic species **104** and immediately builds in a suitable leaving group (Scheme 29). Simple acetyl chloride can be used. These chloride salts formed then react in electrophilic addition reactions at the C(2)-position of pyrroles (**92**) or the C(3)-position of indoles (**90**), generating the corresponding dihydro compounds (*e.g.*, **105**). Subsequent spontaneous elimination of acetic acid affords the hydrochloride salts of the desired products **102/103**. Notably and in line with the substitution pattern, this method is not compatible with pyrroles (**92**) and indoles (**90**) bearing electron-withdrawing groups, *i.e.* nitro or acetyl groups.

The nucleophile can also be an organometallic reactant (Scheme 30). A challenging example is *ortho*-carboranyl lithium






Scheme 29 Proposed reaction mechanism for the S_N^H reaction of 2H-imidazole 1-oxides (7a-c) with pyrroles (92) (cine reaction).



Scheme 30 S_N^H reactions of 2H-imidazole 1-oxides (7) with *ortho*-carboranyl lithium (106) via S_N^H (AE) and oxidative S_N^H (AO) strategies.

(106). The reaction with 2H-imidazole 1-oxides was reported in 2018.⁵³ The nucleophile is particularly interesting, as organoboron compounds and their heterocyclic derivatives are potential agents for boron neutron capture therapy (BNCT) of cancer.⁵³ Given the nucleophilicity of organolithium species, *N*-oxide activation is in this case not required; in fact, it could competitively react at the carbonyl of the ester introduced, and a reaction with acetyl chloride is merely used as a quencher to rearomatize. Besides eliminative S_N^H (AE), an oxidative S_N^H (AO) strategy keeping the *N*-oxide in the structure is possible (Scheme 30). Very recently, the analogous protocol has also been applied for the direct $C(sp^2)$ -H arylation of 2H-imidazole 1-oxides 7 with another organometallic reactant, pentafluorophenyllithium 110 (Scheme 31).⁵⁴ Some of the resulting coupling products 112/113 are considered as promising push-pull



Scheme 31 S_N^H reactions of 2H-imidazole 1-oxides (7) with pentafluorophenyllithium (110) via S_N^H (AE) and oxidative S_N^H (AO) strategies.

fluorophores to be useful in the design of novel fluorometric sensors.

3. Conclusions and outlook

A systematic overview of $C(sp^2)$ -H functionalization reactions for the modification of cyclic azomethine-based compounds has been performed. As functionalizing reactants, both carbon-centered nucleophiles and electrophiles can be used (furthermore, the options for employing heteroatom-centered synthons gradually appear to become accessible as well,⁵⁵ although this area is still in its infancy). Meanwhile, the use of cyclic aldonitrones as starting azomethine substrates turned out to be prevalent over other related compounds, such as imines or hydrazones. Undoubtedly, this is due to the remarkable properties of the nitron *N*-oxide moiety, which can both act as a directing group and as a leaving group.

Although aldonitrones are shown to react readily with a number of electrophilic and nucleophilic synthons, it seems to be rather odd that the examples of radical $C(sp^2)$ -H functionalization of these compounds are scarcely known at the moment.³⁵ After all, nitrones have become widely applicable exactly due to their radical addition reactions affording relatively stable paramagnetic nitroxides useful for EPR analysis. In this connection, we believe that stepping into the above-mentioned *terra incognita* can be a milestone for the further development of the discussed topic.

Another certainly considerable objective is the sustainability of the process. Though very effective synthetic methodologies have been developed in the past decade, there is still quite a lot to be improved in terms of greenness and compliance with the PASE principles. Several approaches surveyed in



this overview deal with transition metal catalysis. Although catalysis is inherently a green methodology, an aryl bromide or iodide reactant partner, as well as noble metals (sometimes with complex ligands), is typically required. Besides, it often appears to be necessary to use (super)stoichiometric quantities of additives (e.g., bases). Indeed, these reactions are highly valuable for modern organic synthesis and open new horizons compared, for instance, with the conventional 'electrophile-nucleophile' interactions, yet there still remains room for further improvement. For example, the use of base metals as catalysts or cross-dehydrogenative procedures, which are inherently more interesting, considering no prefunctionalization of the reactant is required (though this can pose regioselectivity issues in return). However, if no sustainable oxidant is used, nothing can be gained by employing a cross-dehydrogenative approach, as it is still not suitable for scaling, albeit very useful for discovery research purposes. Oxygen,^{26a} hydrogen peroxide,^{26b} and electric current⁵⁶ are interesting oxidants. Photoredox-catalyzed⁵⁷ and organocatalyzed⁵⁸ C-H functionalizations certainly also hold promise with respect to green chemistry. To the best of our knowledge, they have not yet been applied for C(sp²)-H bond functionalization in cyclic azomethines and aldonitrones. Thus, investigations aimed at further improving the sustainability of the current processes are expected to be the focus of research in the next few years.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The research was financially supported by the Russian Science Foundation (Project No. 20-43-01004), a bilateral Russian Science Foundation (RSF) – Fund for Scientific Research Flanders (FWO) project, and the Francqui Foundation.

References

- (a) *Modern Heterocyclic Chemistry*, ed. J. Alvarez-Builla, J. J. Vaquero and J. Barluenga, Wiley-VCH, Weinheim, 2011; (b) *Green Synthetic Approaches for Biologically Relevant Heterocycles*, ed. G. Brahmachari, Elsevier, Oxford, 2014; (c) J. A. Joule and K. Mills, *Heterocyclic Chemistry*, Blackwell Publishing, Chichester, 5th edn, 2013.
- A. Katritzky, C. Ramsden, J. Joule and V. Zhdankin, *Handbook of Heterocyclic Chemistry*, Elsevier, Amsterdam, 3rd edn, 2010.
- M. Smith, *March's advanced organic chemistry: reactions, mechanisms, and structure*, John Wiley & Sons, Inc., Hoboken, 7th edn, 2013.
- (a) R. A. Sheldon, *Chem. Soc. Rev.*, 2012, **41**, 1437–1451; (b) A. P. Dicks and A. Hent, *Green Chemistry Metrics: A Guide to Determining and Evaluating Process Greenness* (*SpringerBriefs in Molecular Science*), Springer, Cham, 1st edn, 2015; (c) L. Summerton and A. Constandinou, in *Green and Sustainable Medicinal Chemistry: Methods, Tools and Strategies for the 21st Century Pharmaceutical Industry*, The Royal Society of Chemistry, 2016, pp. 41–53; (d) J. H. Clark, in *Green and Sustainable Medicinal Chemistry: Methods, Tools and Strategies for the 21st Century Pharmaceutical Industry*, The Royal Society of Chemistry, 2016, pp. 1–11.
- W. Zhang and W.-B. Yi, *Pot, Atom, and Step Economy (PASE) Synthesis*, Springer, Cham, 2019.
- (a) *Metal Free C-H Functionalization of Aromatics. Nucleophilic Displacement of Hydrogen*, ed. V. Charushin and O. Chupakhin, Springer, Cham, 2014; (b) C.-S. Wang, P. H. Dixneuf and J.-F. Soulé, *Chem. Rev.*, 2018, **118**, 7532–7585; (c) C. Sambigioglio, D. Schönbauer, R. Blicke, T. Dao-Huy, G. Pototschnig, P. Schaaf, T. Wiesinger, M. F. Zia, J. Wencel-Delord, T. Besset, B. U. W. Maes and M. Schnürch, *Chem. Soc. Rev.*, 2018, **47**, 6603–6743; (d) D. J. Abrams, P. A. Provencher and E. J. Sorensen, *Chem. Soc. Rev.*, 2018, **47**, 8925–8967; (e) P. Gandeepan, T. Müller, D. Zell, G. Cera, S. Warratz and L. Ackermann, *Chem. Rev.*, 2018, **119**, 2192–2452; (f) S. Rej, Y. Ano and N. Chatani, *Chem. Rev.*, 2020, **120**, 1788–1887.
- M. Beller, A. Varela-Fernandez, J. G. de Vries, S. M. Wong, C. M. So and F. Y. Kwong, in *Applied Homogeneous Catalysis with Organometallic Compounds*, ed. B. Cornils, W. A. Herrmann, M. Beller and R. Paciello, Wiley-VCH, Weinheim, 3rd edn, 2017, pp. 411–464.
- (a) C. R. McElroy, A. Constantinou, L. C. Jones, L. Summerton and J. H. Clark, *Green Chem.*, 2015, **17**, 3111–3121; (b) S. Abou-Shehada, P. Mampuy, B. U. W. Maes, J. H. Clark and L. Summerton, *Green Chem.*, 2017, **19**, 249–258; (c) E. R. Monteith, P. Mampuy, L. Summerton, J. H. Clark, B. U. W. Maes and C. R. McElroy, *Green Chem.*, 2020, **22**, 123–135.
- H. Kayakiri, K. Nakamura, S. Takase, H. Setoi, I. Uchida, H. Terano, M. Hashimoto, T. Tada and S. Koda, *Chem. Pharm. Bull.*, 1991, **39**, 2807–2812.
- B. S. Evans, I. Ntai, Y. Chen, S. J. Robinson and N. L. Kelleher, *J. Am. Chem. Soc.*, 2011, **133**, 7316–7319.
- A. Otero, M.-J. Chapela, M. Atanassova, J. M. Vieites and A. G. Cabado, *Chem. Res. Toxicol.*, 2011, **24**, 1817–1829.
- C. E. Thomas, D. F. Ohlweiler, A. A. Carr, T. R. Nieduzak, D. A. Hay, G. Adams, R. Vaz and R. C. Bernotas, *J. Biol. Chem.*, 1996, **271**, 3097–3104.
- (a) M. Rosselin, B. Poeggeler and G. Durand, *Curr. Top. Med. Chem.*, 2017, **17**, 2006–2022; (b) C. Oliveira, S. Benfeito, C. Fernandes, F. Cagide, T. Silva and F. Borges, *Med. Res. Rev.*, 2018, **38**, 1159–1187.
- (a) Q. Ke, G. Yan, J. Yu and X. Wu, *Org. Biomol. Chem.*, 2019, **17**, 5863–5881; (b) J. Rostoll-Berenguer, G. Blay, J. R. Pedro and C. Vila, *Eur. J. Org. Chem.*, 2020, DOI: 10.1002/ejoc.202000746; (c) P. Ghosh, B. Ganguly and S. Das, *Org. Biomol. Chem.*, 2020, **18**, 4497–4518.



- 15 Selected examples of the C(sp²)-H functionalization in planar azomethine-based heterocycles: (a) S. Laclef, M. Harari, J. Godeau, I. Schmitz-Afonso, L. Bischoff, C. Hoarau, V. Levacher, C. Fruit and T. Besson, *Org. Lett.*, 2015, **17**, 1700–1703; (b) M. Harari, F. Couly, C. Fruit and T. Besson, *Org. Lett.*, 2016, **18**, 3282–3285; (c) S. C. Ruiz, M. Muselli, S. Frippiat, T. M. Diallo, A. Mohamed-Cherif, V. Levacher, C. Baudequin, L. Bischoff and C. Hoarau, *Synlett*, 2020, **31**, 1185–1190; (d) P. Mampuy, T. D. Moseev, M. V. Varaksin, J. De Houwer, C. M. L. Vande Velde, O. N. Chupakhin, V. N. Charushin and B. U. W. Maes, *Org. Lett.*, 2019, **21**, 2699–2703; (e) W. Zhang, Y.-L. Pan, C. Yang, L. Chen, X. Li and J.-P. Cheng, *J. Org. Chem.*, 2019, **84**, 7786–7795; (f) K. Yin and R. Zhang, *Org. Lett.*, 2017, **19**, 1530–1533; (g) W. Wei, L. Wang, H. Yue, P. Bao, W. Liu, C. Hu, D. Yang and H. Wang, *ACS Sustainable Chem. Eng.*, 2018, **6**, 17252–17257; (h) W. Wei, L. Wang, P. Bao, Y. Shao, H. Yue, D. Yang, X. Yang, X. Zhao and H. Wang, *Org. Lett.*, 2018, **20**, 7125–7130; (i) L.-Y. Xie, J.-L. Hu, Y.-X. Song, G.-K. Jia, Y.-W. Lin, J.-Y. He, Z. Cao and W.-M. He, *ACS Sustainable Chem. Eng.*, 2019, **7**, 19993–19999; (j) L. Wang, M. Sun, L. Zhao, Z. Wang and P. Li, *ChemCatChem*, 2020, DOI: 10.1002/cctc.202000459.
- 16 For the Nobel lectures on the development of transition metal-catalyzed cross-coupling reactions, see: (a) E. Negishi, *Angew. Chem., Int. Ed.*, 2011, **50**, 6738–6764; (b) A. Suzuki, *Angew. Chem., Int. Ed.*, 2011, **50**, 6722–6737.
- 17 Selected early examples on the ‘conventional’ cross-coupling reactions of cyclic azomethines: (a) L. N. Pridgen and L. B. Killmer, *J. Org. Chem.*, 1981, **46**, 5402–5404; (b) A. Dondoni, G. Fantin, M. Fogagnolo, A. Medici and P. Pedrini, *Synthesis*, 1987, 693–696; (c) A. I. Meyers and K. A. Novachek, *Tetrahedron Lett.*, 1996, **37**, 1747–1748; (d) W. D. Schmitz and D. Romo, *Tetrahedron Lett.*, 1996, **37**, 4857–4860; (e) P. A. Jacobi and H. Liu, *J. Am. Chem. Soc.*, 1999, **121**, 1958–1959; (f) P. A. Jacobi and H. Liu, *J. Org. Chem.*, 1999, **64**, 1778–1779; (g) A. Nadin, J. M. Sánchez López, A. P. Owens, D. M. Howells, A. C. Talbot and T. Harrison, *J. Org. Chem.*, 2003, **68**, 2844–2852.
- 18 J. C. Lewis, A. M. Berman, R. J. Bergman and J. A. Ellman, *J. Am. Chem. Soc.*, 2008, **130**, 2493–2500.
- 19 Relevant reviews on cross-dehydrogenative coupling chemistry: (a) C.-J. Li, *Acc. Chem. Res.*, 2008, **42**, 335–344; (b) C. S. Yeung and V. M. Dong, *Chem. Rev.*, 2011, **111**, 1215–1292; (c) S. H. Cho, J. Y. Kim, J. Kwak and S. Chang, *Chem. Soc. Rev.*, 2011, **40**, 5068–5083; (d) Y. Yang, J. Lan and J. You, *Chem. Rev.*, 2017, **117**, 8787–8863; (e) C.-Y. Huang, H. Kang, J. Li and C.-J. Li, *J. Org. Chem.*, 2019, **84**, 12705–12721.
- 20 (a) B. Haag, M. Mosrin, H. Ila, V. Malakhov and P. Knochel, *Angew. Chem., Int. Ed.*, 2011, **50**, 9794–9824; (b) D. Tilly, J. Magolan and J. Mortier, *Chem. – Eur. J.*, 2012, **18**, 3804–3820.
- 21 (a) M. A. Voinov, I. A. Grigor’ev and L. B. Volodarsky, *Heterocycl. Commun.*, 1998, **4**, 261–270; (b) M. A. Voinov, I. A. Grigor’ev and L. B. Volodarsky, *Tetrahedron*, 2000, **56**, 4071–4077; (c) M. A. Voinov and I. A. Grigor’ev, *Tetrahedron Lett.*, 2002, **43**, 2445–2447; (d) M. A. Voinov, T. G. Shevelev, T. V. Rybalova, Y. V. Gatilov, N. V. Pervukhina, A. B. Burdukov and I. A. Grigor’ev, *Organometallics*, 2007, **26**, 1607–1615.
- 22 (a) O. N. Chupakhin and I. Y. Postovskii, *Russ. Chem. Rev.*, 1976, **45**, 454–468; (b) O. N. Chupakhin, V. N. Charushin and H. C. van der Plas, *Tetrahedron*, 1988, **44**, 1–34; (c) O. N. Chupakhin, V. N. Charushin and H. C. van der Plas, *Nucleophilic Aromatic Substitution of Hydrogen*, Academic Press, San Diego, 1994; (d) V. N. Charushin and O. N. Chupakhin, *Pure Appl. Chem.*, 2004, **76**, 1621–1631; (e) V. N. Charushin and O. N. Chupakhin, *Mendeleev Commun.*, 2007, **17**, 249–254; (f) A. M. Prokhorov, M. Małkosza and O. N. Chupakhin, *Tetrahedron Lett.*, 2009, **50**, 1444–1446; (g) O. N. Chupakhin and V. N. Charushin, *Tetrahedron Lett.*, 2016, **57**, 2665–2672; (h) O. N. Chupakhin and V. N. Charushin, *Pure Appl. Chem.*, 2017, **89**, 1195–1208; (i) V. N. Charushin and O. N. Chupakhin, *Russ. Chem. Bull.*, 2019, **68**, 453–471.
- 23 (a) D. Döpp, L. Greci and A. M. Nour-el-Din, *Chem. Ber.*, 1983, **116**, 2049–2057; (b) E. Gössinger and B. Witkop, *Monatsh. Chem.*, 1980, **111**, 803–811; (c) E. G. Janzen, Y.-K. Zhang and D. L. Haire, *Magn. Reson. Chem.*, 1994, **32**, 711–720.
- 24 M. V. Varaksin, I. A. Utepova, O. N. Chupakhin and V. N. Charushin, *J. Org. Chem.*, 2012, **77**, 9087–9093.
- 25 (a) S.-I. Murahashi, J. Sun, H. Kurosawa and Y. Imada, *Heterocycles*, 2000, **52**, 557–561; (b) A. Goti, S. Cicchi, V. Mannucci, F. Cardona, F. Guarna, P. Merino and T. Tejero, *Org. Lett.*, 2003, **5**, 4235–4238; (c) M. Marradi, S. Cicchi, I. Delso, L. Rosi, T. Tejero, P. Merino and A. Goti, *Tetrahedron Lett.*, 2005, **46**, 1287–1290.
- 26 (a) H. Sterckx, B. Morel and B. U. W. Maes, *Angew. Chem., Int. Ed.*, 2019, **58**, 7946–7970; (b) S. P. Teong, X. Li and Y. Zhang, *Green Chem.*, 2019, **21**, 5753–5780.
- 27 (a) K. L. Tan, R. G. Bergman and J. A. Ellman, *J. Am. Chem. Soc.*, 2002, **124**, 3202–3203; (b) S. H. Wiedemann, J. C. Lewis, R. G. Bergman and J. A. Ellman, *J. Am. Chem. Soc.*, 2006, **128**, 2452–2462.
- 28 L. Ackermann, S. Barfüsser, C. Kornhaass and A. R. Kapdi, *Org. Lett.*, 2011, **13**, 3082–3085.
- 29 S. Yotphan, R. J. Bergman and J. A. Ellman, *Org. Lett.*, 2009, **11**, 1511–1514.
- 30 (a) H.-Q. Do, R. M. Kashif Khan and O. Daugulis, *J. Am. Chem. Soc.*, 2008, **130**, 15185–15192; (b) H.-Q. Do and O. Daugulis, *J. Am. Chem. Soc.*, 2008, **130**, 1128–1129; (c) H.-Q. Do and O. Daugulis, *J. Am. Chem. Soc.*, 2007, **129**, 12404–12405.
- 31 M. Muselli, C. Baudequin, C. Hoarau and L. Bischoff, *Chem. Commun.*, 2015, **51**, 745–748.
- 32 M. Muselli, C. Baudequin, C. Perrio, C. Hoarau and L. Bischoff, *Chem. – Eur. J.*, 2016, **22**, 5520–5524.
- 33 R. A. Novikov, E. V. Shulishov and Y. V. Tomilov, *Mendeleev Commun.*, 2012, **22**, 87–89.



- 34 Y. V. Tomilov, R. A. Novikov and O. M. Nefedov, *Tetrahedron*, 2010, **66**, 9151–9158.
- 35 (a) S.-I. Murahashi and Y. Imada, *Chem. Rev.*, 2019, **119**, 4684–4716; (b) I. A. Grigor'ev, in *Nitrile Oxides, Nitrones & Nitronates in Organic Synthesis: Novel Strategies in Synthesis*, ed. H. Feuer, John Wiley & Sons, Inc., Hoboken, New Jersey, 2nd edn, 2008, pp. 129–435.
- 36 X. Dai, M. W. Miller and A. W. Stamford, *Org. Lett.*, 2010, **12**, 2718–2721.
- 37 H. Zhao, R. Wang, P. Chen, B. T. Gregg, M. M. Hsia and W. Zhang, *Org. Lett.*, 2012, **14**, 1872–1875.
- 38 L.-C. Campeau, D. R. Stuart, J.-P. Leclerc, M. Bertrand-Laperle, E. Villemure, H.-Y. Sun, S. Lasserre, N. Guimond, M. Lecavallier and K. Fagnou, *J. Am. Chem. Soc.*, 2009, **131**, 3291–3306.
- 39 (a) J. P. Graham, N. Langlade, J. M. Northall, A. J. Roberts and A. J. Whitehead, *Org. Process Res. Dev.*, 2011, **15**, 44–48; (b) S. Ueda, M. Su and S. L. Buchwald, *J. Am. Chem. Soc.*, 2012, **134**, 700–706.
- 40 M. Koszytkowska-Stawińska, E. Mironiuk-Puchalska and W. Sas, *Tetrahedron Lett.*, 2011, **52**, 1866–1870.
- 41 M. Charette and M. W. Gray, *IUBMB Life*, 2000, **49**, 341–351.
- 42 (a) T. Yoshikawa, S. Kimura, T. Hatano, K. Okamoto, H. Hayatsu and S. Arimoto-Kobayashi, *Food Chem. Toxicol.*, 2002, **40**, 1165–1170; (b) M. Monobe, S. Arimoto-Kobayashi and K. Ando, *Mutat. Res.*, 2003, **538**, 93–99.
- 43 *Electron Paramagnetic Resonance*, ed. B. C. Gilbert, M. J. Davies and D. M. Murphy, Royal Society of Chemistry, Cambridge, 2002, vol. 18.
- 44 E. Demory, D. Farran, B. Baptiste, P. Y. Chavant and V. Blandin, *J. Org. Chem.*, 2012, **77**, 7901–7912.
- 45 (a) M. Thiverny, E. Demory, B. Baptiste, C. Philouze, P. Y. Chavant and V. Blandin, *Tetrahedron: Asymmetry*, 2011, **22**, 1266–1273; (b) M. Thiverny, D. Farran, C. Philouze, V. Blandin and P. Y. Chavant, *Tetrahedron: Asymmetry*, 2011, **22**, 1274–1281; (c) M. Thiverny, C. Philouze, P. Y. Chavant and V. Blandin, *Org. Biomol. Chem.*, 2010, **8**, 864–872.
- 46 (a) H.-Y. Sun, S. I. Gorelsky, D. R. Stuart, L.-C. Campeau and K. Fagnou, *J. Org. Chem.*, 2010, **75**, 8180–8189; (b) L. Ackermann, *Chem. Rev.*, 2011, **111**, 1315–1345; (c) D. Garcia-Cuadrado, P. de Mendoza, A. A. C. Braga, F. Maseras and A. M. Echavarren, *J. Am. Chem. Soc.*, 2007, **129**, 6880–6886; (d) D. Garcia-Cuadrado, A. A. C. Braga, F. Maseras and A. M. Echavarren, *J. Am. Chem. Soc.*, 2006, **128**, 1066–1067; (e) S. I. Gorelsky, D. Lapointe and K. Fagnou, *J. Org. Chem.*, 2012, **77**, 658–668.
- 47 M. Carbó-López, G. Royal, C. Philouze, P. Y. Chavant and V. Blandin, *Eur. J. Org. Chem.*, 2014, 4884–4896.
- 48 M. V. Varaksin, I. A. Utepova and O. N. Chupakhin, *Chem. Heterocycl. Compd.*, 2012, **48**, 1213–1219.
- 49 G. A. El-Hiti, K. Smith, A. S. Hegazy, M. B. Alshammari and A. M. Masmali, *ARKIVOC*, 2015, **2015**, 19–47.
- 50 M. V. Varaksin, I. A. Utepova, O. N. Chupakhin and V. N. Charushin, *Tetrahedron*, 2015, **71**, 7077–7082.
- 51 M. Varaksin, T. Moseev, O. Chupakhin, V. Charushin and B. Trofimov, *Org. Biomol. Chem.*, 2017, **15**, 8280–8284.
- 52 A. A. Akulov, M. V. Varaksin, V. N. Charushin and O. N. Chupakhin, *ACS Omega*, 2019, **4**, 825–834.
- 53 L. A. Smyshliaeva, M. V. Varaksin, P. A. Slepukhin, O. N. Chupakhin and V. N. Charushin, *Beilstein J. Org. Chem.*, 2018, **14**, 2618–2626.
- 54 T. D. Moseev, M. V. Varaksin, D. A. Gorlov, V. N. Charushin and O. N. Chupakhin, *J. Org. Chem.*, 2020, **85**, 11124–11133.
- 55 For a recent example of cyclic azomethine C(sp²)-H bond activation towards the novel C–N bond formation, see: A. A. Akulov, M. V. Varaksin, V. N. Charushin and O. N. Chupakhin, *Chem. Heterocycl. Compd.*, 2019, **55**, 783–787.
- 56 (a) M. Yan, Y. Kawamata and P. S. Baran, *Chem. Rev.*, 2017, **117**, 13230–13319; (b) Y. Yuan and A. Lei, *Acc. Chem. Res.*, 2019, **52**, 3309–3324.
- 57 (a) G. E. M. Crisenza and P. Melchiorre, *Nat. Commun.*, 2020, **11**, 803; (b) N. A. Romero and D. A. Nicewicz, *Chem. Rev.*, 2016, **116**, 10075–10166; (c) M. H. Shaw, J. Twilton and D. W. C. MacMillan, *J. Org. Chem.*, 2016, **81**, 6898–6926; (d) D. M. Schultz and T. P. Yoon, *Science*, 2014, **343**, 1239176.
- 58 Y. Qin, L. Zhu and S. Luo, *Chem. Rev.*, 2017, **117**, 9433–9520.

