## Organic & Biomolecular Chemistry



View Article Online

## COMMUNICATION

Check for updates

**Cite this:** *Org. Biomol. Chem.*, 2021, **19**, 10565

Received 2nd November 2021, Accepted 19th November 2021 DOI: 10.1039/d1ob02153b

rsc.li/obc

# Glutathione peroxidase mimics based on conformationally-restricted, *peri*-like, 4,5-disubstituted fluorene dichalcogenides†

Kesar Jagdev, 跑 a Damiano Tanini, 跑 b Jack W. Lownes, ២ a Carlotta Figliola, ២ a Louise Male, ២ a Antonella Capperucci ២ b and Richard S. Grainger ២ \*a

Glutathione peroxidase (GPx) regulates cellular peroxide levels through glutathione oxidation. GPx-mimics based on 4,5-disubstituted fluorene diselenides, their oxides, and ditellurides show catalytic activities consistent with conformational restriction about the dichalcogen bond.

Organoselenium compounds play a central role in biological systems and medicinal chemistry.<sup>1</sup> The selenocysteine-containing enzyme glutathione peroxidase (GPx) catalyses the reduction of peroxides through oxidation of the endogenous thiol glutathione to glutathione disulfide.<sup>2</sup> The build-up of reactive oxygen species such as peroxides is associated with certain disease states, and hence small selenium-containing molecules which can mimic the function of GPx have potential in drug development.<sup>1,2</sup> A wide range of GPx mimics containing diverse selenium functionality has been investigated, with the aminoselenide Ebselen **1** reaching phase 3 clinical trials for a variety of diseases associated with oxidative stress (Fig. 1).<sup>2,3</sup>

Diselenides are promising GPx mimics,<sup>4–7</sup> with even the simple diphenyl diselenide showing two times greater activity than Ebselen.<sup>6</sup> In 2011, Back reported that 1,8-, *peri*-substituted, naphthalene diselenides 2 show an order of magnitude greater GPx-like activity compared with diphenyl diselenide (Fig. 1).<sup>7</sup> Restricting the conformation around the diselenide bond to almost planar, as found in 2, reduces the HOMO–LUMO energy gap and raises the energy of the HOMO compared with conformationally-unrestricted diphenyl diselenide, thereby increases the rate of oxidation of 2 by peroxide in the rate-determining step.



**Fig. 1** Selenium-containing GPx mimics: Ebselen **1**, Back's conformationally-restricted **1**,8-*peri*-substituted naphthalene diselenides **2** and proposed conformationally-restricted **4**,5-disubstituted fluorene diselenides **3** in this study.

In a search for alternative conformationally-restricted aryl diselenides<sup>‡</sup> that show enhanced GPx-like activity and which are amenable to structural variation towards medicinal chemistry applications, we considered the previously unreported 4,5-fluorene diselenides 3 (Fig. 1). As with *peri*-substituted 1,8-naphthalenes 2, the near planarity of fluorene<sup>8</sup> should constrain the geometry of the diselenide bond, and the close proximity of groups in the 4,5-(bay) region should favour dichalcogen bond formation.<sup>9</sup> In this paper we report the first investigation into the synthesis and properties of 3, its mono- and trioxides, the corresponding ditelluride and their GPx-like activity.

Fluorene diselenides 3a (R = Me) and 3b (R = Bu) were synthesized from fluorenes 4a and 4b, through quenching the 4,5dilithiofluorene species, generated using BuLi-TMEDA,9 with elemental selenium (Scheme 1). Diselenide oxidation was investigated, in view of selenium oxides showing potential GPx-like activity, and to compare their behaviour with the analogous 5-membered ring naphthalene bis-selenium species reported by Kice  $(2a)^{10}$  and Back (2b).<sup>7</sup> Oxidation with 1.2 equivalents of mCPBA in Et<sub>2</sub>O gave selenolseleninates 5a and 5b along with recovered starting material. We did not see any evidence of formation of the symmetrical selenenic anhydride in these mono-oxidations, in contrast to the oxidation of 2a, where a mixture of isomeric monoxides is observed.<sup>10</sup> Use of a larger excess (3.5 equivalents) of mCPBA resulted in the precipitation of seleninic anhydrides 6 as single stereoisomers in excellent 85-95% yields. These were assigned as the trans, C2-symmetric stereoisomers, rather than the alternative cis, meso struc-

<sup>&</sup>lt;sup>a</sup>School of Chemistry, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK. E-mail: r.s.grainger@bham.ac.uk

<sup>&</sup>lt;sup>b</sup>University of Florence, Department of Chemistry "Ugo Shiff", Via della Lastruccia 13, 1-50019 Sesto Fiorentino, Italy

<sup>†</sup> Electronic supplementary information (ESI) available: Experimental procedures and analytical data, copies of NMR spectra and X-ray crystallography for **3a** (CCDC 2099301) and **3b** (CCDC 1470801). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1ob02153b

#### **Organic & Biomolecular Chemistry**

tures, on the basis of the equivalent Me groups at C-9 in the <sup>1</sup>H and <sup>13</sup>C NMR of **6a**. Treatment of **6a** with KOH in CD<sub>3</sub>OD formed the dipotassium salt of the bis-seleninic acid **7a**, evidenced by <sup>77</sup>Se NMR, which upon acidification returned the same stereoisomer **6a** in 86% yield, suggesting the *trans*-isomer is thermodynamic preferred. In contrast, naphthalenes **2** give mixtures of diastereomeric seleninic anhydrides in both selenium oxidation and in base-mediated ring-opening – acidification.<sup>7,10</sup> The ditellurides **8a** and **8b** were also prepared from fluorenes **4a** and **4b** using tellurium as the quench for the dilithio species.

The X-ray crystal structure of **3a** is shown in Fig. 2 and that of **3b** (four independent molecules in the unit cell) in the ESI (Fig S1–S5†).<sup>11</sup> In contrast to the essentially planar naphthalene diselenides **2a** and **2b** (C–Se–Se–C dihedral angle **2a**: -2.28(13),<sup>12</sup> **2b**:  $-1.50(7)^{13}$ ), diselenides **3** are non-planar (C4–Se1–Se2–C5 dihedral angle **3a** -41.35(12); **3b** average 42.3

Se–C dihedral angle is still much smaller than in the conformationally unconstrained diphenyl diselenide (85.4(2), -85.5 (3))<sup>15</sup> and in the less constrained biaryl diselenide, dibenzo[c, e][1,2]diselenine (9, Scheme 1) (59.0(3), -59.0(4), -57.0(4)).<sup>16</sup> The Se–Se bond length in **3b** is 2.34416(4) Å, shorter than naphthalene diselenides **2a** (2.3639(5) Å) and **2b** (2.3552(3) Å), but longer than in diphenyl diselenide (2.3066(7) Å; 2.3073(10) Å) and **9** (mean length 2.323(2) Å). The GPx-like catalytic activities of diselenides **3**, selenolseleninates **5**, seleninic anhydrides **6** and ditellurides **8** were determined using Iwoka's NMR assay,<sup>17</sup> which monitors the drop in

(20)§) and cause a twist in the fluorene plane (C4-C11-C12-C5

dihedral angle 3a -10.9(4); 3b average 11.3(15)§).<sup>14</sup> This C-Se-

ninates 5, seleninic anhydrides 6 and ditellurides 8 were determined using Iwoka's NMR assay,<sup>17</sup> which monitors the drop in concentration of dithiotheritol ( $DTT^{red}$ ) as it is oxidized to the disulfide  $DTT^{ox}$  over time (Fig. 3). A solvent system of 2:1 CD<sub>3</sub>OD:CDCl<sub>3</sub> was used to maintain solubility of all com-



Scheme 1 Synthesis of 4,5-disubstituted fluorene diselenides 3, selenolseleninates 5, seleninic anhydrides 6 and ditellurides 8, and structures of dipotassium salt of bis-seleninic acid 7a and related biaryl diselenide 9. Reagents and conditions: (i) *n*-BuLi (4 equiv.), TMEDA (4 equiv.), 60 °C, 4 h, then Se or Te (8 equiv.), THF, -78 °C-rt. (ii) *m*CPBA (1.2 equiv.), Et<sub>2</sub>O, 15 min.



**Fig. 3** Oxidation of **DTT**<sup>red</sup> with  $H_2O_2$  in the presence of selenium- or tellurium-containing catalysts (10 mol%). Reaction conditions:  $[DTT^{red}]^0 = 0.14 \text{ M}$ ,  $[H_2O_2]^0 = 0.14 \text{ M}$ , [catalyst] = 0.014 M, 2:1 CD<sub>3</sub>OD/CDCl<sub>3</sub> solution (0.6 mL). Reaction progress monitored by <sup>1</sup>H NMR. The mean (<u>+</u>) SD values of three separate experiments are reported.



Fig. 2 Two views of the crystal structure of diselenide **3a** with ellipsoids drawn at the 50% probability level.

Table 1 GPx-like activity of chalcogen-containing catalysts

Entry	Catalyst	Initial <b>DDT</b> <sup>red <math>a</math></sup> (%)	$T_{50}^{b}$ (min)
1	2b	100	>300
2	3a	100	>300
3	3b	100	>300
4	5a	88	$141 (\pm 9)^c$
5	5b	92	253 (±17)
6	6a	75	52 (±8)
7	6b	75	$105(\pm 11)$
8	8a	100	<3
9	8b	100	<3
8 9	8a 8b	100 100	<3 <3

<sup>*a*</sup> After addition of 10 mol% catalyst before addition of  $H_2O_2$ . <sup>*b*</sup>  $T_{50}$  is the time required to halve the initial thiol concentration after the addition of  $H_2O_2$ . <sup>*c*</sup> Data in parenthesis are the experimental error.

#### **Organic & Biomolecular Chemistry**

ponents and hence compare catalytic activity under homogenous conditions, although rates in this solvent system are much slower than in the original report of  $D_2O$ .<sup>18</sup> The times taken for the initial concentration of **DTT<sup>red</sup>** to halve ( $T_{50}$ ), after addition of  $H_2O_2$  are shown in Table 1.  $T_{50}$  allows catalysts to be compared where there is a rapid initial reaction, as is the case herein for selenolseleninates and seleninic anhydrides, prior to addition of  $H_2O_2$ . Back's naphthalene diselenide 2**b**, wherein the electron-donating *ortho*-OMe groups were shown to increase catalytic activity over the non-substituted 2**a**, was also included, along with a background reaction (no catalyst).

All of the selenium- and tellurium-containing compounds 3, 5, 6 and 8 catalyse the oxidation of DTT<sup>red</sup> to DTT<sup>ox</sup>. Diselenides 3 have comparable activities to the naphthalene diselenide 2b in this assay, despite lacking activating ortho-OMe substituents (Fig. 1 and Table 1, entries 1-3). The selenolseleninates 5 have shorter  $T_{50}$  than the corresponding diselenides 3 (Table 1, entries 4 and 5). Before adding H<sub>2</sub>O<sub>2</sub>, approximately 10% of DTT<sup>ox</sup> was detected, pointing to an initial fast reaction that occurs prior to the first NMR reading under these homogenous conditions. A more extensive initial reaction occurs with trioxides 6, with approx. 25% DTT<sup>ox</sup> detected, contributing to the overall shorter  $T_{50}$  (entries 6 and 7). In general, 9,9-dimethyl-substituted fluorenes catalyse the oxidation of DTT<sup>red</sup> faster than the butyl-substituted systems (compare entries 4 vs. 5, and entries 6 vs. 7). Oxidation using ditellurides 8a and 8b is two orders of magnitude faster that the corresponding diselenides 3a and 3b (entries 8 and 9), with reactions complete within minutes of adding H<sub>2</sub>O<sub>2</sub>.<sup>19</sup>

In order to gain further mechanistic insight into the catalytic cycle, stoichiometric reactions of selenium-containing catalysts were carried out (Scheme 2). Treatment of diselenide **3a** with a large excess (10 equiv.) of  $H_2O_2$  in 2 : 1 MeOH :  $CH_2Cl_2$  at room temperature gave slow oxidation to monoxide **5a** (Scheme 2, eqn (1)). No higher oxides were detected, and independent treatment of selenolseleninate **5a** or seleninic anhydride **6a** with  $H_2O_2$  under these conditions gave no reaction, suggesting **6a** is not an intermediate in the catalytic cycle.<sup>20</sup>

Diselenide **3a** does not react with (4-(tert-butyl)phenyl) methanethiol  $(10)^{21}$  in CH<sub>2</sub>Cl<sub>2</sub>/MeOH at room temperature (Scheme 2, eqn (2)). However reaction of selenolseleninate **5a** with 10 equivalents of **10** gave an essentially instantaneous and quantitative transformation to diselenide **3a** and disulfide **11** along with 75% recovered thiol **10**, (eqn (3)). Under the same conditions, seleninic anhydride **6a** underwent a similarly rapid and high-yielding transformation to **3a** and **11** (eqn (4)), where the 90% yield of **11** is based on theoretical consumption of 6 molar equivalents of thiol **10** (stoichiometries in the reactions of **5a** and **6a** with thiols are shown in the ESI, Scheme S1<sup>†</sup>).

No intermediate bis-selenium species were observed in the reactions of 5a and 6a with thiol 10. However, reaction of 5a with the bulkier thiol, *t*-BuSH, gave the bis-selenenyl sulfide 12a (Scheme 2, eqn (5)), a potential intermediate in the formation of 3a. Indeed, isolated 12a is slowly transformed over 24 h in



Scheme 2 Mechanistic investigations. % yields of recovered thiol are based on theoretical consumption.

solution to diselenide **3a** and di-*tert*-butyl disulfide (**13**). This rate of this reaction is not changed by addition of 3 equivalents of thiol **10**, and no disulfides derived from **10** were formed, only **13**. The breakdown of bis-selenenyl sulfide **12a** to diselenide **3a** and disulfide **13** is thus presumably intramolecular, but given the steric hindrance provided by the *t*-Bu group, care should be taken in extrapolating these observations to all thiols. Kice reported a similar reaction of *t*-BuSH with the monoxides of naphthalene diselenide **2a** to give isolable **1**,8-bis[(*tert*-butylthio) seleno]naphthalene,<sup>10</sup> which led Back to propose bis-selenenyl sulfides as intermediates in the catalytic cycle of **2b**.<sup>7</sup>

The reaction of seleninic anhydride **6a** with *t*-BuSH also gave bis-selenenyl sulfide **12a** (Scheme 2, eqn (6)), though clearly there are multiple potential intermediates preceding its formation. These intermediates account for the formation of disulfide **13** (98% based on theoretical amount of *t*-BuSH consumed and consistent with recovery of 4 equivalents of thiol, Scheme S1<sup>†</sup>), not observed in the reaction of selenolseleninate **5a** with *t*-BuSH.¶

Based on the above observations, a catalytic cycle directly analogous to that proposed by Back for naphthalene diselenides **2** is suggested<sup>7</sup> (Scheme 3): this cycle is mechanistically distinct from catalysis by other diselenides, which involve initial Se–Se bond cleavage by reaction with thiols.<sup>22</sup> The ratedetermining step is the oxidation of diselenide **3** to selenolseleninate **5**, which in turn rapidly consumes two equivalents of thiol and forms disulfides *via* the intermediates **14** (not observed) and **12** (observed as **12a** for R = Me, R' = *t*-Bu). As noted above, the conversion of **12** to **3** may occur by more than one mechanism and may also be catalysed by thiol: this step is severely slowed in the case of R' = *t*-Bu where nucleophilic attack at sulfur is restricted and where an intramolecular mechanism appears most likely. Oxides **5** and **6** initially circumvent the rate-determining oxidation, resulting in overall



Scheme 3 Proposed catalytic cycle for oxidation of thiols to disulfides.

shorter  $T_{50}$  in the **DDT** NMR assay. The initial rapid reaction of 5 and 6 with **DTT<sup>red</sup>** is evident in Fig. 3. Consumption of 1 equivalent of dithiol **DTT<sup>red</sup>** (2 × SH) with the 10 mol% of catalysts 5a and 5b present at the start of the assay should lead to an immediate 10% reduction in the amount of **DTT<sup>red</sup>**, which is consistent with the approx. 10% observed initial **DTT<sup>red</sup>** (Table 1, entries 4 and 5). Similarly, rapid consumption of 3 equivalents of **DDT<sup>red</sup>** (6 × SH) with the starting 10 mol% of catalysts 5a and 5b should give a theoretical 30% reduction in the amount of **DDT<sup>red</sup>**, with approx. 25% reduction observed in practice (entries 6 and 7).

## Conclusions

In conclusion, readily synthesized, bay-substituted 4,5-fluorene diselenides **3** possess properties analogous to *peri*-substituted 1,8-naphthalenes **2**, including increased GPx-like activity compared with non-conformationally constrained diselenides. Despite a greater twist in the diselenide bond, the catalytic activity of fluorenes **3a** and **3b** is similar to that of naphthalene **2b** in a homogenous **DDT** redox assay, without the need for additional activation by *ortho*-OMe groups on the aromatic rings. Moving forward, the fluorene scaffold is particularly amenable to structural variation through incorporation of different functionality at C-9, for example towards watersoluble GPx mimics,<sup>18,23</sup> and the close proximity of groups in the 4,5-bay positions may be exploited in other applications based on 1,8-*peri*-substituted naphthalenes.<sup>24,25</sup>

## Conflicts of interest

There are no conflicts to declare.

### Acknowledgements

We thank EPSRC (EP/K039245/1) for funding.

## Notes and references

‡ Aryl selenides are less toxic than alkyl selenides. See ref. 3*a*.

 $\$  For 3b the average value calculated from molecules 1–3 for each parameter is given (see ESI†).

 $\P Ph_3CSH$  and 1-adamantylthiol gave the corresponding disulfide and diselenide 3a directly from 6a (see ESI†).

- Reviews: (*a*) C. W. Nogueira, N. V. Barbosa and J. B. T. Rocha, *Arch. Toxicol.*, 2021, **95**, 1179–1226; (*b*) D. Radomska, R. Czarnomysy, D. Radomski and K. Bielawski, *Int. J. Mol. Sci.*, 2021, **22**, 1009; (*c*) Z. Chen, H. Lai, L. Hou and T. Chen, *Chem. Commun.*, 2020, **56**, 179–196.
- 2 (a) N. V. Barbosa, C. W. Nogueira, P. A. Nogara, A. F. de Bem, M. Aschner and J. B. T. Rocha, *Metallomics*, 2017, 9, 1703–1734; (b) L. Orian and S. Toppo, *Free Radicals Biol. Med.*, 2014, 66, 65–74.
- 3 Reviews: (a) K. N. Sands, T. A. Tuck and T. G. Back, *Chem. Eur. J.*, 2018, 24, 9714–9728; (b) G. Mugesh and H. B. Singh, *Chem. Soc. Rev.*, 2000, 29, 347–357.
- 4 Reviews: (a) M. D. Tiezza, G. Ribaudo and L. Orian, *Curr.* Org. Chem., 2019, 23, 1381–1402; (b) M. Álvarez-Pérez, W. Ali, M. A. Marć, J. Handzlik and E. Domínguez-Álvarez, *Molecules*, 2018, 23, 628.
- 5 Recent representative examples: (a) K. Arai, Y. Sato, I. Nakajima, M. Saito, M. Sasaki, A. Kanamori and M. Iwaoka, *Bioorg. Med. Chem.*, 2021, 29, 115866; (b) D. Bhowmick and G. Mugesh, Org. Biomol. Chem., 2015, 13, 9072–9082; (c) V. P. Singh, J.-f. Poon, R. J. Butcher, X. Lu, G. Mestres, M. K. Ott and L. Engman, J. Org. Chem., 2015, 80, 7385–7395; (d) P. Prabhu, B. G. Singh, M. Noguchi, P. P. Phadnis, V. K. Jain, M. Iwaoka and K. I. Priyadarsini, Org. Biomol. Chem., 2014, 12, 2404–2412.
- 6 S. R. Wilson, P. A. Zucker, R.-R. C. Huang and A. Spector, J. Am. Chem. Soc., 1989, 111, 5936–5939.
- 7 D. J. Press and T. G. Back, Org. Lett., 2011, 13, 4104-4107.
- 8 L. A. Chetkina and V. K. Belsky, *Crystallogr. Rep.*, 2013, 58, 26–48.
- 9 V. B. Bonifácio, J. Morgado and U. Scherf, *Synlett*, 2010, 1333–1336.
- 10 J. L. Kice, Y.-H. Kang and M. B. Manek, J. Org. Chem., 1988, 53, 2435–2439.
- 11 CCDC 2099301 (3a) and 1470801 (3b)† contain the supplementary crystallographic data for this paper.
- 12 S. M. Aucott, H. L. Milton, S. D. Robertson, A. M. Z. Slawin and J. D. Woollins, *Heteroat. Chem.*, 2004, **15**, 530–542.
- 13 C. Figliola, L. Male, P. N. Horton, M. B. Pitak, S. J. Coles, S. L. Horswell and R. S. Grainger, *Organometallics*, 2014, 33, 4449–4460.
- A. L. Fuller, L. A. S. Scott-Hayward, Y. Li, M. Bühl,
  A. M. Z. Slawin and J. D. Woollins, *J. Am. Chem. Soc.*, 2010, 132, 5799–5802.

- 15 We have observed a similar twist in a 4,5-carbazole diselenide: I. A. Pocock, A. M. Alotaibi, K. Jagdev, C. Prior, G. R. Burgess, L. Male and R. S. Grainger, *Chem. Commun.*, 2021, 57, 7252–7255.
- 16 M. R. Bryce, A. Chesney, A. K. Lay, A. S. Batsanov and J. A. K. Howard, J. Chem. Soc., Perkin Trans. 1, 1996, 2451–2459.
- 17 K. Kumakura, B. Mishra, K. I. Priyadarsini and M. Iwaoka, *Eur. J. Org. Chem.*, 2010, 440–445.
- 18 The DDT assay has been used in CD<sub>3</sub>OD: K. Arai, A. Tashiro, Y. Osaka and M. Iwaoka, *Molecules*, 2017, 22, 354.
- 19 For higher GPx-like activity of Te derivatives *vs.* Se derivatives, see (*a*) Ref. 7; (*b*) D. Tanini, L. Ricci and A. Capperucci, *Adv. Synth. Catal.*, 2020, **362**, 1323–1332 and references therein.
- 20 For investigation of seleninic anhydrides as GPx mimics see: S.-C. Yu, A. Borchert, H. Kuhn and I. Ivanov, *Chem. – Eur. J.*, 2008, 14, 7066–7071.
- 21 Chosen as a "non-odourous" alkyl thiol mimic of glutathione: M. Node, K. Kumar, K. Nishide, S.-i. Ohsugi and T. Miyamoto, *Tetrahedron Lett.*, 2001, **42**, 9207–9210.
- 22 D. Bhowmick and G. Mugesh, *Org. Biomol. Chem.*, 2015, **13**, 10262–10272.

- 23 N. M. R. McNeil, D. J. Press, D. M. Mayder, P. Garnica, L. M. Doyle and T. G. Back, *J. Org. Chem.*, 2016, **81**, 7884– 7897.
- 24 Reviews of peri-substituted dichalcogenides: (a) P. Kilian, F. R. Knight and J. D. Woollins, Coord. Chem. Rev., 2011, 255, 1387-1413; (b) P. Kilian, F. R. Knight and J. D. Woollins, Chem. - Eur. J., 2011, 17, 2302-2328. For application in other enzyme mimics see: (c) Ref. 13; (d) C. Figliola, L. Male, S. L. Horswell and R. S. Grainger, Eur. J. Inorg. Chem., 2015, 3146-3156; (e) S. Mondal, D. Manna, K. Raja and G. Mugesh, ChemBioChem, 2020, 21, 911-923. For additional applications from our group see: (f) R. S. Grainger, B. Patel and B. M. Kariuki, Angew. Chem., Int. Ed., 2009, 48, 4832-4835; (g) R. S. Grainger, B. Patel, B. M. Kariuki, L. Male and N. Spencer, J. Am. Chem. Soc., 2011, 133, 5843-5852; (h) B. Patel, J. Carlisle, S. E. Bottle, G. R. Hanson, B. M. Kariuki, L. Male, J. C. McMurtrie, N. Spencer and R. S. Grainger, Org. Biomol. Chem., 2011, 9, 2336-2344.
- 25 For increased proton sponge basicity in 4,5-disubstituted fluorene vs. 1,8-naphthalene see: H. A. Staab, T. Saupe and C. Krieger, *Angew. Chem., Int. Ed. Engl.*, 1983, 22, 731–732.