Organic & Biomolecular Chemistry

ROYAL SOCIETY OF CHEMISTRY

CORRECTION

View Article Online
View Journal | View Issue

Cite this: *Org. Biomol. Chem.*, 2021, **19**, 1155

Correction: Rhodium(III)-catalyzed oxidative alkylation of *N*-aryl-7-azaindoles with cyclopropanols

Jidan Liu,*[a] Jinyuan Jiang,[a] Zhenke Yang,[a] Qiaohai Zeng,[a] Jieying Zheng,[a] Siying Zhang,[a] Liyao Zheng,[a] Shang-Shi Zhang[b] and Zhao-Qing Liu*[a]

DOI: 10.1039/d1ob90008k

rsc.li/obc

Correction for 'Rhodium(III)-catalyzed oxidative alkylation of *N*-aryl-7-azaindoles with cyclopropanols' by Jidan Liu *et al.*, *Org. Biomol. Chem.*, 2021, DOI: 10.1039/d0ob02323j.

The authors regret that in Table 1 the additives for entries 7–11 are incorrect. The correct table is shown below.

Table 1 Optimization of reaction conditions[a]



1a **2a** **3a**

Entry	Rh catalyst	Additive	Oxidant	Yield[b] (%)
1	$[Cp^*RhCl_2]_2$	CsOAc	$Cu(OAc)_2$	82
2	$[Cp^*RhCl_2]_2$	CsOAc	$PhI(OAc)_2$	9
3	$[Cp^*RhCl_2]_2$	CsOAc	O_2	Trace
4	$[Cp^*RhCl_2]_2$	CsOAc	Ag_2CO_3	54
5	$[Cp^*RhCl_2]_2$	CsOAc	AgOAc	62
6	$[Cp^*RhCl_2]_2$	CsOAc	Ag_2O	48
7	$[Cp^*RhCl_2]_2$	CsOAc	$Cu(OAc)_2{\cdot}H_2O$	86
8	$[Cp^*RhCl_2]_2$	KOAc	$Cu(OAc)_2{\cdot}H_2O$	83
9	$[Cp^*RhCl_2]_2$	NaOAc	$Cu(OAc)_2{\cdot}H_2O$	75
10	$[Cp^*RhCl_2]_2$	$AgSbF_6$	$Cu(OAc)_2{\cdot}H_2O$	43
11	$[Cp^*RhCl_2]_2$	AgOAc	$Cu(OAc)_2{\cdot}H_2O$	36
12	$Cp^*Rh(OAc)_2$	CsOAc	$Cu(OAc)_2{\cdot}H_2O$	80
13	$Rh(Ph_3P)_3Cl$	CsOAc	$Cu(OAc)_2{\cdot}H_2O$	14
14	$Cp^*Rh(CH_3CN)_3(SbF_6)_2$		$Cu(OAc)_2{\cdot}H_2O$	22
15	$[Cp^*RhCl_2]_2$	CsOAc	$Cu(OAc)_2{\cdot}H_2O$	65[c]
16	$[Cp^*RhCl_2]_2$	CsOAc	$Cu(OAc)_2{\cdot}H_2O$	83[d]
17	$[Cp^*RhCl_2]_2$	CsOAc	$Cu(OAc)_2{\cdot}H_2O$	77[e]
18	$[Cp^*RhCl_2]_2$	CsOAc	$Cu(OAc)_2{\cdot}H_2O$	85[f]
19	$[Cp^*RhCl_2]_2$	CsOAc	$Cu(OAc)_2{\cdot}H_2O/O_2$	18[g]
20		CsOAc	$Cu(OAc)_2{\cdot}H_2O$	nr
21	$[Cp^*RhCl_2]_2$	CsOAc		nr

[a] Reaction conditions: **1a** (0.2 mmol), **2a** (0.3 mmol), Rh catalyst (4 mol%), additive (20 mol%) and oxidant (2 equiv.) in MeOH (1 mL) in a sealed tube at 50 °C under N_2 for 4 h. [b] Isolated yield of **3a** based on **1a**. [c] At 30 °C. [d] At 70 °C. [e] $Cu(OAc)_2{\cdot}H_2O$ (1.8 equiv.) was used. [f] $Cu(OAc)_2{\cdot}H_2O$ (2.1 equiv.) was used. [g] $Cu(OAc)_2{\cdot}H_2O$ (0.2 equiv.) was used.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

[a]*School of Chemistry and Chemical Engineering/Institute of Clean Energy and Materials/Guangzhou Key Laboratory for Clean Energy and Materials, Guangzhou University, Guangzhou, P. R. China. E-mail: jdliu@gzhu.edu.cn, lzqgzu@gzhu.edu.cn*

[b]*Center for Drug Research and Development, Guangdong Pharmaceutical University, Guangzhou, P. R. China*