

CORRECTION

[View Article Online](#)[View Journal](#) | [View Issue](#)Cite this: *Org. Chem. Front.*, 2021, **8**, 5911

Correction: One-pot, three-component regioselective coupling reaction of triphenylamine/carbazole derivatives with [60] fullerene and indoles *via* an “umpolung relay” strategy



Xin-Rui Chen, Qian-Wen Zhang, Ge-Ge Tao, Jun Xuan, Hong-Ping Zhou, Yu-Peng Tian and Fei Li*

DOI: 10.1039/d1qo90077c
rsc.li/frontiers-organicCorrection for ‘One-pot, three-component regioselective coupling reaction of triphenylamine/carbazole derivatives with [60]fullerene and indoles *via* an “umpolung relay” strategy’ by Xin-Rui Chen *et al.*, *Org. Chem. Front.*, 2021, DOI: 10.1039/D1QO01058A.

The authors regret that in Table 2, the chemical structure of **3ab-fi** were incorrectly presented as a 1,2-[60]fullerene adduct rather than a 1,4-[60]fullerene adduct. The corrected table is given below.

Department of Chemistry, Anhui University; Key Laboratory of Structure and Functional Regulation of Hybrid Materials (Anhui University), Ministry of Education; Anhui Province Key Laboratory of Chemistry for Inorganic/Organic Hybrid Functionalized Materials and Key Laboratory of Functional Inorganic Materials of Anhui Province, Hefei, Anhui 230601, China. E-mail: lfe@ahu.edu.cn



Table 2 Results of different triphenylamine derivatives as substrates^{a,b}

^a Unless otherwise specified, all reactions were performed with C₆₀ (0.05 mmol), **1** (0.06 mmol, 1.2 equiv.), KO^tBu (0.10 mmol, 2.0 equiv.), CF₃SO₃H (1 mmol, 20 equiv.), triphenylamine derivatives **2** (0.25 mmol, 5 equiv.) and DDQ (0.25 mmol, 5.0 equiv.) in Schlenk flasks under Ar atmosphere for indicated time. ^b Isolated yield; the values in parentheses are based on consumed C₆₀. ^c The third step of the reaction was conducted at 100 °C for 120 min. ^d The reaction was performed in 1,2-dichlorobenzene/DMSO (4 : 1) and the third step of the reaction was conducted at 150 °C for 60 min. ^e The third step of the reaction was conducted at 110 °C for 60 min. ^f The third step of the reaction was conducted at 130 °C for 120 min.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

