RSC Advances



PAPER

View Article Online
View Journal | View Issue



Cite this: RSC Adv., 2021, 11, 4235

Received 17th November 2020 Accepted 9th January 2021

DOI: 10.1039/d0ra09758f

rsc.li/rsc-advances

Catalyzed ring transformation of cyclic N-arylazadiperoxides with participation of α, ω -dithiols†

Nataliya N. Makhmudiyarova, **D** Kamil R. Shangaraev, Irina R. Ishmukhametova, Askhat G. Ibragimov and Usein M. Dzhemilev

Co(OAc)₂-catalyzed ring transformation reaction of 10-aryl-7,8,12,13-tetraoxa-10-azaspiro[5.7]tridecanes with α , ω -dithiols (ethane-1,2-, propane-1,3-, butane-1,4-, pentane-1,5-, and hexane-1,6-dithiols, 3,6-dioxaoctane-1,8-dithiol) giving 3-aryl-1,5,3-dithiazacyclanes was studied.

Cyclic peroxides attract attention for their antimalarial,1 antibacterial,2 and antitumor3 activities. Among numerous cyclic peroxides, heteroatomic cyclic peroxides occupy a special place owing to their high biological activities.4 The methods of synthesis of heteroatom-containing cyclic peroxides are limited. Recently,5-10 nitrogen- and sulfur-containing cyclic di- and triperoxides with antitumor activity have been synthesized.⁵⁻⁹ The development of efficient methods for the preparation of new cyclic hetero-di(tri)peroxides5-10 promotes active investigation of their transformations. It was shown that the reduction of silatriperoxycycloalkanes with PPh_3 affords siladiperoxycycloalkanes;11 the reaction of spiro{adamantane-[2,3']-(pentaoxacane)} with o-phenylenediamine results in the synthesis of benzodioxazocine.5 The implemented conversion of pentaoxacane with o-phenylenediamine to benzodioxazocine⁵ suggests that cyclic N-containing peroxides can be involved in reactions with binucleophilic reagents, in particular α,ω -dithiols, to give new heterocycles. In contrast to the previously described methods of synthesis5-10 and transformation of the peroxide ring, 5,11 this work for the first time discusses the method of catalytic conversion of tetraoxazaspirotridecane to dithiazacycloalkanes.

It was shown by preliminary experiments that the reaction of 10-phenyl-7,8,12,13-tetraoxa-10-azaspiro[5.7]tridecane 1 with ethane-1,2-dithiol 2 does not proceed without a catalyst. The reaction of azadiperoxide 1 with ethane-1,2-dithiol 2 catalyzed by $Sm(NO_3)_3\cdot 6H_2O,\ H_2SO_4$ or $BF_3\cdot Et_2O$ in THF as a solvent affords 3-phenyl-1,5,3-dithiazepane 8 in 10–15% yield (Scheme 1, Table 1). It was found that the yield of 3-phenyl-1,5,3-dithiazepane¹² is affected by the nature of the catalyst. When the reaction is carried out in a polar solvent (MeOH) in the presence of catalytic amounts of $Sm(NO_3)_3\cdot 6H_2O,\ H_2SO_4$ or $BF_3\cdot Et_2O$, the yield of the target product 8 increases to 30%. In the presence of the $Co(OAc)_2$ catalyst, the

Institute of Petrochemistry and Catalysis, Russian Academy of Sciences, 141 Prospekt Oktyabrya, 450075 Ufa, Russian Federation. E-mail: natali-mnn@mail.ru

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0ra09758f

yield of heterocycle **8** is 85%. When $AlCl_3$ or CuCl catalysts are used, the yields of heterocycle **8** are 55% and 75%, respectively (Table 1). Under these conditions, cyclohexanone is formed and O_2 is released (Scheme 1). All reactions were carried out at room temperature for 20 h.

A probable pathway to the synthesis of 3-phenyl-1,5,3-dithiazepane **8** from 10-phenyl-7,8,12,13-tetraoxa-10-azaspiro [5.7]tridecane **1** includes¹³ coordination of the peroxide oxygen atom to the central atom of the catalyst, nucleophilic addition of ethane-1,2-dithiol to the resulting carbocation, ^{14,15} and the subsequent ring closure giving heterocycle **8** (Scheme 2).


Under conditions including 5 mol% of Co(OAc)₂, 20 °C, MeOH, and 20 h, 10-phenyl-7,8,12,13-tetraoxa-10-azaspiro[5.7] tridecane **1** was allowed to react with propane-1,3- **3**, butane-1,4- **4**, pentane-1,5- **5**, and hexane-1,6-dithiols **6**, which furnished the corresponding 3-phenyl-1,5,3-dithiaazacy-cloalkanes¹⁶ **9–12** in 83–89% yields (Table 1). The ring transformation reaction of azadiperoxide **1** with 3,6-dioxa-1,8-octanedithiol 7 (monooxa derivative is shown in the scheme) under the conditions described above resulted in the synthesis of 6-phenyl-1,11-dioxa-4,8-dithia-6-azacyclotridecane¹⁶ **12** in 91% yield (Scheme 1).

The discovered ring transformation reaction of azadiper-oxide **1** with ethane-1,2-dithiol **2** was also carried out for 10-aryl-7,8,12,13-tetraoxa-10-azaspiro[5.7]tridecanes **14–24**, which produced 3-aryl-1,5,3-dithiazepanes¹² **25–35** in 76–90% yields (Scheme 3).

Scheme 1 Ring transformation reaction of 10-phenyl-7,8,12,13-tet-raoxa-10-azaspiro[5.7]tridecane with α,ω -dithiols.

Table 1 Effect of the catalyst and solvent nature on the yield of 3-phenyl-1,5,3-dithiazacyclanes (\sim 20 °C, 20 h)

No.	Compound	[Cat]	Solvent	Yield, %
1	8	AlCl ₃	THF	45
2	8	AlCl ₃	MeOH	55
3	8	$Co(OAc)_2$	THF	79
4	8	$Co(OAc)_2$	MeOH	85
5	8	$BF_3 \cdot OEt_2$	THF	15
6	8	$BF_3 \cdot OEt_2$	MeOH	30
7	8	CuCl	THF	68
8	8	CuCl	MeOH	75
9	8	H_2SO_4	THF	13
10	8	H_2SO_4	MeOH	25
11	8	$Sm(NO_3)_3 \cdot 6H_2O$	THF	10
12	8	$Sm(NO_3)_3 \cdot 6H_2O$	MeOH	20
13	8	_	THF	_
14	8	_	MeOH	_
15	9	$Co(OAc)_2$	MeOH	87
16	10	$Co(OAc)_2$	MeOH	79
17	11	$Co(OAc)_2$	MeOH	83
18	12	$Co(OAc)_2$	MeOH	89
19	13	Co(OAc) ₂	MeOH	91



Scheme 2 Probable synthesis mechanism for 3-phenyl-1,5,3-dithiazepane 8.

Scheme 3 Ring transformation reaction of 10-aryl-7,8,12,13-tetraoxa-10-azaspiro[5.7]tridecanes with ethane-1,2-dithiol.

m-ClC₆H₄ (19, 30), o-BrC₆H₄ (20, 31), m-BrC₆H₄ (21, 32), p-BrC₆H₄ (22, 33),

In conclusion, we demonstrated that on treatment with α,ω -alkanedithiols and the $Co(OAc)_2$ catalyst, azadiperoxides are converted to *N*-aryl-substituted 1,5,3-dithiazamacroheterocycles in high yields.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgements

The reported study was funded by RFBR according to the research project No. 20-33-90002/20.

Notes and references

- 1 R. Slack, A. Jacobine and G. Posner, *Med. Chem. Commun.*, 2012, 3, 281.
- 2 V. Vil', I. Yaremenko, A. Ilovaisky and A. Terent'ev, *Molecules*, 2017, 22, 1881.
- 3 D. Liu and J. Liu, *Nat. Prod. Bioprospect.*, 2013, 3, 161; M. P. Crespo-Ortiz and M. Q. Wei, Antitumor Activity of Artemisinin and Its Derivatives: From a Well-Known Antimalarial Agent to a Potential Anticancer Drug, *J. Biomed. Biotechnol*, 2012, 257597.
- 4 Y. Tu, Nat. Med., 2011, 17, 1217.
- 5 T. V. Tyumkina, N. N. Makhmudiyarova, G. M. Kiyamutdinova, E. S. Meshcheryakova, K. Sh. Bikmukhametov, M. F. Abdullin, L. M. Khalilov, A. G. Ibragimov and U. M. Dzhemilev, *Tetrahedron*, 2018, 74, 1749.
- 6 N. N. Makhmudiyarova, I. R. Ishmukhametova, L. U. Dzhemileva, T. V. Tyumkina, V. A. D'yakonov, A. G. Ibragimov and U. M. Dzhemilev, *RSC Adv.*, 2019, **9**, 18923.
- N. N. Makhmudiyarova, R. Sh. Rakhimov, T. V. Tyumkina,
 E. S. Meshcheryakova, A. G. Ibragimov and
 U. M. Dzhemilev, Russ. J. Org. Chem., 2019, 5, 620.
- N. N. Makhmudiyarova, K. R. Shangaraev, L. U. Dzhemileva,
 T. V. Tuymkina, E. S. Mescheryakova, V. A. D'yakonov,
 A. G. Ibragimov and U. M. Dzhemilev, RSC Adv., 2019, 9, 29949.
- N. N. Makhmudiyarova, I. R. Ishmukhametova,
 L. U. Dzhemileva, V. A. D'yakonov, A. G. Ibragimov and
 U. M. Dzhemilev, *Molecules*, 2020, 25, 1874.
- N. N. Makhmudiyarova, I. R. Ishmukhametova,
 A. G. Ibragimov and U. M. Dhzemilev, *Dokl. Chem.*, 2020,
 492, 93.
- 11 N. N. Makhmudiyarova, I. R. Ishmukhametova and A. G. Ibragimov, *Russ. J. Org. Chem.*, 2020, **10**, 1495.
- 12 N. N. Murzakova, K. I. Prokof'ev, T. V. Tyumkina and A. G. Ibragimov, Russ. J. Org. Chem., 2012, 48, 588.
- (a) S. Oda, J. Franke and M. Krishce, *J. Chem. Sci.*, 2016, 7, 136; (b) S. Vojacek, K. Beese, Z. Alhalabi, S. Swyter, A. Bodtke, C. C. Schulzke, M. Jung, W. Sippl and A. Link, *Arch. Pharm.*, 2017, 350, e1700097.
- 14 U. J. Wellmar, Heterocyclic Chem, 1998, 35, 1531.
- 15 K. Krohn and S. Cludius-Brandt, Synthesis, 2010, 8, 1344.
- 16 N. N. Makhmudiyarova, L. V. Mudarisova,
 E. S. Meshcheryakova, A. G. Ibragimov and
 U. M. Dzhemilev, *Tetrahedron*, 2015, 71, 259.
- 17 N. N. Makhmudiyarova, G. M. Khatmullina, R. Sh. Rakhimov, E. S. Meshcheryakova, A. G. Ibragimov and U. M. Dzhemilev, *Tetrahedron*, 2016, 72, 3277.

o-CH₃C₆H₄ (23, 34), m-CH₃C₆H₄ (24, 35)