RSC Advances



REVIEW

View Article Online
View Journal | View Issue



Cite this: RSC Adv., 2021, 11, 15060

Natural products in *Cyperus rotundus* L. (Cyperaceae): an update of the chemistry and pharmacological activities

Cyperus rotundus L. (Nutgrass, family Cyperaceae) is a notorious weed which is widespread in temperate tropical and subtropical regions of the world. Owing to its richness and potent pharmacological activities, efforts have been devoted to identify its bioactive constituents. Since 1965, a total of about 192 compounds including terpenoids, flavonoids, stilbenes, aromatics and aliphatic fatty acids have been characterized. This review summarizes the bioactivities and mechanism of action of some of the compounds from *C. rotundus* L.

Received 19th January 2021 Accepted 24th March 2021

DOI: 10.1039/d1ra00478f

rsc.li/rsc-advances

1 Introduction

Cyperus rotundus (CR) L. (Nutgrass, family: Cyperaceae) popularly called "the world's worst weed" is widely distributed in subtropical and tropical regions of the world.¹⁻³ It is a notorious weed and has a destructive effect on agricultural yields after it invades the crop fields.^{4,5} It is a smooth, erect, glabrous, grasslike, fibrous rooted, perennial herb that grows up to 15–60 cm height (Fig. 1) and reproduces widely through rhizomes and tubers.⁶ In Chinese traditional medicine, the rhizomes are used for the treatment of liver diseases, stomach ache, inflammatory

diseases, bowel and menstrual disorders.⁷⁻¹³ They are also recommended in India for the treatment of diabetes, arthritis, diarrhoea, dysentery, leprosy, bronchitis, amenorrhea, dysmenorrhea, fever, arthritis and blood disorders.¹⁴⁻¹⁶ In West Asia, the roots are applied in traditional medicine for the treatment of leprosy, thirst, fever, and blood diseases.^{17,18} In Egyptian folk medicine, the tubers are used as an antihelmintic, aphrodisiac, diuretic, sedative, carminative, stimulant and tonic, and for treating renal colic and stomach ache.¹⁹ This perennial herb has recently received much attention due to its broad range of pharmacological and biological activities.^{5,20-75}



(IITA), Cotonou, Benin.

Smith B. Babiaka is an Assistant Lecturer at the Department of Chemistry, University of Buea (Cameroon), where he received his PhD in Chemistry in 2019. His research interest has been focused on natural products drug discovery and molecular molecular modelling of potentials hits. He has served as a consultant at the Agro-Ecohealth Unit, International Institute of Tropical Agriculture



Aurélien F. A. Moumbock received both BSc (2013) and MSc (2017) degrees in Chemistry at the University of Buea (Cameroon). Since 2018, he is carrying out doctoral studies in pharmaceutical sciences at the University of Freiburg (Germany) under the guidance of Profs Stefan Günther and Henning J. Jessen, with a fellowship from the German Academic Exchange Service (DAAD). His

research focuses on the development and application of chemoinformatics methods and tools to accelerate (nature-inspired) small-molecule drug discovery.

[&]quot;Department of Chemistry, University of Buea, P. O. Box 63, Buea, Cameroon. E-mail: babiaka.smith@ubuea.cm; fidele.ntie-kang@ubuea.cm

^bInstitute of Pharmaceutical Sciences, Albert-Ludwigs-Universität Freiburg, Hermann-Herder-Straße 9, D-79104 Freiburg, Germany

Institute of Pharmacy, Martin-Luther University Halle-Wittenberg, Halle (Saale), Germany

^dInstitute of Botany, Technical University of Dresden, Dresden, Germany



Fig. 1 Photograph of the plant Cyperus rotundus L. (Cyperaceae).

Several reports have stated the presence of terpenoids, flavonoids, stilbene derivatives and other classes compounds. 8,11,17,41,76-78 To the best of our knowledge, a total of about 192 NPs with structural diversity have been isolated from this medicinal weed. 8,12,13,17,41,79-94 Previous reviews have focused on ethnobotanical uses, and pharmacological activities. 6,71,95,96 Also, a majority of researchers have reported in vitro bioactivities and GC-MS analysis of crude extracts of this weed. 2,14,41,97-111 The objective of the present review was to provide an update of NPs derived from this plant species, their bioactivity and the




Fig. 2 Pie chart showing the distribution by compound class.

mechanism of actions of some of the compounds from published data in literature.

Data source and preparation

In this review article, a comprehensive search was performed in the following databases: PubMed, SciFinder, Science Direct, Web of Science, Wiley Online, ResearchGate, Google scholar and other search engines were explored for studies published from 1965-2020. Keywords such as: "C. rotundus", 'bioactive compounds', and "pharmacological activities" were used. We removed duplicated papers, then screened the data, ruled out irrelevant publications. The focus was on research or review articles, work on NPs isolated from this weed. Many of the publications were focused on isolation, structure elucidation and pharmacological activities.

Natural products derived from C. rotundus L. species

After decades of detailed phytochemical investigation, it is evident that this plant species contains two major classes of secondary metabolites, namely, terpenoids and flavonoids



Dr Stefan Günther studied biology and informatics in Germany and was appointed to a JuniorProfessor in Pharmaceutical Bioinformatics in 2009 and to a Full Professor in 2015 at the University Freiburg, Germany. His research area is the development and application of methods from bioinformatics in pharmaceutical sciences. He has a special focus on structurebased drug discovery and the

prediction of theeffects of natural products for therapeutic application.



Fidele Ntie-Kang heads the Molecular Simulations Laboratory, Chemistry Department, University of Buea. He studied Chemistry at the University of Douala (Cameroon) from 1999 to 2005, leading to Bachelor's and Master's degrees. His PhD from University oof Douala (Caeroon) was based on molecular modeling of anti-tubercular drug target to design novel inhibitors, followed by an

Habilitation in Pharmaceutical Chemistry from Martin-Luther University Halle-Wittenberg (Germany), under the supervision of Prof. Wolfgang Sippl. His current focus is the discovery of bioactive natural products from African flora by the use of virtual screening followed by in vitro assays. A major contribution of his research team has been the development of the African natural products database.




Fig. 3 Structures of terpenoids isolated from Cyperus rotundus (1 to 34)

mostly harvested from Asia and Africa (Fig. 2). $^{8,12,13,17,40,79-87,89-94,112-119}$

3.1 Terpenoids

To date, a total of about 131 terpenoids (1–131) have been isolated and identified from *C. rotundus* (CR) (Fig. 3–7). The summaries of the most interesting results for terpenoids isolated from CR have been shown in Table 1. Sesquiterpenes are the major subclass of NPs isolated from this herb. A majority of the secondary metabolites were isolated from the rhizomes/tubers of the plant and they are structurally related, for example, (1–117). 83,120 α -Cyperone (17) isolated from the n-

hexane fraction significantly inhibited prostaglandin E2 (PGE2) production by suppressing lipopolysaccharide (LPS)-induced expression of inducible cyclooxygenase-2 (COX-2) at both RNA and the protein levels. Compound 17 obtained from the tubers of this plant species also showed insecticidal activity. Socyperol (18) has been found to significantly inhibit LPS-induced production of nitrite oxide (NO), PGE2 and suppressed LPS-induced expression of inducible nitric oxide synthase (iNOS) and COX-2 at the mRNA and protein levels in RAW 264.7 macrophages. Extraction of air-dried and chopped rhizomes of CR with hot 70% EtOH followed by purification using GC-MS afforded monoterpenes, sesquiterpenes and

Review

Fig. 4 Structures of terpenoids isolated from Cyperus rotundus (35 to 61).

aromatic compounds (10, 17, 37–43). The sesquiterpenes; valencene (10) and (+)-nootkatone (40) significantly inhibited inducible nitric oxide (iNOS) expression and nitric oxide (NO) production in LPS-simulated RAW264.7 cells. The anti-inflammatory mechanism of CR is due to heme oxygenase-1 (HO-1) induction by compounds 10 and 40.¹⁷ While (+)-nootkatone (40) has been found to have potent inhibitory effect on collagen-, thrombin-, and AA-induced platelet aggregation. Compound 40 was treated with mice and it exhibited significant prolonged bleeding times. It has also shown significant inhibitory effect on rat platelet aggregation *ex vivo*.⁷⁷

Three novel sesquiterpene alkaloids; rotundines A (44), B (45), and C (46) were isolated from the MeOH extract using

standard methods of extraction of alkaloids. The structures of the compounds were determined by comprehensive spectroscopic analyses and chemical methods.¹⁵

61

Ohira *et al.*⁸² isolated the new sesquiterpenoids; 2α -(5-oxopentyl)- 2β -methyl- 5β isopro-penylcyclohexanone (48), 2β -(5-oxopentyl)- 2β -methyl- 5β -isopropenylcyclohexanone (49), cyperolone (50) together with the known compounds 17, 19, 40, 51 and 52 from the roots of CR. The antibacterial activities of the new hits were screened against *Escherichia coli* and *Bacillus subtilis* using the paper disk method. Cyperolone (50) possessed moderate activity against *B. suhtilis* at a concentration of 0.5 mg per disk; the other compounds did not show notable activities.⁸²


Fig. 5 Structures of terpenoids isolated from Cyperus rotundus (62 to 94).

The new sesquiterpenes, cyperusol A3 (53), 3β-hydroxycyperenoic acid (54), along with three known compounds (55–57) were isolated from the ethyl acetate soluble fraction of the rhizomes using a series of column chromatography. The compounds were submitted for their cytotoxic activities against human ovarian cancer cells (A2780) and endometrial adenocarcinoma cells (Ishikawa) using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2*H*-tetrazolium bromide (MTT) assays and 11,12-dihydroxyeudesm-4-en-3-one (57) showed the most potent

cytotoxic activity with observed IC_{50} values of 11.06 and 6.46 mm, respectively.¹²

Jin *et al.*¹⁰ isolated the known compounds; valencene (**10**), α-cyperone (**17**), β-pinene (**37**), limonene (**38**), 4-cymene (**39**), (+)-nootkatone (**40**), 1, 8-cineole (**41**), caryophyllene oxide (**42**), and β-selinene (**43**) and evaluated them for their anti-allergic activity *in vitro* and *in vivo*. In rat basophilic leukemia (RBL)-1 cells, the sesquiterpenes (**10**, **40**, **42**) were reported to strongly inhibit 5-lipoxygenase-catalyzed leukotrienes production. In addition, they inhibited β-hexosaminidase release by antigen-

Review **RSC Advances**



Structures of terpenoids isolated from Cyperus rotundus (95 to 116)

stimulated RBL-2H3 cells, with valencene having the highest inhibitory effect. Authors also found that the most active sesquiterpene (10) inhibited β-hexosaminidase degranulation by inhibiting the initial activation reaction, Lyn phosphorylation, in IgE-stimulated RBL-2H3 cells. Moreover compounds (10, 40), significantly inhibited the delayed-type hypersensitivity reaction in mice when administered orally at 50-300 mg kg⁻¹. 11

The isolation of cyperolone (50) from the essential oil of CR, rekindled the interest of NP Chemists to revisit this plant species. Investigation of the constituents from Chinese origin, led to the isolation of known compounds (19, 50) and new sesquiterpenes; copadiene (58), epoxyguaien (59), rotundone (60) and $4\alpha,5\beta$ -oxidoeudesm-11-en-3 α -ol (61). 79,124-127

Bioactivity and liquid chromatography-mass spectrometry (LC-MS) guided fractionation of 90% EtOH extract using opencolumn, Sephadex LH-20 and semi-preparative high performance liquid chromatography (HPLC) led to the isolation and identification of thirty-seven sesquiterpenoids. 13 compounds include, five new patchoulane-type sesquiterpenoids, 3β-hydroxycyperenoic acid (54), cyperene-3,8-dione (62), 14-hydroxycyperotundone (63), 14-acetoxycyperotundone (64) and sugetriol-3,9-diacetate (65) along with the known NPs 4, 17, 20, 27, 36, 51 and 66-89. Nine eudesmane-type sesquiterpenoids (20, 71-77 and 78-80) significantly inhibited the hepatitis B virus (HBV) DNA replication with IC₅₀ values of 42.7 \pm 75.9, 22.5 ± 71.9 , 13.2 ± 71.2 , 10.1 ± 70.7 , 14.1 ± 71.1 , 15.3 ± 72.7 , 13.8 \pm 70.9, 19.7 \pm 72.1 and 11.9 \pm 70.6 μ M, respectively, of which, compounds 72, 76, 78 and 80 possessed high selectivity index (SI) values of 250.4, 125.5, 259.6 and 127.5, respectively. Two patchoulane-type sesquiterpenoids (54 and 36) effectively suppressed the secretion of HBsAg in a dose-dependent manner with IC₅₀ values of 46.6 \pm 714.3 (SI = 31.0) and 77.2 \pm 713.0 (SI = 1.7) μM, respectively. Compounds 63, 4, 17, 20, 72 and 81 possessed moderate activities against HBeAg secretion with IC50 values of 162.5 \pm 718.9 (SI = 13.3), 399.2 \pm 790.0 (SI = 10.6), 274.7 \pm 770.8 (SI = 5.2), 313.9 \pm 787.5 (SI = 7.2), 334.0 \pm 770.4 (SI = 9.9) and 285.3 \pm 720.9 (SI = 15.5) μ M, respectively.¹³

Antimalarial activity-guided investigation and HPLC separation of the crude hexane extract of CR tubers led to the isolation of the sesquiterpenes; 42, patchoulenone (90) and 10,12-peroxycalamenene (91). The antimalarial activities of these compounds were determined from their effective concentrations (EC50) values against Plasmodium falciparum. The in vitro activity against P. falciparum (EC₅₀) of compounds RSC Advances Review

$$R_{1} = \beta \text{-Ara-}(1 \longrightarrow 6) \text{-Kyl-}(1 \longrightarrow 4) \text{-Gic} \quad R_{2} = \beta \text{-Gic} \quad 117$$

$$R_{1} = \beta \text{-Ara-}(1 \longrightarrow 6) \text{-Kyl-}(1 \longrightarrow 4) \text{-Gic} \quad R_{2} = \text{H} \quad 118$$

$$R = \beta \text{-Gic} \quad 121$$

$$R = \beta \text{-Gic} \quad 121$$

$$R = \beta \text{-Gic} \quad 121$$

$$R = \beta \text{-Gic} \quad 122$$

$$R = \beta \text{-Gic} \quad 123$$

$$R = -\text{Nechesperidoside} \quad 124$$

$$R = \beta \text{-Gic} \quad 129$$

$$R = Ara-(2) \longrightarrow 7 \quad 130$$

$$R = \text{Gal} \quad 131$$

Fig. 7 Structures of terpenoids isolated from Cyperus rotundus (117 to 131).

90 and **91** were in the order of 10^{-4} M (1.08×10^{-4} and 3.45×10^{-4} M respectively).⁸¹

Chemical investigation of the rhizomes of CR led to the isolation of a new norsesquiterpenes, named norcyperone (92), (–)-clovane-2,9-diol (93) and other known compounds. The structure of the novel norsesquiterpene (92) with a tetrahydrofuran ring structure was established on the basis of extensive spectroscopic analyses, including 1D- and 2D-NMR, MS experiments, and single-crystal X-ray diffraction.⁸⁴

A novel norsesquiterpenoid named cyperalin A (94) with the known compound (27) were isolated from freshly-cut rhizomes of CR defatted with n-hexane followed by extraction with methanol and silica gel chromatography. The compounds were screened for their anti-inflammatory activity. Cyperalin A (94) displayed the highest inhibitory activity of PGE2, COX-2, and LOX-5 with IC₅₀s 0.22, 1.03, and 1.37 μ M, respectively compared to indomethacin (IC₅₀s 0.15, 0.69, and 0.81 μ M, respectively). Compound 27, showed significant activity with IC₅₀s 0.57 (PGE2), 1.74 (COX-2) and 2.03 (LOX-5) μ M.⁹³

The new sesquiterpenes, $4\alpha,5\alpha$ -oxidoeudesm-11-en-3-one (95) and cyper-11-ene-3,4-dione (96) together with the known

compounds **4**, **17**, **18** and **47** were obtained from the hexane and dichloromethane fractions of CR. The compounds were examined for their estrogenic activity by E-screen assay on MCF-7 BUS cells. Compound **2**, exhibited the most potent estrogenic activity by increasing transcriptional activities in an estrogen sensitive reporter gene assay. The authors showed that compound **2**, has biphasic activities on estrogen receptors which could be useful as an alternative hormone replacement therapy. Compounds **4** and **17** isolated from the tubers of CR have showed growth inhibitory effects to both shoots and roots on the lettuce seedlings. Les

Rani *et al.*⁸⁷ isolated the sesquiterpenes **40**; solavetivone (97) and aristolone (98) from the acetone extract of CR by silica gel column chromatography and determine their radical scavenging potential compared with standard gallic acid. Among the three sesquiterpenoids isolated, compound **40** possessed the highest radical scavenging potential (IC₅₀ 4.81 g mL⁻¹) followed by **98** (IC₅₀ 5.28 g mL⁻¹) and the new compound 97 (IC₅₀ 6.82 g mL⁻¹) by DPPH radical scavenging assay.⁸⁷

Bioassay-guided fractionation of the methanol extracts of the rhizomes of CR led to the isolation of **41**, **42**, zerumbone (**99**), α -

Table 1 Summary of the bioactivity of derived terpenoids from Cyperus rotundus^a

Compounds	Part (s)/place of harvest of plant studied (VS \neq)	Measured activity	References
Rotundene (1)	NM/China	ND	Paknikar et al., ¹²⁰
1, (-)-cypera-2,4-diene (2), cyprotene (3), cyperotundone (4), (+)-cyperadione (5), (-)-norrotundene (6), (-)-isorotundene (7), γ -gurjunene (8), nootkatene (9), valencene (10), $epi-\alpha$ -selinene (11), α -muurolene (12), γ -muurolene (13), ylanga-2,4-diene (14), γ -calacorene (15), cadalene (16), α -cyperone (17), isocyperol (18), mustakone (19) and cyperol (20) and (-)-cypera-2,4(15)-diene (21)	NM/gift of KD. Protzen, Paul Kadersd GmbH, Hamburg (NM)	ND	Sonwa et al., ⁸³
3,4- <i>O</i> -isopropylideneshikimic acid (22), rotundusolide A (23), rotundusolide B (24), dehydrocostuslactone (25), (+)-alismoxide (26), sugetriol triacetate (27), 2β-hydroxy-α-cyperone (28), eudesma-4(14),11(13)-diene-7α,8α,12-triol (29), rotundusolide C (30), secomacrogenin B (31), and 3,4-seco-mansumbinoic acid (32)	Rhizomes/Purchased from Lanzhou Traditional Chinese Medicine Market (no. ZY2009C002)	ND	Yang et al., ⁸⁹
4, 27, <i>Epi</i> -guaidiol A (33), guaidiol A (34), sugebiol (35) and cyperenoic acid (36)	Rhizomes/Da-Bie-Shan Mountains, Anhui province, China. (No: 20060825)	ND	Xu et al.,85
10 , 17 , β-pinene (37), limonene (38), 4-cymene (39), (+)-nootkatone (40), 1, 8-cineole (41), caryophyllene oxide (42), β-selinene (43)	Rhizomes/Purchased from Kyung Dong market place in Seoul, South Korea. (DKH- 02561)	Anti-inflammatory activity	Tsoyi <i>et al</i> ., ¹⁷
10,17, 37- 43 18	Rhizomes/NM Rhizomes/Kyung Dong Crude Drugs Market, Seoul, South Korea. (CYRO1-2011)	Antiplatelet effects Anti-inflammatory activity	Seo <i>et al.</i> , ⁷⁷ Seo <i>et al.</i> , ¹²³
17	Rhizomes/Kyung Dong Crude Drugs Market, Seoul, South Korea. (CYRO1-2011)	Anti-inflammatory activity	Jung et al., ¹²¹
Rotundine A (44), rotundine B (45), and rotundine C (46)	Rhizomes/Purchased from Uchida Wakanyaku Co., Ltd. (Tokyo, Japan) (lot. 242 118)	ND	Jeong et al., ¹⁵
α-Cyperene (47) 17, 19, 40, 2α-(5-oxopentyl)-2β-methyl-5β- isopropenylcyclohexanone (48), 2β-(5-oxopentyl)-2β- methyl-5β-isopropenylcyclohexanone (49), cyperolone (50), α-rutunol (51) and β-rutunol (52)	NM/China (NM) Roots/Yorishima, Okayama, Japan	ND Antibacterial activity	Komai <i>et al.</i> , ⁸⁰ Ohira <i>et al.</i> , ⁸²
Cyperusol A ₃ (53), 3 β -hydroxycyperenoic acid (54), britanlin E (55), 1 β , 4 α -dihydroxyeudesm-11-ene (56) and 11,12-dihydroxyeudesm-4-en-3-one (57).	Rhizomes/Kyung Dong Crude Drugs Market, Seoul, South Korea. (CYRO1–2011)	Antitumor activity	Ryu et al., 12
10, 37-42	Rhizomes/Kyung Dong Crude Drugs Market, Seoul, South Korea. (NM)	Anti-allergic activity	Jin et al.,10
Copadiene (58), epoxyguaien (59), and rotundone (60)	NM/Japan (NM) NM/India (NM)	ND	Kapadia <i>et al.</i> , ⁷⁹ Kapadia <i>et al.</i> , ¹²
19 50	Rhizomes/Japan (NM)	ND ND	Hikino et al., 125
4α,5β-oxidoeudesm-11- <i>en</i> -3α-ol (61)	Rhizomes/Japan (NM)	ND	Hikino et al., 127
51 and 52	Rhizomes	ND	Hikino et al., 126
4, 17, 20, 27, 36, 51, 54, cyperene-3,8-dione (62), 14-hydroxycyperotundone (63), 14-acetoxycyperotundone (64), sugetriol-3,9-diacetate (65), cyperenol (66), sugeonol (67), scariodione (68), (4a <i>S</i> , 7 <i>S</i>)-7-hydroxy-1,4a-dimethyl-7-(prop-1-en-2-yl)-4,4a,5,6,7,8-hexahydronaphthalen-2(3 <i>H</i>)-one (69), (4a <i>S</i> , 7 <i>S</i> , 8R)-8-hydroxy-1,4a-dimethyl-7-(prop-1-en-2-yl)-4,4a,5,6,7, 8-hexahydronaphtha-len-2(3 <i>H</i>)-one (70), 1β-hydroxy-α-cyperone (71), 10-epieudesm-11-ene-3β, 5α-diol (72), 3β-hydroxyilicic alcohol (11(13)-eudesmene-3,4,12-triol) (73), cyperusol C (74), α-corymbolol (75), 3β, 4α-dihydroxy-7- <i>epi</i> -eudesm-11(13)-ene (76), 2-oxo-α-cyperone (77), 7α (<i>H</i>), 10β-eudesm-4- <i>en</i> -3-one-11,12-diol (78), 2-hydroxy-14-calamenenone (79), rhombitriol (80), 7-	Rhizomes/Purchased from Juhuacun (Kunming, China) (no. 2011041101)	Anti-hepatitis B virus activity	Xu et al., ¹³

Table 1 (Contd.)

Compounds	Part (s)/place of harvest of plant studied (VS \neq)	Measured activity	References
oplopanone (83), 10-hydroxyamorph-4- <i>en</i> -3-one (84), cyperusol D (85), argutosine D (86) and 4,5-seco-guaia-1 (10),11-diene-4,5-dioxo (87), oxyphyllol C (88) and 5-bydroxylygian (80)			
hydroxylucinone (89) 42, patchoulenone (90) and 10,12-peroxycalamenene (91)	Tubers/Purchased from a Thai traditional dispensary, Bangkok, Thailand	Antimalarial activity	Thebtaranontl
Norcyperone (92) and (–)-clovane-2,9-diol (93)	Rhizomes/Dabieshan Mountains of Anhui Province, P.R. China (no: 20060825)	ND	Xu et al.,84
27 and cyperalin A (94)	Rhizomes/King Abdulaziz University campus, Jeddah, Saudi Arabia (2014-CR110)	Anti-inflammatory activity	Ibrahim et al.,
4, 17, 18, 47 , 4α,5α-oxidoeudesm-11 <i>-en</i> -3-one (95) and cyper-11-ene-3,4-dione (96)	Rhizomes/purchased from Kyungdong-Yakryongsi traditional medicine market in Seoul, Korea (SKKU-PH-12-50)	Estrogenic activity	Park et al., ⁹⁴
40, solavetivone (97) and aristolone (98)	Rhizomes/obtained from a registered medicinal plant vendor in Trivandrum (no. 034/2011)	Antioxidant activity	Rani <i>et al.</i> , ⁸⁷
17 41, 42, zerumbone (99), α-humulene (100), (+)-dihydrocarvone (101), (R)-(+)-limonene (102), (1S)- (-)-verbenone (103), (S)-(-)-limonene (104), β- caryophyllene (105), (-)-(E)-pinocarveol (106), (E)-carveol (107), (-)-α-copaene (108), (1R)-(-)-myrtenal (109) and (1R)-(-)-myrtenol (110)	Tubers/Bogor-Indonesia/NM Rhizomes/Purchased from the Boeun medicinal herb shop (Seoul Yangnyeongsi, Seoul, South Korea) (CR-01)	Insecticidal activity Repellent activity	Dadang et al., ⁹² Chang et al., ⁹²
4 and 17	Rhizomes/supplied by Morihiro Kinoshita (Nihon Funmatsu Yakuhin Company, Japan) (NM)	Phytotoxicity	Morimoto et al., 128
4 , 17 , 20 , 43 , 47 , 101 , 109, β-elemene (111), caryophyllene (112), δ-cadinene (113), calamenene (114), patchoulenyl acetate (115) and 6-acetoxy-patchoul-4-en-3-one (116)	Tubers/Islands of Oahu, Maui, Kauai and Hawaii (NM)	ND	Komai <i>et al.</i> , ⁸⁰
Cyprotuside A (117) and cyprotuside B (118)	Rhizomes/Zhanjiang, Guangdong Province of China (No.20090903)	Antidepressant activity	Zhou et al.,90
18- Epi - α -amyrin glucuronoside (119), oleanolic acid arabinoside (120), α -amyrin glucopyranoside (121) and β -amyrin glucopyranoside (122)	Tubers/West Champaran, Bihar, India (no. NISCAIR/ RHMD/Consult/-2008-09/1114/ 145)	ND	Alam et al.,86
Oleanolic acid (123), oleanolic acid-3- <i>O</i> -neohesperidoside (124) and β-sitosterol (125)	Tubers/University of Allahabad campus, Allahabad, India (NM)	ND	Singh et al., 114
2-Methyl cyprot-3-en-2-one-13-oic acid (126), tigmasterol- <i>n</i> -dodecanoate (127), stigmasterol- <i>n</i> - etradecanoate (128), β-sitosterol glucoside (129) and upenyl arabinopyranosyl oleate (130)	Tubers/Purchased from a Delhi market	ND	Sultana et al.,
Sitosteryl (131)	Aerial parts/El-Safa and El- Marwa, Faculty of Agriculture, Al-Azhar University, Assiut, Egypt.	Antifeedant and Cytotoxic activity	Sayed et al., ³³

humulene (100), (+)-dihydrocarvone (101), (R)-(+)-limonene (102), (1S)-(-)-verbenone (103), (S)-(-)-limonene (104), S-caryophyllene (105), (-)-(E)-pinocarveol (106), (E)-carveol (107), (-)- α -copaene (108), (1S)-(-)-myrtenol (109) and (1S)-(-)-myrtenol (110). The compounds were tested for repellency to male

Blattella germanica and the results were compared to N,N-diethyl-3-methylbenzamide (deet). In filter-paper choice tests, 99 was the most repellent compound, and 100 was ineffective, which shows that α,β -unsaturated carbonyl group of 99 contributes to repellency. The article reported that at 81.5 μg

Table 2 Summary of the bioactivity of derived flavonoids from Cyperus rotundus⁴

Compounds	Part (s)/place of harvest of plant studied (VS \neq)	Measured activity	References
Luteolin-7- <i>O</i> -glucoside (132), tricin (133) and aureusidin (134)	Leaves/stem, Australia (K. L. Wilson 3309) (Denistone, N.S.W)	ND	Harborne et al., 141
Afzelechin (135), (+)-catechin (136), luteolin (137), and quercetin (138)	Aerial parts/Monastir region in the Center of Tunisia (Cp.10.04)	Antioxidant and antitumor activities	Kilani-Jaziri <i>et al.</i> , ⁴⁰
Rutin (139)	Rhizomes/Al-Azhar University campus, Assiut Branch, Egypt (2009- CR110)	Hepatoprotective activity	Mohamed, ⁷⁸
Pongamone A (140) and biochanin A (141)	Rhizomes/Collected in Zhanjiang, Guangdong Province of China (No.20090903)	ND	Zhou et al., 11
137	Rhizomes/Purchased from Matsuura-Yakugyo Co. Ltd. (Nagoya, Japan) CP-0901	Antiproliferative activity	Ito et al., ¹⁸
133, isorhamnetin (142), vitexin (143), isovitexin (144), oreintin (145), epiorientin (146), myricetin-3- <i>O</i> -β-p-galactopyranoside (147), luteolin-7- <i>O</i> -β-p-glucuronopyranoside-6"-methyl ester (148), luteolin-4'- <i>O</i> -β-p-glucuronopyranoside (149) and Luteolin-7- <i>O</i> -β-p-glucuronopyranoside (132)	Aerial parts/Experimental Station of El-Safa and El-Marwa, Faculty of Agriculture, Al-Azhar University, Assiut, Egypt	Antioxidant and α-amylase inhibitory activities	Sayed <i>et al.</i> , ⁸
133 and 143	Aerial parts/Experimental Station of El-Safa and El-Marwa, Faculty of Agriculture, Al-Azhar University, Assiut, Egypt (NM)	Antifeedant and Cytotoxic activity	Sayed et al., ³³
143, 145, cinaroside (150), quercetin- <i>O</i> -β-D- glucuronopyranoside (151), cyperaflavoside (152) and myricetin- 3- <i>O</i> -β-D-glucuronopyranoside (153)	Aerial parts/King Abdulaziz University campus, Jeddah, Saudi Arabia (2014-CR110)	5-Lipoxygenase inhibitory activity	Ibrahim <i>et al.</i> , ⁹³
a VC Voucher angeimen number NN			

^a VS≠: Voucher specimen number, NM: not mention, ND: not done.

cm $^{-2}$, enhanced repellency was produced by binary mixtures of **99** and **41**, **101** or **102** (70:30, 50:50 and 30:70 ratios by weight). In Ebeling choice box tests at 652.4 µg cm $^{-2}$, these compounds and deet resulted in complete repellency to intact male *B. germanica*, while they exhibited 35–47% repellency to antennectomized male one. Mixtures of active compounds from this plant species could serve as potential repellents for controlling *B. germanica*. ⁹²

A series of mono and sesquiterpenoids, 4, 17, 20, 43, 47, 101, 109, β -elemene (111), caryophyllene (112), δ -cadinene (113), calamenene (114), petchoulenyl acetate (115) and sugeonyl acetate (116) were isolated from the mature tubers of CR using gas chromatography-mass spectrometry (GC-MS). A new chemotype of CR was found in Hawaii based on the sesquiterpene composition of the mature tubers. The K-type has higher concentrations of 115 and 116 than the three known Asian chemotypes. Information on the distribution of chemotypes could also offer clues to the history of spreading CR weed species. ⁸⁰

Zhou *et al.*⁹⁰ isolated the two novel cycloartane glycosides, cyprotuside A (117) and cyprotuside B (118) from the rhizomes

of CR and evaluated their antidepressant activity by forced swimming test (FST) and tail suspension test (TST) in mice. The preliminary *in vivo* evaluation showed that compounds **117** and **118** exhibited remarkable antidepressant activity in the despair mice models.

Exhaustive extraction of air-dried powdered tubers of CR with methanol followed by chromatography over silica gel column and elution with a gradient of chloroform and methanol afforded two new triterpenic glucosides; 18-epi- α -amyrin glucuronoside (119), oleanolic acid arabinoside (120), and α -amyrin glucopyranoside (121) and β -amyrin glucopyranoside (122).

The sesquiterpenoid, 12-methyl cyprot-3-en-2-one-13-oic acid (126), steroidal esters; stigmasterol-*n*-dodecanoate (127), stigmasterol-*n*-tetradecanoate (128), β-sitosterol glucoside (129) and triterpenoid glycosides; lupenyl arabinopyranosyl oleate (130) were isolated for the first time and they could serve as chromatographic markers for standardization of the tubers of CR.⁹¹ A new steroid glycoside; sitosteryl (131) and other compounds were also isolated from the aerial parts of this plant species harvested in Egypt.³³

Fig. 8 Structures of flavonoids isolated from Cyperus rotundus (132 to 149).

R = β-GlcA-6"-methyl ester

3.2 Flavonoids

Flavonoids continue to attract attention as potentially useful compounds because of their broad spectrum of biological activities. ^{129–140} In this report, summaries of the most interesting results for flavonoids (132–153) isolated from CR have been shown in Table 2, while the chemical structures of the isolated compounds are shown in Fig. 8 and 9. In Table 2, the biological activities of the compounds and the organism studied have been provided.

Harborne *et al.*¹⁴¹ isolated luteolin-7-O-glucoside (132), tricin (133) and aureusidin (134) from the leaves/stems of CR by paper electrophoresis. The structures of the compounds were

identified by standard procedures and co-chromatography with authentic samples carried out in at least 4 solvents.

149

In an effort to search for novel antioxidant and antiproliferative hits, four flavonoids; afzelechin (135), (+)-catechin (136), luteolin (137), and quercetin (138) and other compounds were isolated from the total oligomers flavonoids and ethyl acetate extracts of CR. Compound 137 was the most active in reducing the production of thiobarbituric acid reactive substances (malondialdehyde = 1.5 nM), inhibiting significantly the proliferation of K562 cells ($IC_{50} = 25 \text{ g mL}^{-1}$) and protecting against H_2O_2/UV -photolysis induced DNA damage.⁴⁰ Rutin (139), pongamone A (140) and biochanin A (141) were

Review **RSC Advances**

Structures of flavonoids isolated from Cyperus rotundus (150 to 153).

isolated from the rhizomes of this plant species and the structure was established on the basis of 1D and 2D NMR spectroscopic analyses. 11,78 Luteolin (137) has been isolated from the rhizomes of this plant species.18

Chemical investigation of the aerial parts of CR led to the isolation 133, isorhamnetin (142), vitexin (143), isovitexin (144), oreintin (145), epiorientin (146), myricetin-3-O-β-D-galactopyranoside (147), luteolin-7-O-β-D-glucuronopyranoside-6"-methyl ester (148), luteolin-4'-O-β-p-glucuronopyranoside (149) and luteolin-7-O-β-D-glucuronopyranoside (132). Compounds (146-148) having an ortho-dihydroxyl groups at ring B showed strong antioxidant activity. The presence of a hydroxyl group at position-3 increased the antioxidant activity in 143. Compounds (133, 143, 144, 146, 148) demonstrated strong α -amylase inhibitory activity > 50% influenced by the presence of free hydroxyl groups at 7, 3' and 4' positions. 8,33 The flavonoids 144, **146**, cinaroside (**150**), quercetin-*O*-β-D-glucuronopyranoside (151), cyperaflavoside (152) and myricetin-3-O-β-D-glucuronopyranoside (153) were also isolated from the aerial parts and screened for their 5-lipoxygenase inhibitory potential. All compounds demonstrated 5-lipoxygenase inhibitory potentials with IC₅₀s 5.1 (152), 4.5 (143), 5.9 (145), 4.0 (150), 3.7 (151), and 2.3 μ M (153), respectively, in comparison to indomethacin (IC₅₀ $= 0.98 \text{ M}).^{93}$

Table 3 Summary of the bioactivity of derived stilbenes and derivatives from Cyperus rotundus

Compounds	Part (s)/place of harvest of plant studied (VS \neq)	Measured activity	References
(+)-Cyperusphenol A (154), (-)-(E)-cyperusphenol A (155), (E)-mesocyperusphenol A (156), cyperusphenol C (157), cyperusphenol B (158), cyperusphenol D (159), transscirpusin A (160) and scirpusin B (161)	Rhizomes/Purchased from Matsuura-yakugyo Co. Ltd. (Nagoya, Japan) (CP-0901)	Antiproliferative activity	Ito et al., ¹⁸
4,7-Dimethyl-1-tetralone (162)	Tubers/Purchased from a Thai traditional dispensary, Bangkok, Thailand	Antimalarial activity	Thebtaranonth et al.,81
p-Hydroxybenzoic acid (163)	Rhizomes/Purchased from Uchida Wakanyaku Co., Ltd. (Tokyo, Japan) (lot. 242118)	ND	Jeong et al., ¹⁵
Salicylic acid (164), caffeic acid (165), protocatechuic acid (166) and p-coumaric acid (167)	Aerial parts/El-Safa and El-marwa, Faculty of Agriculture, Al-Azhar University, Assiut, Egypt	Antifeedant and Cytotoxic activity	Sayed et al., ³³
167, and ellagic acid (168)	Rhizomes/Purchased from Kyung Dong market place in Seoul, South Korea. (DKH-02561)	Anti-inflammatory activity	Tsoyi et al., ¹⁷
164–167, methoxycyperotundol (169) and cyperotundol (170)	Rhizomes/Collected in Zhanjiang, Guangdong Province of China (no. 20090903)	ND	Zhou et al., ¹¹
Galloylquinic acid (171), 3-hydroxy- 4-methoxybenzoic acid (172) and ferulic acid (173)	Aerial parts/Monastir region in the Center of Tunisia (Cp.10.04)	Antioxidant and antitumor activities	Kilani-Jaziri <i>et al.</i> , ⁴⁰
Chlorogenic acid (174)	Rhizomes/obtained as Organic Musta Powder (Khandige Organic Health Product, Bangalore, India) (NM)	Anti-inflammatory activity	Rocha <i>et al</i> ., ¹⁰
Methyl-3,4-dihydroxybenzoate (175)	Rhizomes/Al-Azhar University campus, Assiut Branch, Egypt (2009- CR110)	Hepatoprotective activity	Mohamed, ⁷⁸

^a VS≠: Voucher specimen number, NM: not mention, ND: not done.




Fig. 10 Structures of stilbenes, ellagic acid derivatives and other compounds from Cyperus rotundus (154 to 170).




Fig. 11 Structures of stilbenes and derivatives isolated from Cyperus rotundus (171 to 175).

Table 4 Miscellaneous compounds from Cyperus rotundus^a

Compounds	Part (s)/place of harvest of plant studied (VS≠)	Measured activity	References
n-Butyl-β-p-fructopyranoside (176), ethyl-α-p-glucopyranoside (177), adenosine (178) and (–)-(E)- caffeoylmalic acid (179)	Aerial parts/Experimental Station of El-Safa and El-Marwa, Faculty of Agriculture, Al-Azhar University, Assiut, Egypt	Antioxidant and α-amylase inhibitory activities	Sayed et al.,8
Benzo-α-pyrone (180), khellin (181), visnagin (182), ammiol (183) and khellol-β-p-glucopyranoside (184)	Aerial parts/El-Safa and El-Marwa, Faculty of Agriculture, Al-Azhar University, Assiut, Egypt	Antifeedant and Cytotoxic activity	Sayed et al., ³³
Ipolamiide (185) and 6β- hydroxyipolamiide (186)	Rhizomes/Al-Azhar University campus, Assiut Branch, Egypt (2009- CR110)	Hepatoprotective activity	Mohamed, ⁷⁸
1-O-(β-ɒ-glucopyranosyloxy)- (2S,3R,4E,8Z)-2-[(2'R)-2'- hydroxylignoceranoylamino]-4,8- tetradecene-3-diol (187)	Radix/Shandong, China	Anti-proliferation activity	Liu <i>et al.</i> , 143
n-Tricont-1-ol-21-one (188)	Tubers/West Champaran, Bihar, India (No.NISCAIR/RHMD/ Consult/-2008-09/1114/145)	ND	Alam et al., 86
Succinic acid (189), myristic acid (190), palmitic acid (191) and stearic acid (192)	Tubers/upland rice field (sandy loam soil) of Manikganj district, Bangladesh (NM)	Growth inhibitory effects	Quayyum et al., 12

R =β-Glc

Fig. 12 Structures of miscellaneous compounds isolated from Cyperus rotundus (176 to 192).

RSC Advances Review

3.3 Stilbenes and derivatives

Stilbenes are polyphenols containing resveratrol as a basic subunit. These compounds have received much attention because of their cardioprotective effects, but they also display anti-inflammatory, antioxidative, and antimicrobial activities. They are also known as anticancer and cancer-chemopreventive agents. ¹⁴² The stilbenes and derivates (154–176) isolated from CR by several NP research groups have shown interesting biological activities during *in vitro* screening exercise (Table 3). ^{11,17,40,77,78,110} The chemical structures of those isolated from this herb are shown in Fig. 10 and 11.

The novel enantiomeric and meso-stilbene trimers; (+)-cyperusphenol A (154), (-)-(E)-cyperusphenol A (155), (E)mesocyperusphenol A (156), a trimer bearing a novel hexacyclic ring system, cyperusphenol B (158), together with the known stilbenoids, cyperusphenol C (157), cyperusphenol D (159), trans-scirpusin A (160) and scirpusin B (161) were isolated from the rhizomes of CR. The hits were evaluated for their antiproliferative activity employing the Jurkat cell line (human Tcell leukemia cells), while the IC₅₀ potencies of a racemate of compounds 154-156, 158, and 159 were estimated as 27.4, 40.5, 26.4, and 26.3 µM, respectively. The article reported that the suppression of cell growth by compound 159 was due to the induction of apoptosis, which was characterized by nuclear changes and PARP-1 cleavage determined by western blotting.18 Salicylic acid (164), caffeic acid (165), protocatechuic acid (166) and p-coumaric acid (167) isolated from this weed showed significant antioxidant activity.8 The insect antifeedant activity demonstrated by the crude extracts is due to the presence of furochromones having methoxyl group at positions C-5 and/or C-8 positions.33

3.4 Miscellaneous compounds

Steroidal glycosides, furochromones and aromatics (177–185) isolated from the aerial parts of CR collected from Egypt demonstrated antioxidant, α -amylase inhibitory and antifeedant activities. ^{8,33} A summary of these bioactive compounds isolated from CR is provided in (Table 4) and their chemical structures in Fig. 12. New iridoids, cerebroside, known aliphatic fatty acids and coumarin (186–192) from this plant species have also shown hepatoprotective, anti-proliferation, and growth inhibitory properties respectively. ^{78,86,143,144}

3.5 Bioactivities and proposed mechanisms of *C. rotundus* compounds

The pharmacological activities and mechanisms of action of some compounds isolated from *C. rotundus*have been reported extensively.^{8,11,17,41,76-78} The summaries of the most interesting results for some NPs isolated from this weed have been shown in Fig. 12.

Conclusions

Cyperus rotundus L. (Nutgrass, family Cyperaceae) popularly called "the world's worst weed" has attracted particular attention as a medicinal plant, due its broad spectrum of

pharmacological activities. In the past six decades, about 192 NPs have been isolated and characterized from this plant species. Among them, terpenoids and flavonoids are the major bioactive constituents mostly harvested from Asia and Africa. The chemical structures of pure compounds were retrieved from literature sources comprising data collected from articles from major peer-reviewed journals, from all over the world spanning the period 1965 to 2020. The collected data includes region of collection of plant material, voucher specimen number, isolated metabolites and class, and measured biological activities of isolated compounds. The study has provided a survey of the biological activities of 192 NPs and the mechanism of action of some of the compounds isolated from C. rotundus. It is worth mentioning that C. rotundus and its NPs have shown good safety in vitro and in vivo studies. Thus it would be interesting in future to evaluate the toxicities of the NPs from this weed using in silico approaches.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

SBB acknowledges the African-German Network of Excellent in Science (AGNES) junior researcher grant supported by Alexander von Humboldt and sponsored by the Federal Ministry of Education and Research. SBB would like to thank the Agro-EcoHealth Unit, International Institute of Tropical Agriculture (IITA), Cotonou, Benin fortraining in HPLC. FNK would also like to acknowledge funding from the German Academic Exchange Services (DAAD) for a guest professorship at TU Dresden. AFAM was supported by a doctoral research grant from the German Academic Exchange Service (DAAD, Award No. 91653768).

References

- 1 R. Sharma and R. Gupta, Life Sci., 2007, 80, 2389-2392.
- 2 K. H. Kumar, S. Razack, I. Nallamuthu and F. Khanum, *Ind. Crops Prod.*, 2014, **52**, 815–826.
- 3 S. B. Badgujar and A. H. Bandivdekar, *J. Ethnopharmacol.*, 2015, **163**, 39–42.
- 4 A. Stierle, R. Upadhyay and G. Strobel, *Phytochemistry*, 1991, **30**, 2191–2192.
- 5 Y. Jaiswal, Z. Liang, P. Guo, H. M. Ho, H. Chen and Z. Zhao, J. Agric. Food Chem., 2014, 62, 7302–7316.
- 6 T. K. Lim, *Cyperus rotundus. Edible medicinal and non-medicinal plants*, Springer Science + Business Media Dordrecht, 2016, vol. 10, pp.178–208.
- 7 W. G. Seo, H. O. Pae, G. S. Oh, K. Y. Chai, T. O. Kwon, Y. G. Yun, N. Y. Kim and H. T. Chung, *J. Ethnopharmacol.*, 2001, 76, 59–64.
- 8 H. M. Sayed, M. H. Mohamed, S. F. Farag, G. A. Mohamed, O. R. M. Omobuwajo and P. Proksch, *Nat. Prod. Res.*, 2008, 22, 1487–1497.
- 9 M. Kumar, M. Rani and B. Meher, *Curr. Res. Pharm. Sci.*, 2017, **07**, 11–15.

Review

10 J. H. Jin, D. U. Lee, Y. S. Kim and H. P. Kim, *Arch. Pharm. Res.*, 2011, 34, 223–228.

- 11 Z. Zhou and W. Yin, Molecules, 2012, 17, 12636-12641.
- 12 B. Ryu, H. M. Kim, J. S. Lee, Y. J. Cho, M. S. Oh, J. H. Choi and D. S. Jang, *Helv. Chim. Acta*, 2015, **98**, 1372–1379.
- 13 H. B. Xu, Y. B. Ma, X. Y. Huang, C. A. Geng, H. Wang, Y. Zhao, T. H. Yang, X. L. Chen, C. Y. Yang, X. M. Zhang and J. J. Chen, *J. Ethnopharmacol.*, 2015, 171, 131–140.
- 14 N. A. Raut and N. J. Gaikwad, Fitoterapia, 2006, 77, 585-588.
- 15 S. J. Jeong, T. Miyamoto, M. Inagaki, Y. C. Kim and R. Higuchi, *J. Nat. Prod.*, 2000, **63**, 673–675.
- 16 J. H. Nam, D. Y. Nam and D. U. Lee, *J. Nat. Prod.*, 2016, **79**, 1091–1096.
- 17 K. Tsoyi, H. J. Jang, Y. S. Lee, Y. M. Kim, H. J. Kim, H. G. Seo, J. H. Lee, J. H. Kwak, D. U. Lee and K. C. Chang, *J. Ethnopharmacol.*, 2011, 137, 1311–1317.
- 18 T. Ito, H. Endo, H. Shinohara, M. Oyama, Y. Akao and M. Iinuma, *Fitoterapia*, 2012, **83**, 1420–1429.
- 19 L. Boulos and M. N. El-Hadidi, *The weed flora of Egypt*, The American University in Cairo Press, Cairo, 1984, p. 58.
- 20 N. Singh, V. K. Kulshrestha, M. B. Gupta and K. P. Bhargava, Indian J. Med. Res., 1970, 58, 103–109.
- 21 M. B. Gupta, T. K. Palit, N. Singh and K. P. Bhargava, *Indian J. Med. Res.*, 1971, **59**, 76–82.
- 22 K. Komai and K. Ueki, Weed Res. Japan, 1980, 25, 42-47.
- 23 K. Komai, C. S. Tang and R. K. Nishimoto, *J. Chem. Ecol.*, 1991, 17, 1–8.
- 24 P. Kalsi, A. Sharma, A. Singh, I. P. Singh and B. R. Chhabra, *Fitoterapia*, 1995, **66**, 94.
- 25 M. Zhu, H. H. Luk, H. S. Fung and C. T. Luk, *Phytother. Res.*, 1997, 11, 392–395.
- 26 S. C. Umeric and H. O. Ezeuzo, *Bioresour. Technol.*, 2000, 72, 193–196.
- 27 J. H. Ha, K. Y. Lee, H. C. Choi, J. Cho, B. S. Kang, J. C. Lim and D. U. Lee, *Biol. Pharm. Bull.*, 2002, **25**, 128–130.
- 28 M. Zhu, H. H. Luk, H. S. Fung and C. T. Luk, *Phytother. Res.*, 1998, 11, 392–394.
- 29 S. J. Uddin, K. Mondal, J. A. Shilpi and M. T. Rahman, *Fitoterapia*, 2006, 77, 134–136.
- 30 D. K. Pal and S. Dutta, *Indian J. Pahrm. Sci.*, 2006, **68**, 256–258.
- 31 A. Puratchikody, C. N. Devi and G. Nagalakshmi, *Indian J. Pharm. Sci.*, 2006, **68**, 97–101.
- 32 B. Lemaure, A. Touché, I. Zbinden, J. Moulin, D. Courtois, K. Macé and C. Darimont, *Phytother. Res.*, 2007, 21, 724–730.
- 33 H. M. Sayed, M. H. Mohamed, S. F. Farag, G. A. Mohamed and P. Proksch, *Nat. Prod. Res.*, 2007, **21**, 343–350.
- 34 A. Ardestani and R. Yazdanparast, *Int. J. Biol. Macromol.*, 2007, **41**, 572–578.
- 35 Z. A. Nima, M. S. Jabier, R. I. Wagi and H. A. Hussain, *Eng. Technol.*, 2008, **26**, 1156–1159.
- 36 V. Kempraj and S. K. Bhat, *Nat. Prod. Radiance*, 2008, 7, 416–419.
- 37 M. S. Sundaram, T. Sivakumar and G. Balamurugan, *Biomedicine*, 2008, **28**, 302–304.

- 38 S. I. Shivakumar, H. M. Suresh, C. S. Hallikeri, B. C. Hatapakki, J. S. Handiganur, K. Sankh and B. Shivakumar, J. Nat. Remed., 2009, 9, 92–196.
- 39 F. Mojab, H. Vahidi, B. Nickavar and M. Kamali-Nejad, *J. Med. Plant*, 2009, **8**, 91–97.
- Kilani-Jaziri, A. Neffati, I. Limem, J. Boubaker,
 I. Skandrani, M. B. Sghair, I. Bouhlel, W. Bhouri,
 A. M. Mariotte, K. Ghedira, M. G. Dijoux-Franca and
 L. Chekir-Ghedira, Chem. Biol. Interact., 2009, 181, 85–94.
- 41 D. Pal, S. Dutta and A. Sarkar, *Acta Pol. Pharm.*, 2009, **66**, 535–541.
- 42 M. E. Güldür, A. Ozgónül, I. H. Kilic, O. Sögüt, M. Ozaslan, M. Bitiren, M. Yalcin and D. Musa, *Int. J. Pharmacol.*, 2010, 6, 104–110.
- 43 A. Bisht, G. R. S. Bisht, M. Singh, R. Gupta and V. Singh, *Int. J. Res. Pharm. Biomed. Sci.*, 2011, 2, 661–665.
- 44 A. Singh, S. Maurya, R. Singh and U. P. Singh, *Arch. Phytopathol. Plant Prot.*, 2011, 44, 2004–2011.
- 45 A. G. Sunil, K. S. Kesavanarayanan, P. Kalaivani, S. Sathiya, V. Ranju, R. J. Priya, B. Pramila, F. D. S. Paul, J. Venkhatesh and C. S. Babu, *Brain Res. Bull.*, 2011, **84**, 394–405.
- 46 P. G. Daswani, S. Brijesh, P. Tetali and T. J. Birdi, *Indian J. Pharmacol.*, 2011, 43, 340–344.
- 47 G. K. Dang, R. R. Parekar, S. K. Kamat, A. M. Scindia and N. N. Rege, *Phytother. Res.*, 2011, 25, 904–908.
- 48 S. M. Khan and D. U. Lee, Nat. Prod. Sci., 2011, 17, 250-255.
- 49 M. Khalili, Z. Kiasalari, M. Roghani and Y. Azizi, *J. Med. Plant. Res.*, 2011, 5, 1140–1146.
- 50 N. Singh, B. R. Pandey, P. Verma, M. Bhalla and M. Gilca, *Indian J. Nat. Prod. Res.*, 2012, 3, 467–476.
- 51 D. Jebasingh, S. Venkataraman, D. D. Jackson and B. S. Emerald, *Int. J. Appl. Res. Nat. Prod.*, 2012, 5, 1–8.
- 52 G. F. A. El-Kaream, *J. Intercult. Ethnopharmacol.*, 2012, 1, 111–118.
- 53 R. S. Chandratre, S. Chandarana and S. A. Mengi, *Int. J. Pharm. Biol. Arch.*, 2012, 3, 598–600.
- 54 A. Chithran, B. T. Ramesh and N. Himaja, *Int. J. Phytopharmacol.*, 2012, 3, 130–134.
- 55 K. H. Kumar and F. Khanum, *Cell Mol. Neurobiol.*, 2013, 33, 5–17.
- 56 H. Weenen, M. H. H. Nkunya, D. H. Bray, L. B. Mwasumbi, L. S. Kinabo, V. A. E. B. Kilimali and J. B. P. A. Wijnberg, *Planta Med.*, 1990, 56, 371.
- 57 M. Ahmad, M. Mahayrookh, B. R. Asif and N. Jahan, *Pak. J. Pharmacol.*, 2012, **29**, 7–13.
- 58 K. H. Kumar and F. Khanum, *Cell Mol. Neurobio.*, 2013, **33**, 5–17.
- 59 S. Krishna and S. Renu, *J. Drug Del. Ther.*, 2013, 3, 109–113.
- 60 K. H. Kumar, A. Tamatam, A. Pal and F. Khanum, *Neurotoxicol*, 2013, **34**, 150–159.
- 61 Z. Rabiei, M. Hojjati, M. Rafieian-Kopaeia and Z. Alibabaei, *Biomed. Aging Pathol.*, 2013, 3, 185–191.
- 62 H. Imam, Z. Riaz and G. Sofi G, *Int. J. Green Pharm.*, 2013, 7, 37–40.
- 63 G. F. Mohammed, Aesthet. Surg. J., 2014, 34, 298-305.

64 S. E. Park, W. T. Shin, C. Park, S. H. Hong, G. Y. Kim, S. O. Kim, C. H. Ryu, S. H. Hong and Y. H. Choi, *Oncol. Rep.*, 2014, 32, 2461–2470.

RSC Advances

- 65 D. Jebasingh, D. D. Jackson, S. Venkataraman, E. Adeghate and B. S. Emerald, *Pharm. Biol.*, 2014, **15**, 1–12.
- 66 M. Z. Imam and C. D. Sumi, BMC Complement. Altern. Med., 2014, 14, 83.
- 67 H. Imam, S. Zarnigar, A. Ghulamuddin and L. A. Seikh, *Int. J. Nutr. Pharmacol. Nuerol. Dis.*, 2014, 4, 23–27.
- 68 H. H. T. Tran, M. C. Nguyen, H. T. Le, T. L. Nguyen, T. B. Pham and V. M. Chau, *J. Pharm. Biol.*, 2014, 52, 74–77.
- 69 K. Athesh, M. Divakar and P. Brindha, *Asian J. Pharm. Clin. Res.*, 2014, 7, 88–92.
- 70 A. M. Peerzada, H. H. Ali, M. Naeem, M. Latif, A. H. Bukhari and A. Tanveer, *J. Ethnopharmacol.*, 2015, 174, 540–560.
- 71 S. Kasala, K. Ramanjaneyulu, J. Himabindhu, R. Alluri and R. R. Babu, *J. Pharmacogn. Phytochem.*, 2016, 5, 407–409.
- 72 S. Suryavanshi, A. Choudhari, P. Raina and R. Kaul-Ghanekar, J. Ethnopharmacol., 2019, 242, 10.
- 73 F. Wang, X. Song, S. Ma, C. Liu, X. Sun, X. Wang, Z. Liu, D. Liang and Z. Yu, *Biosci. Rep.*, 2019, 39, BSR20190502.
- 74 S. Ma, F. Wang, C. Zhang, X. Wang, X. Wang and Z. Yu, *BMC Complementary Med. Therapies*, 2020, 20, 262.
- 75 Z. Shakerin, E. Esfandiari, S. Razavi, H. Alaei, M. Ghanadian and G. Dashti, *Adv. Biomed. Res.*, 2020, **9**, 17.
- 76 R. S. Dhillon, S. Singh, S. Kundra and A. S. Basra, *Plant Growth Regul.*, 1993, **13**, 89–93.
- 77 E. J. Seo, D. J. Lee, J. H. Kwak, S. M. Lee, Y. S. Kim and Y. S. Jung, *J. Ethnopharmacol.*, 2011, **135**, 48–54.
- 78 G. A. Mohamed, Bull. Fac. Pharm. Cairo Univ., 2015, 53, 5-9.
- 79 V. H. Kapadia, V. G. Naik, M. S. Wadia and S. Dev, *Tetrahedron Lett.*, 1967, 47, 4661-4667.
- 80 K. Komai and C. S. Tang, *Phytochemistry*, 1989, 28, 1883–1886.
- 81 C. Thebtaranonth, Y. Thebtaranonth, S. Wanauppathamkul and Y. Yuthavong, *Phytochemistry*, 1995, **40**, 125–128.
- 82 S. Ohira, T. Hasegawa, K. I. Hayashi, T. Hoshino, D. Takaoka and H. Nozaki, *Phytochemistry*, 1998, 47, 1577–1581.
- 83 M. M. Sonwa and W. A. König, *Phytochemistry*, 2001, **58**, 799–810.
- 84 Y. Xu, H. W. Zhang, C. Y. Yu, Y. Lu, Y. Chang and Z. M. Zou, *Molecules*, 2008, **13**, 2474–2481.
- 85 Y. Xu, H. W. Zhang, X. C. Wan and Z. M. Zoua, *Magn. Reson. Chem.*, 2009, 47, 527–531.
- 86 P. Alam, M. Ali and V. Aeri, J. Nat. Prod. Plant Resour., 2012, 2, 272–280.
- 87 M. P. Rani and K. P. Padmakumari, *J. Chromatogr. B*, 2012, **904**, 22–28.
- 88 G. F. Mohammed, Aesthet. Surg. J., 2014, 34, 298-305.
- 89 J. L. Yang and Y. P. Shi, *Planta Med.*, 2012, 78, 59-64.
- 90 Z. L. Zhou, S. Q. Lin and W. Q. Yin, *J. Asian Nat. Prod. Res.*, 2016, **18**, 662–668.
- 91 S. Sultana, M. Ali and S. R. Mir, *Open Plant Sci. J.*, 2017, **10**, 82–91
- 92 K. S. Chang, J. H. Jeon, G. H. Kim, G. W. Jang, S. J. Jeong, Y. R. Ju and Y. J. Ahn, *Sci. Rep.*, 2017, 7, 16643.

- 93 S. R. M. Ibrahim, G. A. Mohamed, K. Z. Alshali, R. A. Al Haidari, A. A. El-Kholy and M. F. Zayed, *Braz. J. Pharmacog.*, 2018, **28**, 320–324.
- 94 Y. J. Park, H. Zheng, J. H. Kwak and K. H. Chung, *Biomed. Pharmacother.*, 2019, **109**, 1313–1318.
- 95 M. A. Gamal, K. M. K. Hani and I. R. M. Sabrin, *Int. J. Pharmacogn. Phytochem. Res.*, 2015, 7, 83-99.
- 96 A. Kamala, S. K. Middha and C. S. Karigar, 3 Biotech, 2018, 8, 309.
- 97 L. Jirovetz, A. Wobus, G. Buchbauer, M. P. Shafi and P. T. Thampi, *Plant*, 2004, 7, 100–106.
- 98 S. Kilani, A. Abdelwahed, I. Chraief, R. B. Ammar, N. Hayder, M. Hammami, K. Ghedira and L. Chekir-Ghedira, *J. Essent. Oil*, 2005, 17, 695–700.
- 99 S. Kilani, I. Bouhlel, R. Ben, A. Abdelwahed, N. Hayder, A. Mahmoud, K. Ghedira and L. Chekir-Ghedira, *Toxicol. Environ. Chem.*, 2005, 87, 415–425.
- 100 R. Yazdanparast and A. Ardestani, J. Med. Food, 2007, 10, 667–674.
- 101 S. Kilani, S. M. Ben, I. Limem, I. Bouhlel, J. Boubaker, W. Bhouri, I. Skandrani, A. Neffatti, R. Ben-Ammar, M. G. Dijoux-Franca, K. Ghedira and L. Chekir-Ghedira, *Bioresour. Technol.*, 2008, 99, 9004–9008.
- 102 S. Kilani, J. Ledauphin, I. Bouhlel, M. Ben-Sghaier, J. Boubaker, I. Skandrani, R. Mosrati, K. Ghedira, D. Barillier and L. Chekir-Ghedira, *Chem. Biodivers.*, 2008, 5, 729–742.
- 103 O. A. Lawal and A. O. Oyedeji, *Molecules*, 2009, **14**, 2909–2917.
- 104 S. Kilani-Jaziri, W. Bhouri, I. Skandrani, I. Limem, L. Chekir-Ghedira and K. Ghedira, *S. Afr. J. Bot.*, 2011, 77, 767–776.
- S. Kilani-Jaziri, D. Mhalla, F. Châbane, Z. Ghedira,
 I. Limem, K. Ghedira and L. Chekir-Ghedira, BMC Complement. Alternat. Med., 2013, 13, 28.
- 106 A. Sharma, R. Verma and P. Ramteke, J. Herbal Med., 2014, 4, 74–82.
- 107 P. Mannarreddy, M. Denis, D. Munireddy, R. Pandurangan, K. P. Thangavelu and K. Venkatesan, *Biomed. Pharmacother.*, 2017, 95, 1375–1387.
- 108 Q. P. Hu, X. M. Cao, D. L. Hao and L. L. Zhang, *Sci. Rep.*, 2017, 7, 45231.
- 109 R. A. Abo-Altemen, A. M. Al-Shammari and M. S. Shawkat, *Energy Procedia*, 2019, **157**, 1462–1474.
- 110 F. G. Rocha, M. D. M. Brandenburg, P. L. Pawloski, B. D. S. Soley, S. C. A. Costa, C. C. Meinerz, I. P. Baretta, M. F. Otuki and D. A. Cabrini, *J. Ethnopharmacol.*, 2020, 254, 112709.
- 111 B. Huang, J. Liu, S. Fu, Y. Zhang, Y. Li, D. He, X. Ran, X. Yan, J. Du, T. Meng, X. Gao and D. Liu, *Front. Pharmacol.*, 2020, 11, 281.
- 112 K. Komai and K. Ueki K, *J. Weed Sci. Technol.*, 1975, **20**, 66–71.
- 113 K. Komai, J. Iwamura and K. Ueki, *J. Weed Sci. Technol.*, 1977, 22, 14–18.
- 114 P. N. Singh and S. B. Singh, Phytochemistry, 1980, 19, 2056.

Review

115 S. K. Kim, B. Y. Hwang, S. J. Kang, J. J. Lee, J. S. Ro and K. Lee, *Koran I. Pharmacogn.*, 2000, **31**, 1–6.

- 116 S. J. Kim, H. J. Kim, Y. P. Jang, M. S. Oh and D. S. Jang, *Bull. Korean Chem.*, 2012, 33, 3115–3118.
- 117 H. G. Kim, J. Hong, Y. Huh, C. Park, D. S. Hwang, J. H. Choi and M. S. Oh, *J. Ethnopharmacol.*, 2013, **148**, 322–328.
- 118 S. J. Kim, B. Ryu, H. Y. Kim, Y. I. Yang, J. Ham, J. H. Choi and D. S. Jang, *Bull. Korean Chem. Soc.*, 2013, 34, 2207– 2210.
- 119 Y. X. Li, J. Chem. Pharm. Res., 2014, 6, 1496-1500.
- 120 S. K. Paknikar, O. Motl and K. K. Chakravarti, *Tetrahedron Lett.*, 1977, 24, 2121–2124.
- 121 S. H. Jung, S. J. Kim, B. G. Jun, K. T. Lee, S. P. Hong, M. S. Oha, D. S. Jang and J. H. Choi, *J. Ethnopharmacol.*, 2013, 147, 208–214.
- 122 A. Dadang, K. Ohsawa, S. Kato and I. Yamamoto, *J. Pesticide Sci.*, 1996, **21**, 444–446.
- 123 Y. J. Seo, M. Jeong, K. T. Lee, D. S. Jang and J. H. Choi, *Int. Immunopharmacol.*, 2016, 38, 61–69.
- 124 V. H. Kapadia, B. A. Nagasampagi, V. G. Naik and S. Dev, *Tetrahedron*, 1965, **21**, 607–618.
- 125 H. Hikino, K. Aota and T. Takemoto, *Tetrahedron*, 1967, 23, 2169–2172.
- 126 H. Hikino, K. Aota, D. Kuwano and T. Takemoto, *Tetrahedron*, 1971, 27, 4831–4836.
- 127 H. Hikino and K. Aota, Phytochemistry, 1976, 16, 1265-1266.
- 128 M. Morimoto and K. Komai, *Weed Biol. Manage.*, 2005, 5, 203–209.
- 129 S. Rune, F. Torgils and M. V. Ingunn, *J. Agric. Food Chem.*, 2007, 55, 10067.
- 130 P. R. Jensen, K. M. Jenkins, D. Porter and W. Fenical, Appl. Environ. Microbiol., 1998, 64, 1490.

- 131 C. N. Liu, S. H. Kuo and M. C. Chung, *J. Nat. Prod.*, 1997, **60**, 851.
- 132 H. W. Chu, H. T. Wu and Y. J. Lee, *Tetrahedron*, 2004, **60**, 2647.
- 133 S. Li, C. Y. Lo and C. T. Ho, J. Agric. Food Chem., 2006, 54, 4176.
- 134 J. H. Yoon, T. G. Lim, K. M. Lee, A. J. Jeon, S. Y. Kim and K. W. Lee, *J. Agric. Food Chem.*, 2011, **59**, 222.
- 135 S. E. Nielsen, V. Breinholt, C. Cornett and L. O. Dragsted, *Food Chem. Toxicol.*, 2000, **38**, 739.
- 136 Z. Cheng, S. Surichan, K. Ruparelia, R. Arroo and M. R. Boarder, *Br. J. Pharmacol.*, 2011, **162**, 1781.
- 137 S. Li, M. H. Pan, C. S. Lai, C. Y. Lo, S. Dushenkov and C. T. Ho, *Bioorg. Med. Chem.*, 2007, 15, 3381.
- 138 G. Casano, A. Dumetre, C. Pannecouque, S. Hutter, N. Azas and M. Robin, *Bioorg. Med. Chem.*, 2010, **18**, 6012.
- 139 N. Beldjoudi, L. Mambu, M. Labbaied, P. Grellier, R. David, R. Philippe, T. M. Mare and F. Frappier, *J. Nat. Prod.*, 2003, **66**, 1447.
- 140 D. Paola, B. Ivana, R. Carla, R. Marina, D. Paola, S. Luca and M. Luigi, *Flavour Fragrance J*, 2010, **26**, 34.
- 141 J. B. Harborne, C. A. Williams and K. L. Wilson, *Phytochemistry*, 1982, 21, 2491–2507.
- 142 K. Arraki, P. Totoson, A. Decendit, A. Badoc, A. Zedet, J. Jolibois, M. Pudlo, C. Demougeot and C. Girard-Thernier, J. Nat. Prod., 2017, 80, 2432–2438.
- 143 P. Liu, L. Liu, Y. P. Tang, J. A. Duan and N. Y. Yang, *Chin. Chem. Lett.*, 2010, 21, 606–609.
- 144 H. A. Quayyum, A. U. Mallik, D. M. Leach and C. Gottardo, *J. Chem. Ecol.*, 2000, **26**, 2221–2231.