


 Cite this: *RSC Adv.*, 2021, **11**, 28530

 Received 14th July 2021
 Accepted 16th August 2021

 DOI: 10.1039/d1ra05405h
rsc.li/rsc-advances

Selective oxidation of alcohol- d_1 to aldehyde- d_1 using MnO_2 †

Hironori Okamura, Yoko Yasuno, Atsushi Nakayama, Katsushi Kumadaki, Kohei Kitsuwa, Keita Ozawa, Yusaku Tamura, Yuki Yamamoto and Tetsuro Shinada *

The selective oxidation of alcohol- d_1 to prepare aldehyde- d_1 was newly developed by means of $NaBD_4$ reduction/activated MnO_2 oxidation. Various aldehyde- d_1 derivatives including aromatic and unsaturated aldehyde- d_1 can be prepared with a high deuterium incorporation ratio (up to 98% D). Halogens (chloride, bromide, and iodide), alkene, alkyne, ester, nitro, and cyano groups in the substrates are tolerated under the mild conditions.

1. Introduction


Deuterium (2H , d) is a stable, non-radioactive, and safe isotope of hydrogen (1H). Since its discovery,¹ d has been widely utilized in organic chemistry, biochemistry, analytical chemistry, pharmaceutical science, and drug discovery.^{2,3} Because of the high demand for d-labelled molecules in the scientific research fields, many efforts have been devoted to developing a new method for the synthesis of d-labelled molecules.

Aldehyde- d_1 2 has received significant attention as a synthetic target due to the facts that aldehyde 1 is a useful feedstock in organic synthesis. Various methods have been performed in the synthesis of alkyl and aryl aldehyde- d_1 . For example, more than 40 syntheses (25 different reaction conditions) of benzaldehyde- d_1 (PhCDO) were conducted even since 2018 in the studies to develop new d-incorporation method or reaction mechanism using PhCDO.⁴⁻⁸

The previous synthetic approaches to access d-labelled molecules are classified into 5 types; (A) addition of D^- followed by oxidation, (B) carbonyl Umpolung approach, (C) radical reaction, (D) transition metal-catalysed reaction, and (E) others. Recently, mild, one-step, and catalytic syntheses of aldehyde- d_1 2 have been achieved by deuteration of the Breslow intermediates,⁹ deuteration of acyl radicals,^{6,10} and transition metal-catalysed deuterium incorporation.¹¹ However, the previous synthetic methods including the modern direct syntheses often suffered from drawbacks such as over deuteration, requirements of harsh conditions (high and low temperature, and strong base and acids), and the use of

expensive catalysts. Moreover, the synthetic examples of substituted acrolein and propynal- d_1 are much less than those of alkyl and aryl aldehyde- d_1 ,^{12,13} though recently developed NHC-catalysed H-D exchange reactions allowed access to various substituted acrolein- d_1 derivatives.⁹ In this context, development of a new d-incorporation method which allows flexible synthesis of aromatic and unsaturated aldehyde- d_1 2 remains to be a challenging synthetic task (Scheme 1).

Method A using D^- as a deuterium source has been recognized as a robust and conventional synthetic method to prepare aldehyde- d_1 2 (Scheme 2). The synthesis is typically performed





Scheme 1 Synthesis of aldehyde- d_1 derivatives. (A) reduction of the formyl with D^- and oxidation, (B) carbonyl Umpolung approach, (C) radical H-D exchange, (D) transition metal-catalysed H-D exchange, and (E) others.

Graduate School of Science, Osaka City University, Sugimoto, Sumiyoshi, Osaka, 558-8585 Japan. E-mail: shinada@sci.osaka-cu.ac.jp

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d1ra05405h

‡ HO and Y. Yasuno contributed equally to this study.



Previous work: Oxidation of alcohol-*d*₂This work: Selective oxidation of alcohol-*d*₁

28 examples, Mild conditions,
Short reaction period, Simple operation,
Compatibility with ester, nitro, nitrile, alkyne, olefin


Scheme 2 Oxidation of deuterated alcohols. Eqn (1): oxidation of alcohol-*d*₂, eqn (2): selective oxidation of alcohol-*d*₁.

in two steps; (i) reduction of carboxylic acid derivatives using LiAlD_4 to provide alcohol-*d*₂ 3 and (ii) oxidation to aldehyde-*d*₁ 2 (eqn (1)).¹⁴ In this approach, the deuterium incorporation ratio (%D) of the commercially available D^- sources such as LiAlD_4 (~ 98 atom) is reliably transferred into the product. On the other hand, the use of highly reactive LiAlD_4 often limits the synthetic scope. Under the conditions, various functional groups such as nitro, nitrile, ester, and acid moieties, and alkene and alkyne with electron-withdrawing group(s) are not tolerated. To overcome the limitation, we emerged selective oxidation of alcohol-*d*₁ derivatives 4 (Scheme 2(eqn (2))). It is expected that various alcohol-*d*₁ 4 can be prepared by the mild NaBD_4 reduction. The next selective oxidation of D (H/D selectivity) is the key to this approach. Recently, oxidation of benzyl alcohol-*d*₁ (PhCDHOH) with PCC or PDC was conducted to prepare PhCDO with $\sim 85\%$ D.^{4a,4e,4q} On the other hand, further efforts to improve the selectivity (%D) in the selective oxidation have not been well-examined. Herein, we would like to report that NaBD_4 reduction followed by activated MnO_2 oxidation ($\text{NaBD}_4/\text{MnO}_2$ system). The simple and mild protocol allows expansion of the synthetic range of aldehyde-*d*₁ 2 including not only aromatic aldehyde-*d*₁ derivatives but also substituted acrolein-*d*₁ and propynal-*d*₁ derivatives with high %D (up to 98%).

2. Results and discussion

In a similar manner to the previous synthetic examples of NaBH_4 reduction of aldehyde 1, the reduction with NaBD_4 gave the corresponding alcohol-*d*₁ derivatives 4 with excellent functional group compatibility and yields (Scheme 3). Chloride, bromide, iodide, methoxy, ethoxy, or methylene acetal, nitrile, ester, nitro, and alkyne groups on the aromatic ring of 4c–4q were tolerated under the conditions. Substituted acrolein and propynal 1r–1aa also underwent smooth NaBD_4 reduction to provide 4r–4aa without loss of the alkyne and alkene moieties, and tetrahydropyranyl (THP), benzoyl (Bz), and *tert*-butyldimethylsilyl (TBS) protecting groups.

We next examined the key oxidation of alcohol-*d*₁ 4 using 4-phenylbenzyl alcohol-*d*₁ (4a) (Table 1). As a result, activated MnO_2 was found to be superior to other general oxidation



Scheme 3 Reduction of aldehyde 1 with NaBD_4 .

reagents (entry 1, Table 1). Treatment of 4a with 23 eq. of MnO_2 in CH_2Cl_2 gave aldehyde-*d*₁ 2a with 92% D in 2 h. The use of pyridinium dichlorochromate (PDC) gave 2a in good selectivity (88% D).^{4e} However, the isolate yield was moderate (entry 2). Dess–Martin periodinane oxidation, 2,2,6,6-tetramethylpiperidine 1-oxyl (TEMPO) oxidation in the presence of PhI(OAc)_2 , and Parikh–Doering oxidation (sulfur trioxide–pyridine complex in dimethyl sulfoxide (DMSO)) resulted in lower selectivities (74, 76 and 66%D, entries 3–5).

Activated MnO_2 oxidation was successfully expanded to the synthesis of various aldehyde-*d*₁ 2a–2aa with high %D (85–96% D) (Scheme 4A–C). Chloride, bromide, iodide, methoxy, ethoxy, or methylene acetal, nitrile, ester, nitro, and alkyne groups on the aromatic ring of 4c–4q are preserved under the mild oxidation conditions (Scheme 4A). Substituted acrolein 4r–4v and propynal 4w–4aa smoothly underwent MnO_2 oxidation to provide 2r–2aa without loss of the alkyne and alkene moieties (Scheme 4B and C). The synthetic utility was further demonstrated by the synthesis of 2v with a bromo group at the α -position of cinnamaldehyde. In addition, Bz, THP, and TBS protecting groups of 4y, 4z, and 4aa were also maintained under the conditions. These propargyl alcohols 4y, 4z, and 4aa were smoothly converted to the corresponding propynal derivatives 2y, 2z, and 2aa with high %D, respectively.

In conjunction with our recent efforts toward elucidation of biosynthetic reaction mechanisms of terpene synthases using d-labelled prenols,^{15,16} we needed geranylgeraniol-*d*₂ (6) as an enzyme substrate. Previously, the synthesis of 6 (ref. 17) and other acyclic prenol-*d*₂ derivatives¹⁸ was performed in four steps from 5 via reduction of ester 7 with LiAlD_4 . However, commercially available LiAlD_4 is almost out of stock in recently years. In addition, low temperature conditions (-20°C) is required for the LiAlD_4 reduction to avoid the undesired 1,4-reduction. We




Table 1 Oxidation of alcohol-*d*₁ 4a^a

| Entry | Conditions | Yield ^b (%) | %D ^c |
|-------|---|------------------------|-----------------|
| 1 | MnO ₂ (23 eq.), 1 h | 92 | 92 |
| 2 | PDC (1.2 eq.), MS4A, 2 h | 51 | 88 |
| 3 | Dess–Martin periodinane (1.5 eq.), 5 min | 84 | 74 |
| 4 | TEMPO (0.01 eq.), Bu ₄ NHSO ₄ (0.05 eq.), NaOCl (1.2 eq.), 1 h | 96 | 76 |
| 5 | DMSO (10 eq.), SO ₃ -pyridine (4 eq.), <i>i</i> Pr ₂ NET (5 eq.), 1.5 h | 75 | 66 |

^a 0.5 mmol scale. ^b Isolated yield. ^c %D for 2a is calculated based on the integration ratios of aldehyde and aromatic proton. MnO₂ = activated manganese dioxide, PDC = pyridinium dichlorochromate, MS4A = molecular sieves 4A, TEMPO = 2,2,6,6-tetramethylpiperidine 1-oxyl, DMSO = dimethyl sulfoxide.


expected that NaBD₄/MnO₂ system would be an alternative to the LiAlD₄ procedure to prepare **6**, conveniently. According to the literature,¹⁹ geranylgeranol (**5**) was converted to aldehyde **8** by MnO₂ oxidation (Scheme 5). Aldehyde **8** was subjected to NaBD₄/MnO₂ to deliver d-enriched aldehyde **9** which was

subsequently reduced by NaBD₄ to provide geranylgeranol-*d*₂ (**6**) in 70% yield over four steps with satisfactory deuterium incorporation ratio (94% D). Under the conditions, the undesired 1,4-addition reaction was not observed. Thus, an



Scheme 4 MnO₂ oxidation of alcohol-*d*₁ 4. (A) synthetic examples of aromatic aldehyde-*d*₁ 2, (B) synthetic examples of substituted acrolein-*d*₁ 2, and (C) synthetic examples of substituted propynal-*d*₁ 2. ^a2 h, ^b6 h, ^c12 h.





Scheme 5 Synthesis of geranylgeraniol-*d*₂ (6) from geranylgeraniol (5).

operationally simple and mild deuteration of prenols-*d*₂ was achieved by application of NaBD₄/MnO₂ system.

Previously, Brecker *et al.* investigated ¹³C kinetic isotope effects (KIEs) in the oxidation of cinnamyl alcohol using MnO₂, Dess–Martin periodinane, and Swern oxidation (DMSO/(COCl)₂/Et₃N) to gain insight into the reaction mechanism.²⁰ Comparison of the kinetic isotope of effects revealed the following order MnO₂ > Dess–Martin oxidation ≈ Swern oxidation. The higher ¹³C KIE using MnO₂ displayed that the C–H bond breaking in the intermediate is irreversible and rate-determining, and the oxidation proceeded *via* energy rich transition state. On the other hand, the lower ¹³C KIEs observed in Swern oxidation and Dess–Martin oxidation indicated that the intramolecular C–H bond cleavage in these oxidation reaction processes would not be slower to be rate-limiting.

Experimental results in Table 1 clearly shows that the degrees of %D are as follows MnO₂ > PDC > TEMPO ≈ Dess–Martin > SO₃–pyridine/DMSO. It is speculated that higher %D of MnO₂ oxidation and lower %D of SO₃–pyridine/DMSO oxidation would correlate to the ¹³C KIE data (MnO₂ > Swern oxidation). It is interesting to note that the %D value in Scheme 4 depended on the substrates. The oxidation of propargyl alcohols **4w–4aa** resulted in higher %D than those of the other alcohols. The oxidation of **4w–4aa** needed a longer reaction period to complete the reactions. As mentioned in the previous ¹³C KIE studies, the rate limiting steps of the MnO₂ oxidation relies on the C–H cleavage step of the reaction intermediate. It is considered that the slower C–H cleavage would provide the higher %D.

3. Conclusions

We have established a facile synthesis of aldehyde-*d*₁ derivatives by NaBD₄/MnO₂ system. The new method is characterized by a high degree of functional group compatibility and a wide range of substrate scope including the synthesis of d-containing unsaturated aldehydes. Aromatic aldehyde-*d*₁ derivatives such as **2c** and **2g** would be a useful synthetic intermediate for olefination, amination, hydride reduction, Suzuki cross coupling, and Sonogashira coupling reactions.^{9e,10c} Substituted acroleins and propynals can be used for Michael addition reaction,

cycloaddition reaction, and transition metal catalysed transformations. In this context, NaBD₄/MnO₂ system would offer vital opportunity to the synthesis of highly functionalized d-labelled molecules *via* facile preparation of aromatic and unsaturated aldehyde-*d*₁ **2**. Deuterium-labelled compounds are often needed for the investigation of the mechanisms or determination of the rate-limiting step. The present synthetic method supports the studies from the viewpoint of the facile preparation of aldehyde-*d*₁ **2** and its derivatives. Further application and mechanism studies are ongoing in our laboratory.

Author contributions

Y. Yasuno and HO are contributed equally. Y. Yasuno, HO, and TS designed the synthetic route. TS wrote the manuscript. HO, Y. Yasuno, and AN prepared ESI.† HO, Y. Yasuno, AN, K. Kumadaki, K. Kitsuwa, KO, YT, and Y. Yamamoto performed syntheses of **2**.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was financially supported from JSPS Kakenhi (JP19H04661 for TS, JP20K09396 for AN, and JP21K14629 for YYasuno), the OCU ‘Think globally, act locally’ Research Grant for Young Scientists 2021 through the hometown donation fund of Osaka City for AN, the Uehara Memorial Foundation for AN, and the Sasakawa Scientific Research Grant from The Japan Science Society for HO.

Notes and references

- H. C. Urey, F. G. Brickwedde and G. M. Murphy, *Phys. Rev.*, 1932, **39**, 164.
- J. Yang, in *Deuterium: Discovery and Applications in Organic Chemistry*, Elsevier, Amsterdam, 2016.
- (a) T. Pirali, M. Serafini, S. Cargnini and A. A. Genazzani, *J. Med. Chem.*, 2019, **62**, 5276; (b) J. Atzrodt, V. Derda, W. J. Kerr and M. Reid, *Angew. Chem., Int. Ed.*, 2018, **57**, 1758; (c) A. Mullard, *Nat. Rev. Drug Discovery*, 2016, **15**, 219; (d) A. Katsnelson, *Nat. Med.*, 2013, **19**, 656; (e) J. M. Herbert, *J. Labelled Compd. Radiopharm.*, 2010, **53**, 658.
- Synthesis of PhCDO (type A): (a) A. S. K. Raj, A. S. Narode and R.-S. Liu, *Org. Lett.*, 2021, **23**, 1378; (b) M. Zeng, C. Lou, J.-W. Xue, H. Jiang, L. Kaiwen, Z. Chen, S. Fu and G. Yin, *Appl. Organomet. Chem.*, 2021, **35**, e6093; (c) L. Kang, J. Zhang, H. Yang, J. Quan and G. Jinlong, *Org. Lett.*, 2020, **22**, 9118; (d) J.-Y. Gu, W. Zhang, S. R. Jackson, Y.-H. He and Z. Guan, *Chem. Commun.*, 2020, **56**, 13441; (e) J.-L. Xu, H. Tian, J.-H. Kang, W.-X. Kang, W. Sun, R. Sun, Y.-M. Li and M. Sun, *Org. Lett.*, 2020, **22**, 6739; (f) C. Lázaro-Milla, J. Macicior, H. Yanai and P. Almendros, *Chem.-Eur. J.*, 2020, **26**, 8983; (g) Y. Yang, X. Zhang, L.-P. Zhong, J. Lan, X. Li, C.-C. Li and L. W. Chung, *Nat. Commun.*, 2020, **11**,



1850; (h) Y. Zhang, W. Schilling, D. Riemer and S. Das, *Nat. Protoc.*, 2020, **15**, 822; (i) F. Hu, Z. Chen, Y. Tan, D. Xu, S. Huang, S. Jia, X. Gong, W. Qin and H. Yan, *Org. Lett.*, 2020, **22**, 1934; (j) J.-X. Wang, X.-T. Zhou, Q. Han, X.-X. Guo, X.-H. Liu, C. Xue and H.-B. Ji, *New J. Chem.*, 2019, **43**, 19415; (k) T. Kaicharla, B. M. Zimmermann, M. Oestreich and J. F. Teichert, *Chem. Commun.*, 2019, **55**, 13410; (l) S.-S. Meng, L.-R. Lin, X. Luo, H.-J. Lv, J.-L. Zhao and A. S. C. Chan, *Green Chem.*, 2019, **21**, 6187; (m) G.-F. Pan, X.-L. Zhang, X.-Q. Zhu, R.-L. Guo and Y.-Q. Wang, *iScience*, 2019, **20**, 229; (n) G. Nicolau, G. Tarantino and C. Hammond, *ChemSusChem*, 2019, **12**, 4953; (o) A. K. Bains, A. Kundu, S. Yadav and D. Adhikari, *ACS Catal.*, 2019, **9**, 9051; (p) J. Qian, J. Zhang, H. Yang, L. Kang and G. Jiang, *Chem. Sci.*, 2019, **10**, 8812; (q) R. D. Kardile and R.-S. Liu, *Org. Lett.*, 2019, **21**, 6452; (r) J. L. DiMeglio, A. G. Breuhaus-Alvarez, S. Li and B. M. Bartlett, *ACS Catal.*, 2019, **9**, 5732; (s) T. Yatabe, N. Mizuno and K. Yamaguchi, *ACS Catal.*, 2018, **8**, 11564; (t) D. Wang, W. Liu, Y. Hong and X. Tong, *Org. Lett.*, 2018, **20**, 5002; (u) M. Zhang, Y. Zhai, S. Ru, D. Zang, S. Han, H. Yu and Y. Wei, *Chem. Commun.*, 2018, **54**, 10164; (v) Z.-P. Wang, Y. He and P.-L. Shao, *Org. Biomol. Chem.*, 2018, **16**, 5422; (w) T. C. Malig, D. Yu and J. E. Hein, *J. Am. Chem. Soc.*, 2018, **140**, 9167; (x) S. Shirase, K. Shinohara, H. Tsurugi and K. Mashima, *ACS Catal.*, 2018, **8**, 6939; (y) W. Schilling, D. Riemer, Y. Zhang, N. Hatami and S. Das, *ACS Catal.*, 2018, **8**, 5425; (z) Z.-Z. Zhou, M. Liu, L. Lv and C.-J. Li, *Angew. Chem., Int. Ed.*, 2018, **57**, 2616.

5 Synthesis of PhCDO (type B): (a) D.-X. Zhu, H. Xia, J.-G. Liu, L. W. Chung and M.-H. Xu, *J. Am. Chem. Soc.*, 2021, **143**, 2608; (b) S. K. Mahesh, J. B. Nanubolu and G. Sudhakar, *J. Org. Chem.*, 2019, **84**, 7815.

6 Synthesis of PhCDO (type C): Y. Kuang, H. Cao, H. Tang, J. Chew, W. Chen, X. Shi and J. Wu, *Chem. Sci.*, 2020, **11**, 8912.

7 Synthesis of PhCDO (type D): (a) Z. Liu, P. Wang, Z. Yan, S. Chen, D. Yu, X. Zhao and T. Mu, *Beilstein J. Org. Chem.*, 2020, **16**, 645; (b) Z. Liu, Z. Yang, B. Yu, X. Yu, H. Zhang, Y. Zhao, P. Yang and Z. Liu, *Org. Lett.*, 2018, **20**, 5130.

8 Synthesis of PhCDO (type E): (a) A. Baykal and B. Plietker, *Eur. J. Org. Chem.*, 2020, 1145; (b) S. K. Mahesh, J. B. Nanubolu and G. Sudhakar, *J. Org. Chem.*, 2019, **84**, 7815; (c) N. A. Eberhardt, N. P. N. Wellala, Y. Li, J. A. Krause and H. Guan, *Organometallics*, 2019, **38**, 1468; (d) T. Li, B.-H. Xu, D.-P. Zhu, Y.-F. Wang and S.-J. Zhang, *Org. Chem. Front.*, 2018, **5**, 1933.

9 (a) V. G. Landge, K. K. Shrestha, A. J. Grant and M. C. Young, *Org. Lett.*, 2020, **22**, 9745; (b) S. C. Gadekar, V. Dhayalan, A. Nandi, I. L. Zak, S. Barkai, M. S. Mizrachi, S. Kozuch and A. Milo, *ChemRxiv*, 2020, 1; (c) A. Palazzolo, T. Naret, M. Daniel-Bertrand, D.-A. Buisson, S. Tricard, P. Lesot, Y. Coppel, B. Chaudret, S. Feuillastre and G. Pieters, *Angew. Chem., Int. Ed.*, 2020, **59**, 20879; (d) Y. Sawama, Y. Miki and H. Sajiki, *Synlett*, 2020, **31**, 699; (e) H. Geng, X. Chen, J. Gui, Y. Zhang, Z. Shen, P. Qian, J. Chen, S. Zhang and W. Wang, *Nat. Catal.*, 2019, **2**, 1071; (f) W. Liu, L.-L. Zhao, L.-M. Melaimi, L. Cao, X. Xu, J. Bouffard, G. Bertrand and X. Yan, *Chem.*, 2019, **5**, 2484.

10 (a) J.-Y. Dong, W.-T. Xu, F.-Y. Yue, H.-J. Song, Y.-X. Liu and Q.-M. Wang, *Tetrahedron*, 2021, **82**, 131946; (b) Y. Zhang, P. Ji, Y. Dong, Y. Wei and W. Wang, *ACS Catal.*, 2020, **10**, 2226; (c) J. Dong, X. Wang, Z. Wang, H. Song, Y. Liu and Q. Wang, *Chem. Sci.*, 2020, **11**, 1026.

11 (a) M. Y. S. Ibrahim and S. E. Denmark, *Angew. Chem., Int. Ed.*, 2018, **57**, 10362; (b) W. J. Kerr, R. M. Reid and T. Tuttle, *Angew. Chem., Int. Ed.*, 2017, **56**, 7808.

12 Synthesis of cinnamaldehyde-*d*₁: (a) B. D. Mokar and C. S. Yi, *Organometallics*, 2019, **38**, 4625; (b) X. Li, S. Wu, S. Chen, Z. Lai, H.-B. Luo and C. Sheng, *Org. Lett.*, 2018, **20**, 1712; (c) Y. Shi, B. Jung, S. Torker and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2015, **137**, 8948; (d) T. Fujihara, C. Cong, T. Iwai, J. Terao and Y. Tsuji, *Synlett*, 2012, **23**, 2389.

13 Synthesis of 3-phenylpropinal-*d*₁: (a) B. Zhou, Q. Wu, Z. Dong, J. Xu and Z. Yang, *Org. Lett.*, 2019, **21**, 3594; (b) A. Verlee, T. Heugebaert, T. van der Meer, P. Kerchev, F. Van Breusegem and C. V. Stevens, *Org. Biomol. Chem.*, 2018, **16**, 9359; (c) A. Vallet, A. Janin and R. Romanet, *J. Labelled Compd.*, 1971, **7**, 80.

14 Alternative practical synthesis of PhCDO from benzyl, see ref. 4*c*, 4*i*, 4*p*, and 4*w*.

15 (a) R. Stepanova, H. Inagi, K. Sugawara, K. Asada, T. Nishi, D. Ueda, Y. Yasuno, T. Shinada, K. Miki, M. Fujihashi and T. Sato, *ACS Chem. Biol.*, 2020, **15**, 1517; (b) Q. Chen, J. Li, Z. Liu, T. Mitsuhashi, Y. Zhang, H. Liu, Y. Ma, J. He, T. Shinada, T. Sato, Y. Wang, H. Liu, I. Abe, P. Zhang and G. Wang, *Plant Commun.*, 2020, 100051; (c) Y. Totsuka, S. Ueda, T. Kuzuyama and T. Shinada, *Bull. Chem. Soc. Jpn.*, 2015, **44**, 575; (d) A. Meguro, Y. Motoyoshi, K. Teramoto, S. Ueda, Y. Totsuka, Y. Ando, T. Tomita, S.-Y. Kim, T. Kimura, M. Igarashi, R. Sawa, T. Shinada, M. Nishiyama and T. Kuzuyama, *Angew. Chem., Int. Ed.*, 2015, **54**, 4353.

16 For a review using stable isotopes in the study of terpene biosynthesis: J. Rinkel and J. S. Dickschat, *Beilstein J. Org. Chem.*, 2015, **11**, 2493.

17 V. A. Clausen, R. L. Edelstein and M. D. Distefano, *Biochemistry*, 2001, **40**, 3920.

18 N. Duhamel, D. Martin, R. Larcher, B. Fedrizzi and D. Barker, *Tetrahedron Lett.*, 2016, **57**, 4496.

19 M. P. Doyle and M. Yan, *ARKIVOC*, 2002, (viii), 180.

20 L. Brecker, M. F. Kögl, C. E. Tyl, R. Kratzerb and B. Nidetzky, *Tetrahedron Lett.*, 2006, **47**, 4045.

