Chemical Science



EDGE ARTICLE

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2021, 12, 4126

d All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 27th December 2020 Accepted 1st February 2021

DOI: 10.1039/d0sc07042d

rsc.li/chemical-science

Palladium-catalyzed remote para-C-H activation of arenes assisted by a recyclable pyridine-based template†


Xiaoxi Chen,^{ab} Shuai Fan,^a Meng Zhang,^a Yuzhen Gao, ^{ba} Shangda Li^a and Gang Li^{b*ab}

Direct para-selective C-H functionalization of arenes remains a daunting challenge and is still significantly restricted to a few scaffolds. Herein, we report an unprecedented pyridine-based para-directing template (DT) assisted, Pd-catalyzed para-C-H alkenylation of three classes of arenes, i.e. phenylpropanoic acids, 2-phenyl benzoic acids and benzyl alcohols, with a series of alkenes including perfluoroalkenes. Notably, the pyridine-based para-DT could be easily synthesized and readily recycled under mild conditions. These results may find application in rapid construction of para-substituted arenes and stimulate the exploration of novel methods for para-C-H functionalization of arenes.

Introduction

Transition-metal-catalyzed C-H activation assisted by directing groups (DGs) has proven to be a powerful strategy for achieving regioselective C-H functionalizations in a general manner, which is able to differentiate C-H bonds with similar electronic and steric properties in an organic molecule. The DG strategy has enabled the development of a large number of ortho-C-H and some meta-C-H transformations of arenes. 1,2 In contrast, although quite a few para-C-H transformations of arenes have been disclosed by making use of arene's electronic and steric biases,3,4 the para-C-H transformation beyond the control of such biases remains a daunting challenge and is still significantly restricted to a few scaffolds. In this endeavor, Nakao and co-workers reported a breakthrough in para-C-H alkylation and borylation of arenes by cooperative nickel/aluminum and iridium/aluminum catalysis (Scheme 1a, left).5 Chattopadhyay,6 Phipps^{7a} and Maleczka^{7b} also independently realized elegant Ircatalyzed, noncovalent interactions assisted para-C-H borylation of arenes (Scheme 1a, left). A series of para-C-H alkylations of arenes through alkyl radical addition have also been developed by Li, 8a Frost, 8b Zhao, 8c-e,i Zhou, 8f Zeng, 8g and others (Scheme 1a, right).8 However, these non-directed para-C-H functionalizations are restricted in transformation patterns and substrate classes. Alternatively, Maiti and co-workers developed

the first D-shaped nitrile-based *para*-directing template (DT) to assist several *para*-C–H functionalizations of toluene and phenol derivatives, demonstrating the possibility of using the DT strategy as a general means to achieve diverse *para*-C–H transformations (Scheme 1b, left). Dai and Yu also reported a *para*-C–H acetoxylation of benzoic acid derivatives assisted by an amide linked DT (Scheme 1b, right). Unfortunately, this DT



Scheme 1 Para-C-H activation beyond steric & electronic control.

[&]quot;State Key Laboratory of Structural Chemistry, Center for Excellence in Molecular Synthesis, Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences (CAS), 155 West Yang-Qiao Road, Fuzhou, Fujian 350002, P. R. China. E-mail: gangli@fjirsm.ac.cn

^bFujian College, University of Chinese Academy of Sciences, Beijing, 100049, China † Electronic supplementary information (ESI) available: Experimental procedures and data for new compounds. See DOI: 10.1039/d0sc07042d

strategy is restricted to very limited substrate classes at present. Most recently, by using the cooperation of a meta-DT and norbornene relay palladation, Yu and Houk11a as well as Maiti11b independently reported two examples of para-C-H arylation of arenes such as the phenylpropanoic acids. However, 2,6-disubstitution of the substrate is unavoidable for this method to achieve the desired selectivity, making it not a general solution. Due to the aforementioned limitations and the good performance of pyridine-based DGs in promoting ortho-1 and meta-C-H activations, 2d,12 the development of a versatile method for para-C-H activation of arenes by taking the advantage of the directing ability of pyridines is highly desirable.

Herein, we report an unprecedented Pd-catalyzed paraselective C-H alkenylation of three classes of arenes, i.e. phenylpropanoic acids, 2-phenyl benzoic acids and benzyl alcohols, with a series of alkenes including perfluoroalkenes assisted by a novel pyridine-based para-DT (Scheme 1c). Notably, the distance between the DT and the target C-H bond is longer than previous examples using the nitrile-based para-DT, 9,10 and the pyridine-based para-DT could be easily synthesized in two steps and readily recycled under mild conditions.

Results and discussion

Phenylpropionic acids are highly important structural motifs in drug molecules such as baclofen, and p-phenylenedipropionic acids is a class of important molecular unit of functional materials such as the gelator molecule of nanofiber. 13 Therefore, phenylpropionic acid was selected as the model substrate for our investigation of para-C-H activation of arenes. Based on our recent work on meta-C-H activation of phenylpropionic acid14 and inspired by recent research on meta-C-H activation assisted by pyridine-based DTs,12 we designed a series of potential pyridine-based para-DTs with different substitutions to attach with phenylpropionic acid, leading to the amide

Table 1 Evaluation of pyridine-based para-DT^a

substrates (Table 1, see ESI† for more DTs evaluated), where the nitrogen atom of pyridine could coordinate to the transition metal to induce site-selective C-H activation. The design of the structure of the template was also inspired by previous reported nitrile-based biphenyl containing para-directing template. 9,10 After selecting C-H alkenylation as the model reaction, it was found simple template without a substituent led to low yield and moderate selectivity $(1a_1)$. Introducing an electrondonating methoxy group resulted in no desired product (1a2). It was found an electron-withdrawing substituent, which could help to reduce the coordination ability of the pyridine group, was beneficial for the reaction to produce good yield of desired product $(1a_3-1a_7)$. Since the amide group could coordinate with the Pd catalyst to possibly influence the efficiency and selectivity of the reaction, the templates bearing an electron-withdrawing protecting group of the amide linkage to reduce the coordination ability were prepared, leading to higher selectivities ($1a_5$ vs. $1a_3$ and $1a_8$ vs. $1a_7$). Thus, the best preliminary results were obtained with substrate 1a₈ using Pd(OAc)₂ as the catalyst with 3-(trifluoromethyl)pyridin-2-ol (L1) as the ligand in HFIP (hexafluoroisopropanol) at 80 °C for 12 h, affording desired paraolefinated phenylpropionic acid derivative 3a₈ as the major product with good site-selectivity.

With the optimal pyridine-based para-DTs, we continued optimize the reaction conditions. After extensive

Table 2 Optimization of reaction conditions. a,b

Entry	Oxidant (equiv.)	Additive (equiv.) Yield (%) (<i>p</i> : others)
1	AgOAc (3)	_	$90(79)^{j}/15:1$
2^c	AgOAc (3)	_	70/11:1
3^d	AgOAc (3)	_	76/12:1
4^e	$Cu(TFA)_2$ (3)	_	21/7:1
5^e	CuCl ₂ (3)	_	_
6^e	CuI (3)	_	11/5:1
7^e	$Cu(OAc)_2$ (3)	_	40/8:1
8 ^f	$Cu(OAc)_2 (0.5)$	AcOH (2.0)	68/12:1
9 ^{f,g}	$Cu(OAc)_2 (0.5)$	AcOH (2.0)	72/12:1
$10^{f,h}$	$Cu(OAc)_2 (0.5)$	AcOH (2.0)	$80(72)^{j}/12:1$
$11^{f,i}$	$Cu(OAc)_2 (0.5)$	AcOH (2.0)	53/12:1
$12^{f,h}$	$Cu(OAc)_2 (0.5)$	AcOH (1.0)	76/12:1
$13^{f,h}$	$Cu(OAc)_2(0.5)$	AcOH (5.0)	73/12:1
	CF ₃	NOH CF3	Me 3u N t-Bu
	L1	L2 L3	L4

^a 1a₈ (0.1 mmol), Pd(OAc)₂ (0.01 mmol), 2a (2 equiv.), L1 (0.02 mmol), oxidant, additive, HFIP (1.5 mL), 90 °C, air (1 atm), 12 h. b Yield was determined by ¹H NMR with CH₂Br₂ as internal standard; products ratios were determined from crude ¹H NMR, and products of others were mainly *m*- and *o*-isomers. ^c **L2** (20 mol%). ^d **L3** (20 mol%). ^e 18 h. ^f 2a (3 equiv.), L4 (20 mol%), 60 °C, 48 h. ^g 70 °C. ^h 75 °C. ⁱ 80 °C. ^j Isolated yield of desired product in parentheses.

^a Reaction conditions: 1a (0.1 mmol), 2a (2 equiv.), Pd(OAc)₂ (0.01 mmol), 3-(trifluoromethyl)pyridin-2-ol (L1) (0.02 mmol), AgOAc (0.3 mmol), HFIP (1.0 mL), 80 °C, 12 h. Ratio = p: others; Ts: tosyl; Bs: phenylsulfonyl.

investigation of the reaction conditions (see ESI† for details), desired product 3a₈ was produced in 79% isolated yield with elevated temperature at 90 °C and lowered concentration based on the preliminary reaction conditions (Table 2, entry 1), together with excellent site-selectivity. The yield was decreased when other pyridine-based ligands were tested (entries 2 and 3). We then tried to use a cheaper copper salt as the oxidant, and Cu(OAc)₂ was found to be promising (entries 4-7). Pleasingly, a substoichiometric amount of Cu(OAc)₂ could be enough to produce higher yields of 3a8 with good site-selectivity (entries 8-11) in the presence of air as the terminal oxidant and two equivalents of AcOH as the additive (see ESI† for the evaluation of other acids) using L4 as the ligand for 48 h, and the best results using Cu(OAc)2 were obtained at 75 °C (entry 10). Notably, the mount of AcOH was also important, since slightly lower yields of 3a8 would be received with lower (entry 12) of higher (entry 13) loading of this additive.

With the optimized reaction conditions in hand, the generality of this reaction was investigated with a range of phenylpropionic acids as well as some representative olefins (Table 3). It was found that a number of substituents that are electron-donating such as methyl and methoxy groups or electron-withdrawing such as choloro, trifluoromethoxy, trifluoromethyl and fluoro groups at the *ortho-* or *meta-*

positions were tolerated in this reaction, delivering the target products in generally moderate to good yields with overall high *para*-selectivity (3a-3j). Notably, several di-substituted substrates were also viable in this reaction to afford generally good yields of desired products (3k-3p). Moreover, substitution at the benzylic position with a methyl (3q) or ethyl (3r) groups was tolerated. Interestingly, *para*-C-H functionalization of the phenylalanine derivative was also feasible (3s). Finally, a series of electron-deficient alkenes were examined to give generally good yields of the desired products with substrate 1h (3ha-3hi), albeit electron-rich alkenes were not viable.

Subsequently, perfluoroalkenes were also evaluated in our study since fluorine-containing compounds play an important role in drug development and material sciences. ¹⁵ Although *meta*-C-H perfluoroalkenylation of arenes had been disclosed, ¹⁶ there was no report of *para*-C-H perfluoroalkenylation of arenes. We were glad to find that this reaction was compatible with several perfluoroalkenes including brominated perfluoroalkene (Table 4). Phenylpropionic acid derivatives bearing mono-substitution at the benzylic position or on the aryl group were well tolerated to react with tridecafluorooct-1-ene to give desired products with excellent *para*-selectivity (4a-4h). Di-substituted substrates were also feasible to produce desired products (4i-4l). In

Table 3 Scope of phenylpropionic acids and alkenes^a

^a Reaction conditions A: **1** (0.1 mmol), Pd(OAc)₂ (10 mol%), **2** (3 equiv.), Cu(OAc)₂ (50 mol%), **L4** (20 mol%), AcOH (2 equiv.), air (1 atm), HFIP (1.5 mL), 75 °C, 48 h. ^b Conditions B: Oxidant: AgOAc (3 equiv.), ligand: **L1** (20 mol%), 24 h; no AcOH added. ^c **2** (2 equiv.), 90 °C. ^d O₂ (1 atm). ^e **2** (3 equiv.), HFIP (1.0 mL), 100 °C. ^f **3ha**₁ (51%, terminal alkene), **3ha**₂ (22%, internal alkene). Ratio = p: others. Isolated yield of pure desired para-isomer was reported herein.

Table 4 Reaction with perfluoroalkenes

^a Reaction conditions: 1 (0.1 mmol), Pd(OAc)₂ (10 mol%), 2 (3 equiv.), Cu(OAc)₂ (50 mol%), L4 (20 mol%), AcOH (2 equiv.), HFIP (1.5 mL), air (1 atm), 70 °C, 48 h. ^b AgOAc (3 equiv.), L1 (20 mol%), 80 °C, 24 h; no AcOH added. Ratio = p: others. Isolated yield of pure desired para-isomer was reported herein.


addition, other fluorinated alkenes were also evaluated to give moderate to good yields of the target molecules (4m-4p).

The generality of the newly designed pyridine-based para-DT was then tested with other types of molecular structures. Pleasingly, 2-phenyl benzoic acid derivatives bearing several substitution patterns were also smoothly alkenylated at the remote para-position with acrylate or fluorinated alkenes (Table 5, 6a-6i). Moreover, several benzyl alcohol derivatives were also tested, and good yields of desired products could be produced with good to excellent para-selectivity (6j-6p). It should also be mentioned that the scopes of these two types substrates were not as good as that of phenylpropionic acids at current stage.

Finally, the synthetic potential of our method was investigated briefly. The DT could be readily removed and recycled with excellent yield of para-alkenylated phenylpropionic acid under mild reaction conditions (Scheme 2a). The recovered DT (7) was identical to that synthesized from Suzuki coupling reaction (see ESI†). The reaction was able to be scaled up to 3 mmol to give about one gram of 3h using lower loading of Pd(OAc)₂ (5 mol%) (Scheme 2b). In addition, p-phenylenedipropionic acids, which could be synthesized from our para-alkenylated-products after hydrolysis and hydrogenation, were employed to synthesize gelator of nanofiber, including uncommon ones (Scheme 2c).13a

Table 5 Scope of 2-phenyl benzoic acids and benzyl alcohols^a

^a Reaction conditions: 5 (0.1 mmol), 2 (3 equiv.), Pd(OAc)₂ (10 mol%), L1 (20 mol%), AgOAc (3 equiv.), HFIP (1.0 mL), 100 °C, 24 h. b 2 (2 equiv.), HFIP (1.5 mL), 90 °C. Ratio = p: others. Isolated yield of pure desired para-isomer was reported herein.



Scheme 2 Synthetic Potential Investigation.

Conclusions

In conclusion, we have developed an unprecedented pyridine-based *para*-DT assisted, Pd-catalyzed *para*-selective C–H alkenylation of phenylpropanoic acids with a series of alkenes including perfluoroalkenes. This protocol was also compatible with 2-phenyl benzoic acids and benzyl alcohols with high *para*-selectivity. Notably, air could be used as the terminal oxidant for this reaction and the pyridine-based *para*-DT could be easily synthesized and readily recycled under mild conditions. It is expected these results would find application in rapid construction of *para*-substituted arenes and stimulate the exploration of novel methods for *para*-C–H functionalization of arenes.

Author contributions

X. C. and G. L. conceived the project. X. C. performed the experiments and developed the method. S. F. and M. Z. prepared some of the substrates. S. F., Y. G. and S. L. repeated some of the reactions and checked the data. G. L. directed the project and wrote the manuscript with the feedback from other authors.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We gratefully thank the financial supports from NSFC (Grant No. 22022111, 22071248, 21871257), the Natural Science Foundation of Fujian Province (2020J02008), the Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDB20000000), and the Youth Innovation Promotion Association of the Chinese Academy of Sciences (No. 2020306).

Notes and references

1 For recent reviews: (a) T. W. Lyons and M. S. Sanford, Chem. Rev., 2010, 110, 1147; (b) T. Satoh and M. Miura, Chem.-Eur. J., 2010, 16, 11212; (c) D. A. Colby, R. G. Bergman and J. A. Ellman, Chem. Rev., 2010, 110, 624; (d) C. S. Yeung and V. M. Dong, Chem. Rev., 2011, 111, 1215; (e) B.-J. Li and Z.-J. Shi, Chem. Soc. Rev., 2012, 41, 5588; (f) G. Song, F. Wang and X. Li, Chem. Soc. Rev., 2012, 41, 3651; (g) K. M. Engle, T.-S. Mei, M. Wasa and J.-Q. Yu, Acc. Chem. Res., 2012, 45, 788; (h) S. A. Girard, T. Knauber and C.-J. Li, Angew. Chem., Int. Ed., 2014, 53, 74; (i) O. Daugulis, J. Roane and L. D. Tran, Acc. Chem. Res., 2015, 48, 1053; (j) Z. Chen, B. Wang, J. Zhang, W. Yu, Z. Liu and Y. Zhang, Org. Chem. Front., 2015, 2, 1107; (k) T. Gensch, M. N. Hopkinson, F. Glorius and J. Wencel-Delord, Chem. Soc. Rev., 2016, 45, 2900; (l) G. He, B. Wang, W. A. Nack and G. Chen, Acc. Chem. Res., 2016, 49, 635; (m) R. Shang, L. Ilies and E. Nakamura, Chem. Rev., 2017, 117, 9086; (n) Y. Park, Y. Kim and S. Chang, Chem. Rev., 2017, 117, 9247;

- (o) P. Gandeepan, T. Muller, D. Zell, G. Cera, S. Warratz and L. Ackermann, *Chem. Rev.*, 2019, **119**, 2192; (p) Q. Zhang and B.-F. Shi, *Chin. J. Chem.*, 2019, 37, 647; (q) Y. Hu and C. Wang, *Acta Phys.-Chim. Sin.*, 2019, 35, 913; (r) S. Rej, Y. Ano and N. Chatani, *Chem. Rev.*, 2020, **120**, 1788; (s) A. Trowbridge, S. M. Walton and M. J. Gaunt, *Chem. Rev.*, 2020, **120**, 2613; (t) Y. Qiu, C. Zhu, M. Stangier, J. Struwe and L. Ackermann, *CCS*, 2020, **2**, 1529.
- 2 (a) S. R. Neufeldt and M. S. Sanford, Acc. Chem. Res., 2012, 45, 936; (b) J. Li, S. De Sarkar and L. Ackermann, Top. Organomet. Chem., 2015, 55, 217; (c) L. Ackermann and J. Li, Nat. Chem., 2015, 7, 686; (d) C. G. Frost and A. J. Paterson, ACS Cent. Sci., 2015, 1, 418; (e) J. A. Leitch and C. G. Frost, Chem. Soc. Rev., 2017, 46, 7145; (f) M. Font, J. M. Quibell, G. J. P. Perry and I. Larrosa, Chem. Commun., 2017, 53, 5584; (g) Y. Gao and G. Li, Aldrichimica Acta, 2017, 50, 61; (h) M. T. Mihai, G. R. Genov and R. J. Phipps, Chem. Soc. Rev., 2018, 47, 149; (i) C. Haldar and B. Emdadul Chattopadhyay, Tetrahedron Lett., 2018, 59, 1269; (j) M. Ghosh and S. De Sarkar, Asian J. Org. Chem., 2018, 7, 1236; (k) A. Dey, S. K. Sinha, T. K. Achar and D. Maiti, Angew. Chem., Int. Ed., 2019, 58, 10820; (1) J. Wang and G. Dong, Chem. Rev., 2019, 119, 7478; (m) G. Meng, N. Y. S. Lam, E. L. Lucas, T. G. Saint-Denis, P. Verma, N. Chekshin and J.-Q. Yu, J. Am. Chem. Soc., 2020, 142, 10571; (n) G. Liao, Y.-J. Wu and B.-F. Shi, Acta Chim. Sin., 2020, 78, 289.
- 3 (a) A. Dey, S. Maity and D. Maiti, Chem. Commun., 2016, 52, 12398; (b) X. Wang, D. Leow and J.-Q. Yu, J. Am. Chem. Soc., 2011, 133, 13864; (c) J. Karthikeyan and C. H. Cheng, Angew. Chem., Int. Ed., 2011, 50, 9880; (d) C. L. Ciana, R. J. Phipps, J. R. Brandt, F. M. Meyer and M. J. Gaunt, Angew. Chem., Int. Ed., 2011, 50, 458; (e) K. Sun, Y. Li, T. Xiong, J. Zhang and Q. Zhang, J. Am. Chem. Soc., 2011, 133, 1694; (f) X. C. Cambeiro, T. C. Boorman, P. Lu and I. Larrosa, Angew. Chem., Int. Ed., 2013, 52, 1781; (g) W. Liu and L. Ackermann, Org. Lett., 2013, 15, 3484; (h) A. M. Wagner, A. J. Hickman and M. S. Sanford, J. Am. Chem. Soc., 2013, 135, 15710; (i) Z. Wu, F. Luo, S. Chen, Z. Li, H. Xiang and X. Zhou, Chem. Commun., 2013, 49, 7653; (j) Z. Yu, B. Ma, M. Chen, H. H. Wu, L. Liu and J. Zhang, J. Am. Chem. Soc., 2014, 136, 6904; (k) Y. Saito, Y. Segawa and K. Itami, J. Am. Chem. Soc., 2015, 137, 5193; (1) N. A. Romero, K. A. Margrey, N. E. Tay and Nicewicz, Science, 2015, 349, 1326; G. B. Boursalian, W. S. Ham, A. R. Mazzotti and T. Ritter, Nat. Chem., 2016, 8, 810; (n) Y. Ma, B. Wang, L. Zhang and Z. Hou, J. Am. Chem. Soc., 2016, 138, 3663; (o) K. Naksomboon, J. Poater, F. M. Bickelhaupt and M. A. Fernandez-Ibanez, J. Am. Chem. Soc., 2019, 14, 6719; (p) K. Liu, S. Tang, T. Wu, S. Wang, M. Zou, H. Cong and A. Lei, Nat. Commun., 2019, 10, 639; (q) F. Berger, M. B. Plutschack, J. Riegger, W. Yu, S. Speicher, M. Ho, N. Frank and T. Ritter, *Nature*, 2019, 567, 223; (r) S. C. Fosu, C. M. Hambira, A. D. Chen, J. R. Fuchs and D. A. Nagib, Chem, 2019, 5, 417; (s) F. de Azambuja, M. H. Yang, T. Feoktistova, M. Selvaraju, A. C. Brueckner,

Edge Article

M. A. Grove, S. Koley, P. H. Cheong and R. A. Altman, *Nat. Chem.*, 2020, **12**, 489.

- 4 (a) H. Xu, M. Shang, H. X. Dai and J.-Q. Yu, *Org. Lett.*, 2015, 17, 3830; (b) Y. X. Luan, T. Zhang, W. W. Yao, K. Lu, L. Y. Kong, Y. T. Lin and M. Ye, *J. Am. Chem. Soc.*, 2017, 139, 1786.
- (a) Y. Nakao, Y. Yamada, N. Kashihara and T. Hiyama, J. Am. Chem. Soc., 2010, 132, 13666; (b) S. Okumura, S. Tang, T. Saito, K. Semba, S. Sakaki and Y. Nakao, J. Am. Chem. Soc., 2016, 138, 14699; (c) S. Okumura and Y. Nakao, Org. Lett., 2017, 19, 584; (d) L. Yang, K. Semba and Y. Nakao, Angew. Chem., Int. Ed., 2017, 56, 4853.
- 6 M. E. Hoque, R. Bisht, C. Haldar and B. Chattopadhyay, J. Am. Chem. Soc., 2017, 139, 7745.
- 7 (a) M. T. Mihai, B. D. Williams and R. J. Phipps, *J. Am. Chem. Soc.*, 2019, 141, 15477; (b) J. R. Montero Bastidas,
 T. J. Oleskey, S. L. Miller, M. R. Smith 3rd and
 R. E. Maleczka Jr, *J. Am. Chem. Soc.*, 2019, 141, 15483.
- 8 (a) X. Guo and C.-J. Li, Org. Lett., 2011, 13, 4977; (b) J. A. Leitch, C. L. McMullin, A. J. Paterson, M. F. Mahon, Y. Bhonoah and C. G. Frost, Angew. Chem., Int. Ed., 2017, **56**, 15131; (c) C. Yuan, L. Zhu, R. Zeng, Y. Lan and Y. Zhao, Angew. Chem., Int. Ed., 2018, 57, 1277; (d) G. Tu, C. Yuan, Y. Li, J. Zhang and Y. Zhao, Angew. Chem., Int. Ed., 2018, 57, 15597; (e) C. Yuan, L. Zhu, C. Chen, X. Chen, Y. Yang, Y. Lan and Y. Zhao, Nat. Commun., 2018, 9, 1189; (f) Z. Jiao, L. H. Lim, H. Hirao and J. S. Zhou, Angew. Chem., Int. Ed., 2018, 57, 6294; (g) P. Liu, C. Chen, X. Cong, J. Tang and X. Zeng, Nat. Commun., 2018, 9, 4637; (h) K. D. Mane, A. Mukherjee, K. Vanka and G. Suryavanshi, J. Org. Chem., 2019, 84, 2039; (i) W. T. Fan, Y. Li, D. Wang, S. J. Ji and Y. Zhao, J. Am. Chem. Soc., 2020, 142, 20524; (j) W. K. Tang, F. Tang, J. Xu, Q. Zhang, J. J. Dai, Y. S. Feng and H. J. Xu, Chem. Commun., 2020, 56, 1497; (k) J. Zhou, F. Wang, Z. Lin, C. Cheng, Q. Zhang and J. Li, Org. Lett., 2020, 22, 68.
- 9 (a) S. Bag, T. Patra, A. Modak, A. Deb, S. Maity, U. Dutta, A. Dey, R. Kancherla, A. Maji, A. Hazra, M. Bera and D. Maiti, J. Am. Chem. Soc., 2015, 137, 11888; (b) T. Patra, S. Bag, R. Kancherla, A. Mondal, A. Dey, S. Pimparkar, S. Agasti, A. Modak and D. Maiti, Angew. Chem., Int. Ed., 2016, 55, 7751; (c) A. Maji, S. Guin, S. Feng, A. Dahiya, V. K. Singh, P. Liu and D. Maiti, Angew. Chem., Int. Ed., 2017, 56, 14903; (d) A. Maji, A. Dahiya, G. Lu, T. Bhattacharya, M. Brochetta, P. Liu, G. Zanoni and D. Maiti, Nat. Commun., 2018, 9, 3582; (e) U. Dutta,

- S. Maiti, S. Pimparkar, S. Maiti, L. R. Gahan, E. H. Krenske, D. W. Lupton and D. Maiti, *Chem. Sci.*, 2019, **10**, 7426; (*f*) S. Pimparkar, T. Bhattacharya, A. Maji, A. Saha, R. Jayarajan, U. Dutta, G. Lu, D. W. Lupton and D. Maiti, *Chem.-Eur. J.*, 2020, **26**, 11558.
- 10 M. Li, M. Shang, H. Xu, X. Wang, H.-X. Dai and J.-Q. Yu, Org. Lett., 2019, 21, 540.
- (a) H. Shi, Y. Lu, J. Weng, K. L. Bay, X. Chen, K. Tanaka,
 P. Verma, K. N. Houk and J.-Q. Yu, *Nat. Chem.*, 2020, 12,
 399; (b) U. Dutta, S. Porey, S. Pimparkar, A. Mandal,
 J. Grover, A. Koodan and D. Maiti, *Angew. Chem., Int. Ed.*,
 2020, 59, 20831.
- 12 (a) L. Chu, M. Shang, K. Tanaka, Q. Chen, N. Pissarnitski, E. Streckfuss and J.-Q. Yu, ACS Cent. Sci., 2015, 1, 394; (b) Z. Jin, L. Chu, Y. Q. Chen and J.-Q. Yu, Org. Lett., 2018, 20, 425; (c) H. Xu, M. Liu, L. J. Li, Y. F. Cao, J.-Q. Yu and H. X. Dai, Org. Lett., 2019, 21, 4887; (d) M. Liu, L.-J. Li, J. Zhang, H. Xu and H.-X. Dai, Chin. Chem. Lett., 2020, 31, 1301; For pyrimidine-based meta-DT: (e) S. Bag, R. Jayarajan, R. Mondal and D. Maiti, Angew. Chem., Int. Ed., 2017, 56, 3182; (f) R. Jayarajan, J. Das, S. Bag, R. Chowdhury and D. Maiti, Angew. Chem., Int. Ed., 2018, 57, 7659; (g) S. Fang, X. Wang, F. Yin, P. Cai, H. Yang and L. Kong, Org. Lett., 2019, 21, 1841; (h) B. Jiao, Z. Peng, Z.-H. Dai, L. Li, H. Wang and M.-D. Zhou, Eur. J. Org. Chem., 2019, 2019, 3195.
- 13 (a) J. Liu, F. Yuan, X. Ma, D. Y. Auphedeous, C. Zhao, C. Liu, C. Shen and C.-L. Feng, Angew. Chem., Int. Ed., 2018, 57, 6475; (b) X. Wang, H. Alshammary, R. Zhang, A. Seifpour, J. T. Villalva, Z. Xu, C. Zheng, J. Li and X.-Y. Huang, Polyhedron, 2008, 27, 3439; (c) S. Tanaka, A. Yagyu, M. Kikugawa, M. Ohashi, T. Yamagata and K. Mashima, Chem.-Eur. J., 2011, 17, 3693; (d) S. Sanram, J. Boonmak and S. Youngme, Polyhedron, 2016, 119, 151; (e) S. Roelens and R. Torriti, J. Am. Chem. Soc., 1998, 120, 12443.
- 14 S. Li, H. Wang, Y. Weng and G. Li, *Angew. Chem., Int. Ed.*, 2019, **58**, 18502.
- 15 (a) K. Muller, C. Faeh and F. Diederich, Science, 2007, 317, 1881; (b) R. Berger, G. Resnati, P. Metrangolo, E. Weber and J. Hulliger, Chem. Soc. Rev., 2011, 40, 3496; (c) J. Wang, M. Sanchez-Rosello, J. L. Acena, C. del Pozo, A. E. Sorochinsky, S. Fustero, V. A. Soloshonok and H. Liu, Chem. Rev., 2014, 114, 2432.
- 16 M. Brochetta, T. Borsari, S. Bag, S. Jana, S. Maiti, A. Porta, D. B. Werz, G. Zanoni and D. Maiti, *Chem.-Eur. J.*, 2019, 25, 10323.