Volume 12 Number 15 21 April 2021 Pages 5333-5690

Chemical Science

rsc.li/chemical-science



ISSN 2041-6539



EDGE ARTICLE Dóra Papp and Gábor Czakó Facilitated inversion complicates the stereodynamics of an $S_N 2$ reaction at nitrogen center

Chemical Science

EDGE ARTICLE

Check for updates

Cite this: Chem. Sci., 2021, 12, 5410

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 26th January 2021 Accepted 12th March 2021

DOI: 10.1039/d1sc00490e

rsc.li/chemical-science

1. Introduction

The bimolecular nucleophilic substitution $(S_N 2)$ reaction is one of the most important reaction types in organic and biological chemistry. The traditional stereo-selective back-side-attack Walden inversion mechanism of S_N2 reactions at carbon center has been known for more than a hundred years,¹ however, it took half a century to understand its atomic-level details: the process goes through a transition state with a nucleophile-central atom-leaving group geometry and results stereospecifically in the inversion of configuration around the central C atom.² Although with a very low probability, the retention of the initial configuration around a C atom in a S_N2 reaction is also possible: through the front-side-attack mechanism, known also for decades,2-4 or following the recently revealed double-inversion reaction path,5-7 where Waldeninversion is preceded by a proton-abstraction-induced first inversion. Double inversion, which is shown to be a non-IRC reaction route,⁸ is especially important in the case of a strong nucleophile, such as F⁻, when it becomes the lower-energy, and thus the primary retention pathway.⁶ Note that numerous reaction mechanisms resulting in inversion of configuration

Facilitated inversion complicates the stereodynamics of an S_N2 reaction at nitrogen center⁺

Dóra Papp 🕩 * and Gábor Czakó 🕩 *

Bimolecular nucleophilic substitution (S_N2) reactions at carbon center are well known to proceed with the stereospecific Walden-inversion mechanism. Reaction dynamics simulations on a newly developed high-level *ab initio* analytical potential energy surface for the $F^- + NH_2Cl$ nitrogen-centered S_N2 and proton-transfer reactions reveal a hydrogen-bond-formation-induced multiple-inversion mechanism undermining the stereospecificity of the N-centered S_N2 channel. Unlike the analogous $F^- + CH_3Cl S_N^2$ reaction, $F^- + NH_2Cl \rightarrow Cl^- + NH_2F$ is indirect, producing a significant amount of NH_2F with retention, as well as inverted NH_2Cl during the timescale within the unperturbed NH_2Cl molecule gets inverted with only low probability, showing the important role of facilitated inversions *via* an $FH...NHCl^-$ -like transition state. Proton transfer leading to $HF + NHCl^-$ is more direct and becomes the dominant product channel at higher collision energies.

were also reported for S_N2@C, such as roundabout,9 H-bonded complex formation,10 direct rebound11 and stripping11 pathways;12 as well as an interesting leaving-group effect13 and frontside complex formation¹⁴ have also been revealed. However, even with strong nucleophiles, in $S_N 2$ (a)C reactions the system usually has to overcome a substantial energy barrier to follow a retention pathway, while Walden inversion is often a barrierless (energies of all involved stationary points are below the reactant asymptote) route, supported by deep pre- and postreaction ion-dipole, or H-bonded minima.7,12 Thus, inversion is unequivocally the dominant outcome of a C-centered S_N2 reaction ensuring almost exclusive stereospecificity. Replacing the central carbon atom with nitrogen, having at most three ligands and the capability of their self-inversion through a rather low-energy barrier,15 as well as different electronic structure and symmetry properties than carbon, is expected to make both the energetics and the stereodynamics of an S_N2 reaction considerably distinct from those of S_N2@C.

 S_N2 reactions at N center were found to play an important role in carcinogenesis¹⁶⁻¹⁸ in the early 1990s arousing interest in these types of reactions. Double labeling experiments then suggested the presence of a classic transition state for an S_N2 reaction at N center¹⁹ and later a twice larger reactivity was observed for the F^- + NH₂Cl reaction when compared to F^- + CH₃Cl in gas-phase selected ion flow tube experiments.²⁰ Theoretical investigations of the S_N2 reactions at N center began with the characterization of the potential energy surfaces (PESs) of X⁻ + NH₂Y and N(CH₃)₂Y [X, Y = F, Cl, Br, I] using lower-level *ab initio*²¹⁻²⁴ and density functional theory (DFT) methods.²⁵⁻²⁷ Later, Bickelhaupt and co-workers studied the impact of replacing the central carbon atom in S_N2 reactions with several

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence. Open Access Article. Published on 24 March 2021. Downloaded on 8/9/2025 11:55:41 AM



View Article Online

View Journal | View Issue

MTA-SZTE Lendület Computational Reaction Dynamics Research Group, Interdisciplinary Excellence Centre and Department of Physical Chemistry and Materials Science, Institute of Chemistry, University of Szeged, Rerrich Béla tér 1, Szeged H-6720, Hungary. E-mail: dorapapp@chem.u-szeged.hu; gczako@chem. u-szeged.hu

[†] Electronic supplementary information (ESI) available: Structures, energies, harmonic frequencies and T_1 diagnostics of the stationary points of the F⁻ + NH₂Cl reaction, entrance channel 1D scan of the PES and snapshots from a quasi-classical trajectory. See DOI: 10.1039/d1sc00490e

other atoms by performing an activation strain analysis.^{28,29} In 2018 one of the present authors reported benchmark classical CCSD(T)-F12b/complete-basis-set-quality and adiabatic stationary-point energies and geometries taking also into account the correlation of core electrons both for identity and non-identity X^- + NH₂Y [X, Y = F, Cl, Br, I] reactions.³⁰ The energetics of the OH⁻ + NH₂Cl³¹ and the F⁻ + NH₂Cl³² reactions was investigated in water as well by Wang and co-workers using a QM/MM approach. Not only the PESs, but the dynamics of S_N2@N reactions were also studied in the framework of direct dynamics simulations, which compute the gradients of the PES "on-the fly" and therefore allow using only low-level ab initio or DFT methods, and running a few hundred/thousand of quasiclassical trajectories. First, the $OH^- + NH_2Y [Y = F, Cl]^{33}$ and the F⁻ + NH₂F³⁴ reactions were subjects of such calculations using the MP2 method with small basis sets, followed by the direct dynamics investigation of the F^- + NH₂Cl reaction in 2017 by Liu et al.35 at the B3LYP/aug-cc-pVDZ level of theory. They proposed indirect dynamics for the S_N2 reaction and suggested minor competiveness of the proton-transfer channel even at the higher of the two collision energies considered. They also observed various S_N2 mechanisms, such as direct rebound, direct stripping and indirect hydrogen-bonding. Li and Wang later proposed a proton-abstraction roundabout mechanism followed by Walden-inversion for the $F^- + NH_2Cl \rightarrow Cl^- + NH_2F$ reaction based on M06-2X/aug-cc-pVDZ-level direct dynamics simulations, along with identifying a double-inversion transition state and the corresponding reaction path.³⁶

Taking a significant step forward from the previous direct dynamics investigations, here we report the first fulldimensional global analytical PES for the F^- + NH₂Cl reaction fitted on high-quality CCSD(T)-F12b/aug-cc-pVTZ energy points, which allows for a detailed and statistically reliable quasiclassical dynamics study based on more than half a million trajectories covering a wide range of collision energies. Moreover, our analysis makes it possible to track the change of the initial configuration, and therefore is capable of providing new insights into the so far unstudied stereodynamics of this prototypic $S_N 2$ @N reaction.

2. Methods

As a starting point for the potential energy surface development we randomly displace the Cartesian coordinates of the CCSD(T)-F12b/aug-cc-pVTZ optimized geometries of the stationary points of the F^- + NH₂Cl reaction taken from ref. 30 in the 0.0– 0.4 Å interval, and in the case of the reactants and products we also scatter the two fragments in random orientation around each other in the 2.5–15.0 Å distance range. The random geometries generated this way are then subjected to MP2/augcc-pVDZ^{37,38} single-point computations, and the obtained dataset is cut at a 100 kcal mol⁻¹ energy threshold above the global minimum of the set. The energies of the remaining 5775 geometries are recalculated at the CCSD(T)-F12b/aug-ccpVTZ^{38,39} level of theory, and then serve as the starting dataset for the automated PES development carried out with the ROBOSURFER program package,⁴⁰ using also the same level of theory. For all quantum chemical computations the MOLPRO program package⁴¹ is used. The permutationally invariant polynomial method^{42,43} is applied for fitting the PES ensuring that it is invariant under the permutation of like atoms. For this, the potential energy points are fitted using a full-dimensional analytical function which is an expansion of the polynomials of the $y_{ij} = \exp(-r_{ij}/a)$ Morse-like variables with r_{ij} being the atom-atom distances and the *a* parameter, which defines the asymptotic behavior of the functions, is 3.0 bohr. The highest polynomial order used for this PES is 6, and the fit requires 4285 coefficients. A weighted least-squares fit is carried out, where a given energy E relative to the global minimum has a weight of $(E_0/(E_0 + E)) \times (E_1/(E_1 + E))$ with $E_0 = 0.1$ hartree and $E_1 = 0.5$ hartree. In the ROBOSURFER program a hard upper limit of 100 kcal mol^{-1} is set relative to the energy of the reactants. During development a target accuracy of 0.5 kcal mol^{-1} is demanded up to 50 kcal mol⁻¹ relative to the reactants. New geometries, carefully selected and added to improve the PES, are generated in quasi-classical trajectory (QCT) simulations and by the Holebuster subprogram.⁴⁰ ROBOSURFER iterations are performed at the following collision energies (kcal mol^{-1}) used in the dynamics computations with the maximal values of the *b* impact parameter (the distance between the velocity vectors of the reactants in bohr) in parentheses: 64, 27, 13, 11, 13, 27, 17, 62, 36, and 18 iterations with 1.0(20.0), 5.0(15.0), 10.0(10.0), 20(10.0), 30(10.0), 40(10.0), 50(10.0), 80(10.0),100(10.0) and 50(10.0), in the last bunch of iterations executing the Holebuster subprogram too. The final PES is built of 16 780 geometries and the corresponding CCSD(T)-F12b/aug-cc-pVTZ energies, and features 0.56(0.93) kcal mol⁻¹ root mean square deviations in the 0–63(63–126) kcal mol^{-1} interval with respect to the global minimum of the final dataset.

Quasi-classical trajectory simulations are performed at seven collision energies: 0.9, 6.9, 10.0, 20.0, 30.0, 40.0, and 46.1 kcal mol⁻¹ to investigate the dynamics of the F⁻ + NH₂Cl reaction. The spatial orientation of the reactants is randomly sampled, and the initial distance between the F^- ion and the center of mass of the NH₂Cl molecule is set to $\sqrt{x^2 + b^2}$ bohr, where x = 30.0 bohr and the *b* impact parameter is varied between 0 and b_{max} , the distance where the reaction probability becomes zero, with a step size of 1.0 bohr. 5000 trajectories are run at each b value; thus the total number of trajectories is more than half a million. At the beginning of the trajectories the zeropoint energy (ZPE) of the NH₂Cl molecule is set by using standard normal mode sampling.44 Each trajectory is propagated with a 0.0726 fs time step until the largest interatomic distance becomes larger than the largest initial one by 1 bohr. 2000 trajectories with zero collision energy are run until a 500 000 time-step limit to monitor the configuration change of the unperturbed NH₂Cl reactant in its ground vibrational state while the F^- ion is placed 50 bohr far from NH₂Cl. Integral cross-sections (ICSs) of the two reaction channels are calculated by a *b*-weighted numerical integration of the P(b) opacity functions at each collision energy. For the proton-transfer channel, three different ZPE-constraints are set for the reaction probabilities and ICSs: (1) soft: the sum of the classical vibrational energy of the NHCl⁻ product and the internal energy of the HF

product must be larger than the sum of the ZPE of NHCl⁻ and the ZPE corresponding to the actual rotational state of HF, (2)hard: the above restrictions are set separately for each product, and (3) NHCl⁻ ZPE: the restriction is set only for NHCl⁻. The variationally determined rovibrational energy levels of the HF molecule are taken from ref. 45. The scattering angle distributions of the products are obtained by binning the cosine of the angle (θ) of the relative velocity vectors of the products and the reactants into 10 equidistant bins from -1 to 1. $\cos(\theta) = -1$ (θ $= 180^{\circ}$) corresponds to backward scattering. The rotational quantum number of the HF product is determined as detailed in ref. 46. The configuration of the NH₂Cl reactant (and the 'product' in non-reactive trajectories) and the NH₂F product is identified as retained (retention) if the sign of the scalar product of the normal vector of the HNH plane and the NX (X = Cl, F)vector was the same as in the initial reactant configuration, otherwise it is identified as inverted (inversion).47

3. Results and discussion

Compared to the analogous S_N^2 reaction at carbon center, $F^- + CH_3Cl \rightarrow Cl^- + CH_3F$, the energy landscape of the $F^- + NH_2Cl \rightarrow Cl^- + NH_2F$ reaction, shown in Fig. 1, is markedly different. The $F^- + NH_2Cl$ reaction also follows the two typical pathways of an ion-molecule reaction: S_N^2 and proton transfer (PT). The S_N^2 path features a -13.6 kcal mol⁻¹ exothermicity, which is significantly less than the -31.9 kcal mol⁻¹ value for the C-centered reaction, while the PT channel has only a 4.0 kcal mol⁻¹ adiabatic endothermicity, opening at a much lower energy than in the case of carbon center.⁵ The $F^- + NH_2Cl \rightarrow HF + NHCl^-$ reaction has a submerged barrier, -9.7 kcal mol⁻¹, considerably deeper than for $F^- + CH_3Cl \rightarrow HF + CH_2Cl^-$ (12.6 kcal mol⁻⁵), and both lack a kinetic barrier. Based on the above, the proton-transfer channel promises to be

much more competitive with the S_N2 path at N-center. As also seen in Fig. 1, the adiabatic barrier height for the inversion of the reactant NH₂Cl molecule is only 7.9 kcal mol⁻¹, which makes self-inversion possible to occur even at low energies, turning the stereodynamics of the S_N2 pathway more complex and thus more difficult to predict, in sharp contrast to the $S_N 2$ reaction at C-center, where there is no possibility for the reactant to be self-inverted. The barrier of the usual back-side-attack Walden-inversion mechanism of the N-centered S_N2 reaction is also negative $(-11.9 \text{ kcal mol}^{-1})$ and is of similar relative energy to that of the C-centered one, however, the former reaction is missing the typical pre-reaction arrangement, a prior minimum structure with high symmetry,^{7,12,48} to reach the Walden transition state (W TS), where a nucleophile-central atom bond is being formed. In the case of the F^- + NH₂Cl reaction only a Hbonded minimum (H premin) is found in the entrance channel $(-30.2 \text{ kcal mol}^{-1})$ ³⁰ twice as deep as for the F⁻ + CH₃Cl reaction, relative to which the proton-transfer transition state (PT TS) lies only slightly higher than the Walden-TS, in contrast to the C-centered case, where the PT TS lies 12.6 kcal mol^{-1} above the reactant asymptote.5 In addition, the configurational space covered near the H premin is more favorable for proton transfer, which makes it more likely to occur while preventing the system from passing through the Walden-TS easily. Accordingly, it is not surprising that the visual analysis of the dynamics simulations shows that the system spends most of the 'prereaction time' near the H premin. On the other hand, from the H premin arrangement, the most favorable transformation is not proton-transfer, but an inversion through a H-bonded, FH...NHCl⁻-like transition state, called traditionally as the double-inversion TS (DI TS),5 also observed in the direct dynamics computations of Wang and co-workers.³⁶ In contrast to the case of carbon center, where it has a 16.4 kcal mol^{-1} relative energy,^{5,6} the DI TS lies here quite deep,



Fig. 1 Schematic representation of the energetics and possible reaction paths of the $F^- + NH_2Cl$ reaction showing classical and adiabatic relative energies of the minima and transition states comparing benchmark values³⁰ with those obtained on the newly developed PES. For details about the stationary points see the ESI.[†]

-23.8 kcal mol⁻¹ below the reactants. As seen in Fig. 1, the Hbond forming between the reactants reduces the adiabatic inversion barrier of NH₂Cl from 7.9 kcal mol⁻¹ to 5.1 kcal mol⁻¹ (owing probably to the weakened N-H bond), and thus due both to the configurational and energetic preferences (about one quarter of the barrier height) from the H premin to the pre-DI TS with respect to the Walden-TS, several induced inversion events through the pre-DI TS are expected to occur before finding a favorable arrangement for the substitution to take place. In the exit channel a similarly deep H-bonded postreaction minimum (H postmin) can be found, however, with a considerably higher-lying post-DI TS, energetically close to the product asymptote. This is another major difference compared to the analogous $F^- + CH_3Cl \rightarrow Cl^- + CH_3F$ reaction, where the post-DI TS lies 66.1 kcal mol⁻¹ higher than the reactants⁶ and 98.0 kcal mol⁻¹ higher than the products,^{6,48} and thus inversion through this TS is almost negligible, while the product asymptote is only 9.7 kcal mol⁻¹ above the H postmin complex, driving the system to form the products rather directly. In the F^{-} + NH₂Cl case a front-side pre-reaction complex (FS premin) featuring a halogen-halogen bond, known to have an important role in S_N2 dynamics for C center,^{13,14} can also be identified submerged below the reactants $(-12.1 \text{ kcal mol}^{-1})$, significantly deeper than the -2.7 kcal mol⁻¹ value observed for F⁻ + CH₃Cl.¹⁴ Such a deep minimum with a non-reactive orientation is expected to have a significant prohibiting effect on the reaction dynamics as it was shown for the $F^- + CH_3I \rightarrow I^- + CH_3F$ reaction.14 The front-side-attack transition state (FS TS) for the $F^- + NH_2Cl \rightarrow Cl^- + NH_2F$ reaction lies relatively high above the reactants (almost at the same height as for $F^- + CH_3Cl^5$), thus if the NH₂F product is found with retained configuration at low collision energies, the system is sure to have followed a double or more likely a multiple-inversion mechanism through the DI TSs. Taken together, owing to the deep non-reactively oriented and H-bonded minima, the H-bond-lowered barrier height of inversion at the pre-DI TS, the lack of the traditional N-Fbonded pre- and post-reaction minima, the relatively high Walden-TS from the H premin and the strongly competitive proton-transfer channel, we can expect much more indirect dynamics for the S_N2 reaction at N-center than at C-center. Furthermore, the indirect reaction proceeding through a mechanism that is very likely to involve multiple configuration inversions also raises several questions regarding stereospecificity in S_N2@N.

To test the hypotheses made based purely on the energy landscape characterized above, we investigate the dynamics of the F^- + NH₂Cl reaction in detail. For this we first develop a high-quality full-dimensional *ab initio* analytical PES using a permutationally invariant fitting of the gold-standard CCSD(T)-F12b/aug-cc-pVTZ energy points, on which we run quasi-classical trajectories at seven collision energies, 0.9, 6.9, 10.0, 20.0, 30.0, 40.0 and 46.1 kcal mol⁻¹, covering the chemically interesting energy range of the reaction. Looking at Fig. 2 we can see that the excitation function, *i.e.* the integral crosssections (ICSs) as a function of collision energy, of the S_N2 channel decreases sharply, as it is expected in the case of a barrierless highly exothermic reaction. For the proton-transfer



Fig. 2 Integral cross-sections (ICSs) as a function of collision energy for the S_N2 and proton-transfer (PT) product channels of the F^- + NH_2CI reaction obtained on the newly developed PES. The inset shows ICS values obtained with different zero-point-energy constraints regarding the products of the PT reaction (for details see the Methods section).

ICSs we applied different ZPE-constraints (see the Methods section), among which the hard-restricted ICSs, where each product must have higher vibrational energy then its own ZPE, are the most realistic. ZPE-constraints seem to have a significant effect on the reactivity, changing the fast decaying shape of the unrestricted excitation function to a much flatter one with a maximum and in the soft- and hard-constrained cases also with a threshold. After the proton-transfer channel becomes thermodynamically available at around 4 kcal mol⁻¹ collision energy, it features a commeasurable reactivity with $S_N 2$, and slightly above 10 kcal mol⁻¹ it turns to be the dominant reaction path, in contrast to the findings of the statistically considerably less reliable direct dynamics computations of Liu *et al.*³⁵

Fig. 3(A) shows that the probability for the $S_N 2$ reaction to occur is the highest at the lowest collision energy, reaching even 40% at zero impact parameter, and decreases rapidly with increasing collision energy. This demonstrates that translational energy basically inhibits a barrierless exothermic reaction by reducing the time available for finding a favorable condition (orientation or vibrational phase) for reaction. The maximum value of the impact parameter, where the reactivity vanishes, is the largest (26 bohr) at 0.9 kcal mol^{-1} collision energy, and decreases fast as translational energy increases, underlining the promoting effect of reaction time by allowing the ion-dipole interaction between the reactants to guide the substitution reaction. The monotonically decaying shape of the opacity functions, *i.e.* the reaction probability as a function of the impact parameter, of Fig. 3(A) along with the scattering angle distributions (Fig. 3(B)), which show an increasing preference for forward over backward scattering with increasing collision energy, give rise to a direct rebound mechanism at lower translational energies, occurring mostly at small impact parameters and a more and more dominative direct stripping mechanism at higher energies. The latter takes place at large



Fig. 3 (A) Opacity functions (reaction probabilities as a function of the *b* impact parameter), (B) product scattering angle distributions, (C) internal energy distributions of the NH₂F product (the vertical black line refers to the ZPE of the product) and (D) relative translational energy distributions of the products as a function of collision energy (given in kcal mol⁻¹ and denoted by different colors) for the $F^- + NH_2Cl \rightarrow Cl^- + NH_2F S_N2$ reaction obtained on the newly developed PES.

impact parameters, where the approaching F^- ion strips away the NH₂ fragment of the reactant molecule while retaining its initial direction. However, the primarily isotropic scattering angles indicate a rather indirect reaction. This is in contrast to the C-centered analogous reaction,48 where forward scattering and the corresponding stripping mechanism are mostly observed at low collision energies, and the overall dynamics is essentially direct. The internal energy distribution of the NH₂F product, seen in Fig. 3(C), peaks near the maximum accessible internal energies (the sum of the ZPE of the reactant, the collision energy and the absolute value of the classical reaction energy) at lower collision energies, which indicates a highly indirect reaction where almost all the available energy, which is considerable due to the high exothermicity, is trapped in the internal degrees of freedom while the system is stuck in the preand post-reaction H-bonded minima and/or DI TSs inducing repetitive inversions. The elongated N-F bond in the W TS structure with respect to that in NH₂F³⁰ is also expected to lead to vibrationally highly excited products.49 The translational energy distribution of the S_N2 products, shown in Fig. 3(D), peaking near zero (except the two highest energies), also underlies the indirect mechanism, implying that the products can only separate with minimal translational energy at the rare

occasions when enough of the excess internal energy flows into the translational degree of freedom. As collision energy increases, the maxima gradually shift toward higher energies and the internal energy distributions adopt a more Gaussianlike shape as a sign of an increasingly direct reaction, in consistence with the broadening relative translational energy distributions. As seen from the internal energy distributions, the number of ZPE violating trajectories is practically negligible.

After the opening of the proton-transfer channel the hard ZPE-constrained opacity functions of Fig. 4(A) show enhancing reaction probability as collision energy increases. Note that at $0.9 \text{ kcal mol}^{-1}$ the hard-restricted reaction probability is zero, in accord with the 3.8 kcal mol⁻¹ adiabatic endothermicity and the experiments of ref. 20. We have seen in Fig. 1 that PT is also a barrierless reaction, however, the system is much less likely to stick in the deep H-bonded pre-reaction minimum, from which the PT TS is almost as far as the W TS, but the configuration of the H premin favors proton transfer. This and a shallower exit-channel well give rise to a more direct mechanism, that is why the increase of the initial translational energy promotes the reaction noticeably. At high collision energies the stripping mechanism becomes dominant as seen from the substantial reaction probabilities at relatively large impact parameters of 3–



Fig. 4 (A) Opacity functions (reaction probabilities as a function of the *b* impact parameter) obtained with a hard ZPE constraint, (B) product scattering angle distributions, (C) internal energy distributions of the NHCl⁻ product (the vertical black line refers to the ZPE of the product) and (D) relative translational energy distributions of the products as a function of collision energy (given in kcal mol⁻¹ and denoted by different colors) for the $F^- + NH_2Cl \rightarrow NHCl^- + HF$ proton-transfer reaction obtained on the newly developed PES.

6 bohr and from the forward-scattering preference of the scattering angle distributions of the products (Fig. 4(B)) in these cases. The b_{max} values are much smaller for the PT reaction than for S_N2, the largest is 13 bohr at 10 kcal mol⁻¹ energy, as another sign of direct reaction dynamics. From the internal energy distributions of the NHCl⁻ product, shown in Fig. 4(C), it is also clear that a large number of the low-energy trajectories is ZPE-violating, making it necessary to apply the above-described ZPE-constraints. Fig. 4(C) and (D) also suggest a more direct reaction than it is observed in the case of the S_N2 channel with colder and broader Gaussian-like internal energy distributions along with hotter and broader relative translational energy distributions of the products both with blue-shifting maxima as collision energy increases.

After revealing a considerably more indirect $S_N 2$ reaction dynamics, caused mainly by the deep H-bonded minima and low-lying submerged TSs for nitrogen center compared to C center, we arrive at the most interesting question: how much stereospecificity is affected by the central atom? The low-barrier umbrella motion of the ligands around the N atom has two consequences that could play an important role in the stereodynamics: (1) on one hand, unlike the case of carbon center, at nitrogen center self-inversion is a feasible internal motion of the NH₂Cl reactant and of the NH₂F product (with only

7.9 kcal mol⁻¹ and 14.4 kcal mol⁻¹ adiabatic barrier heights, respectively, the latter being higher due maybe to the stronger N-F bond relative to N-Cl); (2) on the other hand, the H-bond formed with the F⁻ ion further lowers this barrier height for the reactant (to 5.1 kcal mol⁻¹), while the somewhat weaker Hbond with Cl⁻ does not really affect the inversion barrier of the product (15.2 kcal mol^{-1}), but still leaves the door open for the inversion to take place in the exit channel as well. To investigate the time-scale and impact of (1), we analyzed the motion of the NH₂Cl molecule in an unperturbed state, with zero collision energy and allowing only zero-point vibration. The fraction of retention and inversion of NH2Cl as the trajectories are propagated in time is shown in Fig. 5(A). It can be seen that the initial configuration starts to invert after 1 ps without any external action and the average retention-inversion ratio gradually converges to the racemic limit, however, at quite a low pace: it is still 58-42% after 36 ps. To study the effect of the inversion motion induced by the H-bond formed with F⁻ we also analyze the non-reactive trajectories (with NH2Cl as 'product') run at different collision energies, and show the ratio of those that result in the retention of configuration in Fig. 5(B). Taking a look at the long trajectories of the low collision energies (the time distribution of the non-reactive trajectories is plotted in Fig. 5(C)) one can observe that above a certain trajectory length



Fig. 5 (A) Fraction of unperturbed NH_2CI molecules in the ground vibrational state found with retained or inverted initial configuration as a function of time (for further details see the Methods section). (B) Ratios of ICSs of non-reactive trajectories whose NH_2CI outcome is found with retention of configuration as a function of trajectory length at different collision energies (kcal mol⁻¹). (C) Time-distributions of non-reactive trajectories at different collision energies (kcal mol⁻¹). (D) Ratio of the retention and inversion ICSs as a function of collision energy in the case of S_N2 and non-reactive trajectories (the inset zooms in the S_N2 data).

NH₂Cl comes with roughly 50% probability in inverted or in retained configuration, which means that stereospecificity basically vanishes in these cases. With increasing collision energy, the length of the trajectories decreases, however, due to faster collisions, inversion may take place over a shorter period of time, which is indicated in Fig. 5(B) by a faster 'convergence' to the racemic limit with increasing trajectory length. By comparing Fig. 5(A) and (B) it is clear that induced inversion plays a significant role in the dynamics, since non-reactive trajectories reach a near racemic average outcome usually in 2-4 ps, while only 10% of the NH₂Cl molecules are inverted after 4 ps in an unaffected zero-point vibrational state. It is also true for the lowest 0.9 kcal mol⁻¹ collision energy (see insets of Fig. 5(B) and (C)), where the 50–50% probability is reached at about 12 ps trajectory length, and kept smoothly for longer trajectories as well, whereas the unperturbed reactant is found inverted only in 20% at 12 ps. The above observations suggest that a pre-reaction inversion can readily occur and it is most likely to happen through the pre-DI TS. As seen from Fig. 3(C), NH₂F is produced with considerable internal energy owing to the high exothermicity, and, due to its formation via inversion at Walden-TS a significant part of the excess energy may be responsible to excite the umbrella motion of the ligands. Thus, inversion of the product, despite its higher barrier than that of

the reactant, can also take place easily both through the post-DI TS or on its own. Moreover, not only one, but multiple inversions can and do occur both in the entrance and exit channels, since longer trajectories eventually reach a racemic outcome of the S_N2 reaction, as shown in Fig. 5(B). These findings are also strengthened by the visual analysis of S_N2 trajectories showing that the system frequently alters the configuration around the N atom through the pre-DI TS. Visual inspection also underlies the post-reaction inversion mechanisms, however, while the attacking F⁻ ion recurrently abstracts one of the protons from N, the product Cl⁻ ion seems very unlikely to be capable of that. We also observe a roaming/roundabout mechanism⁵⁰ in the entrance channel: after proton-abstraction the HF molecule does not have enough energy to drift from the NHCl⁻ fragment, and thus it begins to stroll around the rotating NHCl⁻ until finally putting the proton back. This process can then be followed by actual proton-transfer, but can precede an S_N2 reaction as well, in consistence with the observations of Li and Wang.36 Such an indirect pathway makes the reaction dynamics even more complex. Selected snapshots from QCT simulations of the above mechanisms are presented in Fig. S1 in the ESI.† As shown in Fig. 5(D) the impact of the multiple-inversion mechanism is also reflected in the ICS values of the S_N2 channel: at the lowest collision energy the retention/inversion ICS ratio is

almost 1 and it slowly decays to 0.2 as trajectories are getting shorter with increasing collision energy. This finding is in sharp contrast to what is observed for the analogous C-centered reaction, where the ICS fraction of retention is smaller than 2%,⁵ showing the clear stereospecificity of the Walden-inversion mechanism for S_N2(a)C. The H-bond-induced inversion in the case of non-reactive trajectories is also substantial: the retention/inversion ratio at the lowest collision energy is only 2.3, indicating that if enough time is available, induced inversion is very likely to occur. With higher energies the trajectory length is dropping rapidly which prevents effectively the inversion of configuration. The energetics of the $F^- + NH_2Cl \rightarrow Cl^- +$ NH₂F reaction explored in water by Wang and co-workers³² suggests that although the barrier heights of both the pre-DI TS and the W TS become positive, the former is much less increased, and thus the difference between the two increases by about 50%, this way promoting the pre-reaction inversional motion in aqueous solution. Taken together, replacing the central carbon atom with nitrogen in an S_N2 reaction inherently questions its stereospecificity, so far taken for granted regarding this family of chemical reactions, and although it raises new complexities, it might open new doors as well.

4. Conclusions

In the present study we have developed a full-dimensional global analytical potential energy surface fitted on CCSD(T)-F12b/aug-cc-pVTZ energy points for the F⁻ + NH₂Cl reaction, which allows for performing accurate dynamics simulations based on more than half a million quasi-classical trajectories covering a wide range of collision energies. The F⁻ + NH₂Cl \rightarrow Cl⁻ + NH₂F reaction serves as a prototypic S_N2@N reaction, for which we report the first detailed dynamical investigations. Furthermore, our newly developed PES and the size of the system open the route for studying the dynamics of the F⁻ + NH₂Cl reaction by quantum methods in the near future, even in full dimensions. Such simulations will be able to complement the present QCT results by capturing tunneling effects which can be important during inversion or when proton transfer takes place.

It is clear from the internal, translational and scattering angle distributions of the products that, in contrast to its C-centered analogue, the N-centered S_N2 reaction exhibits essentially indirect characteristics, originating from deep pre- and post-reaction H-bonded minima, low lying H-bonded inversion transition states, the lack of traditional ion-dipole minima, and a strongly competitive proton-transfer reaction. Increasing initial translational energy is found to counteract the exothermic barrierless S_N2 reaction, allowing less time for finding a favorable condition for reaction; whereas it enhances the reactivity of the endothermic (but also barrierless) proton-transfer pathway, which becomes dominant above 10 kcal mol⁻¹ collision energy. The latter turns out to feature more direct dynamics, being less likely to get stuck in deep minima.

Our simulations, capable of tracking the configurational changes around the central N atom, feasible even at room temperature and further facilitated by the H-bond formed with the attacking F^- ion, also give new insights into the so far unstudied stereodynamics of this prototypical $S_N 2$ @N reaction. Compared to $S_N 2$ @C, where stereospecificity *via* Walden inversion is the conventional route, our simulations clearly demonstrate the important role of the H-bond-induced multiple-inversion mechanisms at N center, accelerating the inversion of NH₂Cl compared to the unperturbed reactant and producing a significant fraction of the NH₂F products with retention of the initial configuration. While double inversion is a minor pathway for $S_N 2$ @C, its multiple-inversion analogue can be the key mechanism for N-centered $S_N 2$ reactions.

Author contributions

G. C. designed research, D. P. performed research and analyzed data, G. C. and D. P. discussed the results and D. P. wrote the paper.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank the National Research, Development and Innovation Office–NKFIH, K-125317; the Ministry of Human Capacities, Hungary Grant 20391-3/2018/FEKUSTRAT; and the Momentum (Lendület) Program of the Hungarian Academy of Sciences for financial support.

References

- 1 P. Walden, Ber. Dtsch. Chem. Ges., 1896, 29, 133.
- 2 C. K. Ingold, *Structure and Mechanisms in Organic Chemistry*, Cornell Univ. Press, Ithaca, NY, 1953.
- 3 M. N. Glukhovtsev, A. Pross, H. B. Schlegel, R. D. Bach and L. Radom, *J. Am. Chem. Soc.*, 1996, **118**, 11258.
- 4 A. P. Bento and F. M. Bickelhaupt, J. Org. Chem., 2008, 73, 7290.
- 5 I. Szabó and G. Czakó, Nat. Commun., 2015, 6, 5972.
- 6 I. Szabó and G. Czakó, J. Phys. Chem. A, 2015, 119, 3134.
- 7 I. Szabó and G. Czakó, J. Phys. Chem. A, 2017, 121, 9005.
- 8 Y.-T. Ma, X. Ma, A. Li, H. Guo, L. Yang, J. Zhang and W. L. Hase, *Phys. Chem. Chem. Phys.*, 2017, **19**, 20127.
- 9 J. Mikosch, S. Trippel, C. Eichhorn, R. Otto, U. Louderaj, J.-X. Zhang, W. L. Hase, M. Weidemüller and R. Wester, *Science*, 2008, **319**, 183.
- 10 J. Zhang, J. Mikosch, S. Trippel, R. Otto, M. Weidemüller, R. Wester and W. L. Hase, *J. Phys. Chem. Lett.*, 2010, 1, 2747.
- 11 J. Mikosch, J. Zhang, S. Trippel, C. Eichhorn, R. Otto, R. Sun, W. A. de Jong, M. Weidemüller, W. L. Hase and R. Wester, *J. Am. Chem. Soc.*, 2013, 135, 4250.
- 12 J. Xie and W. L. Hase, Science, 2016, 352, 32.
- 13 M. Stei, E. Carrascosa, M. A. Kainz, A. H. Kelkar, J. Meyer, I. Szabó, G. Czakó and R. Wester, *Nat. Chem.*, 2016, **8**, 151.
- 14 I. Szabó, B. Olasz and G. Czakó, J. Phys. Chem. Lett., 2017, 8, 2917.

- 15 S. A. Jarrett-Sprague and I. H. Hillier, J. Chem. Soc., Faraday Trans., 1990, 86, 3991.
- 16 M. Novak, K. A. Martin and J. L. Heinrich, *J. Org. Chem.*, 1989, 54, 5430.
- 17 J. S. Helmick, K. A. Martin, J. L. Heinrich and M. Novak, *J. Am. Chem. Soc.*, 1991, **113**, 3459.
- 18 R. Ulbrich, M. Famulok, F. Bosold and G. Boche, *Tetrahedron Lett.*, 1990, **31**, 1689.
- 19 P. Beak and J. Li, J. Am. Chem. Soc., 1991, 113, 2796.
- 20 R. Gareyev, S. Kato and V. M. Bierbaum, J. Am. Soc. Mass Spectrom., 2001, 12, 139.
- 21 M. Bühl and H. F. Schaefer III, J. Am. Chem. Soc., 1993, 115, 9143.
- 22 M. Bühl and H. F. Schaefer III, J. Am. Chem. Soc., 1993, 115, 364.
- 23 M. N. Glukhovtsev, A. Pross and L. Radom, *J. Am. Chem. Soc.*, 1995, **117**, 9012.
- 24 Y. Ren and H. Zhu, J. Am. Soc. Mass Spectrom., 2004, 15, 673.
- 25 J. Yang, Y. Ren, H. Zhu and S.-Y. Chu, *Int. J. Mass Spectrom.*, 2003, **229**, 199.
- 26 Y.-M. Xing, X.-F. Xu, Z.-S. Cai and X.-Z. Zhao, *J. Mol. Struct.: THEOCHEM*, 2004, **671**, 27.
- 27 X. Liu, J. Zhang, L. Yang and R. Sun, *J. Phys. Chem. A*, 2016, **120**, 3740.
- 28 J. Kubelka and F. M. Bickelhaupt, *J. Phys. Chem. A*, 2017, **121**, 885.
- 29 T. A. Hamlin, M. Swart and F. M. Bickelhaupt, ChemPhysChem, 2018, 19, 1315.
- 30 B. Hajdu and G. Czakó, J. Phys. Chem. A, 2018, 122, 1886.
- 31 J. Lv, J. Zhang and D. Wang, Phys. Chem. Chem. Phys., 2016, 18, 6146.
- 32 X. Niu, P. Liu and D. Wang, J. Phys. Chem. A, 2020, 124, 141.

- 33 F. Yu, L. Song and X. Zhou, *Comput. Theor. Chem.*, 2011, 977, 86.
- 34 F. Yu, J. Comput. Chem., 2012, 33, 401.
- 35 X. Liu, C. Zhao, L. Yang, J. Zhang and R. Sun, *Phys. Chem. Chem. Phys.*, 2017, **19**, 22691.
- 36 Y. Li and D. Wang, Phys. Chem. Chem. Phys., 2018, 20, 12106.
- 37 C. Møller and M. S. Plesset, Phys. Rev., 1934, 46, 618.
- 38 T. H. Dunning Jr, J. Chem. Phys., 1989, 90, 1007.
- 39 T. B. Adler, G. Knizia and H.-J. Werner, J. Chem. Phys., 2007, 127, 221106.
- 40 T. Győri and G. Czakó, J. Chem. Theory Comput., 2020, 16, 51.
- 41 H.-J. Werner, P. J. Knowles, G. Knizia, F. R. Manby and M. Schützet al., Molpro, version 2015.1, a package of ab initio programs, see http://www.molpro.net.
- 42 B. J. Braams and J. M. Bowman, *Int. Rev. Phys. Chem.*, 2009, 28, 577.
- 43 J. M. Bowman, G. Czakó and B. Fu, *Phys. Chem. Chem. Phys.*, 2011, **13**, 8094.
- 44 W. L. Hase, *Encyclopedia of Computational Chemistry*, Wiley, New York, 1998, pp. 399–407.
- 45 G. Czakó, B. C. Shepler, B. J. Braams and J. M. Bowman, *J. Chem. Phys.*, 2009, **130**, 084301.
- 46 D. Papp, V. Tajti, T. Győri and G. Czakó, *J. Phys. Chem. Lett.*, 2020, **11**, 4762.
- 47 P. Papp, V. Tajti and G. Czakó, *Chem. Phys. Lett.*, 2020, 755, 137780.
- 48 I. Szabó, A. G. Császár and G. Czakó, *Chem. Sci.*, 2013, 4, 4362.
- 49 J. C. Polanyi, Science, 1987, 236, 680.
- 50 D. Townsend, S. A. Lahankar, S. K. Lee, S. D. Chambreau, A. G. Suits, X. Zhang, J. Rheinecker, L. B. Harding and J. M. Bowman, *Science*, 2004, **306**, 1158.