



Cite this: *Chem. Sci.*, 2021, **12**, 10191

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 28th April 2021  
Accepted 18th June 2021

DOI: 10.1039/d1sc02360h  
[rsc.li/chemical-science](http://rsc.li/chemical-science)


## Introduction

Epoxides are versatile intermediates for the synthesis of countless irreplaceable compounds.<sup>1</sup> Since the introduction of various versatile metal-catalyzed epoxidations pioneered by Sharpless, Jacobsen, Katsuki and others,<sup>2</sup> alkene epoxidation has become one of the most studied reactions in organic synthesis, both in industry and academia.<sup>2</sup> After the successful developments employing metal catalysts, much effort has been devoted to the development of protocols where a more sustainable metal-free catalyst is employed. The organocatalytic epoxidation has flourished<sup>1a,1b,2</sup> in particular with ketone,<sup>2</sup> acid,<sup>3</sup> nitrile<sup>4</sup> or iminium salt<sup>2</sup> catalysts (Scheme 1, A). Pioneered by Adam,<sup>5</sup> Curci,<sup>6</sup> Yang,<sup>7</sup> Denmark,<sup>8</sup> Shi,<sup>9</sup> Miller,<sup>10</sup> Page<sup>11</sup> and others, a variety of oxidants, such as Oxone, *tert*-butyl hydroperoxide (TBHP) or less commonly hydrogen peroxide ( $H_2O_2$ ) were utilized to *in situ* generate the key dioxirane, peracid, imidoperoxoic acid, or oxaziridine intermediates.


Over the last few years, the epoxidation of alkenes with environmentally friendly aqueous  $H_2O_2$ <sup>12</sup> as the oxidant is drawing increasing attention, since it is inexpensive, relatively safe and affords water as byproduct. However, hydrogen peroxide by itself is a poor oxidant for organic oxidations<sup>2,13</sup> and thus has to be activated by a catalyst to form a reactive intermediate in order to enable the epoxidation of unactivated

alkenes. As one of the most common functionalities in organic chemistry, a wide range of aldehydes are readily available and have been employed as catalysts.<sup>14</sup> However, one of the most interesting characteristics is its auto-oxidation, where the aldehyde reacts with molecular oxygen, leading to a variety of activated intermediates.<sup>15</sup> For instance, in the versatile Mukaiyama epoxidation, overstoichiometric amounts of aldehydes have been employed as the mediators of the reaction (Scheme 1, B).<sup>16–18</sup> As a result, a particularly mild and selective


### A Organocatalytic Epoxidations of Non-Activated Alkenes (known)



### B Mukaiyama Epoxidation



### C Aldehyde-catalyzed Epoxidation of Non-Activated Alkenes (this work)



<sup>a</sup>Laboratory of Organic Chemistry, Department of Chemistry, National and Kapodistrian University of Athens, Panepistimiopolis 15771, Athens, Greece. E-mail: [cokotos@chem.uoa.gr](mailto:cokotos@chem.uoa.gr)

<sup>b</sup>Department of Chemistry, University of Basel, St. Johanns-Ring 19, Basel 4056, Switzerland. E-mail: [christof.sparr@unibas.ch](mailto:christof.sparr@unibas.ch)

† Electronic supplementary information (ESI) available: Experimental data, optimisation studies, mechanistic and HRMS studies. See DOI: [10.1039/d1sc02360h](https://doi.org/10.1039/d1sc02360h)

Scheme 1 Established epoxidations and conceptualization.




process allowed the epoxidation of an exceptionally broad range of alkenes at room temperature employing molecular oxygen as the terminal oxidant. The process can thereby be performed with<sup>16</sup> or without transition metal catalysts<sup>17</sup> or in the presence of *N*-hydroxyphthalimide as the catalyst.<sup>18</sup> The mechanisms of these epoxidations were studied in great detail. Nonetheless, the nature of the active oxygen species remains a subject of debate.<sup>16</sup> Considering the broad utility of the Mukaiyama epoxidation and the ideal attributes of aqueous H<sub>2</sub>O<sub>2</sub> as oxidant for sustainable synthesis, we anticipated the development of an aldehyde-catalyzed epoxidation of unactivated alkenes by accommodating the formyl functionality within a spatially well-defined site of an atropisomeric multiaxis system. We describe herein, that the atropisomeric two-axis aldehyde catalyst **3a** that was identified as an efficient catalyst for the activation of aqueous hydrogen peroxide, allows the synthesis of a broad range of epoxides **2** from a variety of unactivated alkenes **1** with notable turnover (Scheme 1, C).

## Results and discussion

We initiated our epoxidation studies using several atropisomeric two-, three- and four-axis systems **3a–3e**, which were recently prepared.<sup>19</sup> To our delight, 5 mol% of catalyst **3a** provided excellent activity for the epoxidation of  $\alpha$ -methylstyrene (**1a**) using H<sub>2</sub>O<sub>2</sub> as the oxidant (Scheme 2), which is in stark contrast to literature precedents where overstoichiometric amounts of aldehyde were required.<sup>16–18</sup> We next compared this remarkable finding for the epoxidation of unactivated alkenes with the other atropisomeric aldehydes comprising two to four stereogenic axes, as well as commercially available aldehydes

under otherwise identical conditions (2 eq. H<sub>2</sub>O<sub>2</sub> : MeCN in *t*-BuOH/aq. 0.6 M K<sub>2</sub>CO<sub>3</sub>; 4  $\times$  10<sup>-5</sup> M EDTA tetrasodium salt, pH = 11). When aldehyde **3a** was employed as the catalyst, the desired epoxide **2a** was formed in 74% yield. Interestingly, the diastereomeric aldehyde **3b** provided epoxide **2a** in significantly lower yield (34%), underscoring the prerequisite spatial positioning of aryl and bromide moieties to augment catalyst turnover. Furthermore, the diastereomeric catalysts **3c** and **3d** with three stereogenic axes as well as the atropisomeric four-axis system **3e** also gave considerably lower yields. While the relative configuration of **3a** notably impacts activity conclusively by governing the conformation of catalytic intermediates, the enantioselectivity remained low for all cases ( $\leq 10\%$  ee). The catalytic activity trends were also confirmed with the commercially available aldehydes without rotationally restricted axes (**3f–i**). Interestingly, an enhanced yield was also observed for *ortho*-bromonaphthaldehyde (**3i**) as compared to parent 2-naphthaldehyde (**3h**), supporting the notion that a bromide in proximity to the active site is beneficial for the reaction outcome. Nonetheless, the atropisomeric two-axis aldehyde **3a** outperformed all other catalysts, highlighting the impact of bromine and aryl groups in proximity to the active site to improve the catalytic activity and turnover.

Having identified the unique characteristics of catalyst **3a**, we focused on optimizing the reaction conditions (Table 1). With 5 mol% of **3a** in the presence of 6 eq. H<sub>2</sub>O<sub>2</sub> and aqueous buffer in *t*-BuOH, the desired  $\alpha$ -methyl styrene oxide (**2a**) was formed in 98% yield (entry 2). Furthermore, control experiments without catalyst, oxidant, buffer or MeCN confirmed that all components are necessary for the epoxidation (Table 1, entry 3–6). In agreement with our previous findings,<sup>20a,20b</sup> a lower pH provided the desired product in compromised yield, validating



Scheme 2 Survey of aldehyde catalysts. <sup>a</sup>K<sub>2</sub>CO<sub>3</sub>/EDTA buffer (pH = 11). The yields were determined by <sup>1</sup>H NMR using an internal standard.

Table 1 Optimization of the reaction conditions<sup>a</sup>

Entry	3a [mol%]	Solvent	Deviation	Yield (%)
1	5.0	<i>t</i> -BuOH	—	74%
2	5.0	<i>t</i> -BuOH	6 eq. H <sub>2</sub> O <sub>2</sub>	98%
3	0	<i>t</i> -BuOH	—	9%
4	5.0	<i>t</i> -BuOH	No H <sub>2</sub> O <sub>2</sub>	0%
5	5.0	<i>t</i> -BuOH	No buffer	0%
6	5.0	<i>t</i> -BuOH	No MeCN	0%
7	5.0	<i>t</i> -BuOH	pH = 10	41%
8	5.0	EtOAc	—	67%
9	5.0	MeCN	—	82%
10	5.0	CH <sub>2</sub> Cl <sub>2</sub>	—	15%
11	5.0	CHCl <sub>3</sub>	—	12%
12	5.0	MeOH	—	quant.
13	1.0	MeOH	—	83%

<sup>a</sup> K<sub>2</sub>CO<sub>3</sub>/EDTA buffer (pH = 11). Yield determined by <sup>1</sup>H NMR using an internal standard.




that  $\text{pH} = 11$  is ideal for the generation of Payne's intermediate<sup>20b,21</sup> (peroxycarboximidic acid **V** from MeCN and  $\text{H}_2\text{O}_2$ , Table 1, entry 7 and mechanism in Scheme 5). The importance of the generation of Payne's intermediate for the success of our protocol is also highlighted by the fact that in the absence of MeCN, no epoxidation is taking place (Table 1, entry 6). However, Payne's intermediate alone cannot promote the reaction to completion as can be extracted by the results of Table 1, entry 3. Using different solvents (entries 8–12), methanol was determined as the most effective medium for the reaction, forming the desired  $\alpha$ -methyl styrene oxide in quantitative yield (entry 12). On the other hand, a decrease of the


catalyst loading to 1 mol% led to a reduced yield of 83% (entry 13). Furthermore, after performing the epoxidation (Table 1, entry 12), the catalyst **3a** was recovered intact (94% recovery by  $^1\text{H-NMR}$ , 82% recovery after column chromatography). To our knowledge, this is the first example, where an aldehyde is employed in substoichiometric amounts as the catalyst for an epoxidation reaction (5 mol% *versus* usually 10–50 mol% of an activated ketone in literature<sup>7–10</sup> or 5–10 equivalents of pivaldehyde<sup>16–18</sup>).

With the optimized reaction conditions defined, we set out to explore the substrate scope of the aldehyde-catalysed epoxidation of unactivated alkenes (Scheme 3). A variety of mono-substituted styrenes were first tested, providing excellent isolated yields for the unfunctionalized and methoxy-substituted styrenes (**2b** and **2c**), but low yields for **2d** and **2e** (in the case of **2e**, 28% of the corresponding diol was also isolated). In contrast to known procedures which lead to low amounts of epoxide when terminal aliphatic alkenes are employed, synthetically meaningful yields were obtained for corresponding epoxides **2f** and **2g**. Gratifyingly, when studying disubstituted alkenes, geminal and vicinal disubstituted olefins afforded an excellent outcome [**2a**, **2h** (99 : 1 *trans* : *cis* from 99 : 1 *trans* : *cis* **1h**) and **2i** (only *trans* from only *trans* **2i**)] and also cyclic disubstituted alkenes provided epoxides in up to 97% isolated yield (**2j**–**2m**). To explore the limitations of the method, also trisubstituted unactivated alkene substrates were investigated, leading to a 68% yield for a double epoxidation to product **2n** and an inferior performance for aryl-substituted **2o**. Unfortunately, tetrasubstituted alkenes or linear *cis* alkenes are not viable substrates for this method, as in the former case, the desired epoxide is not formed, while in the latter case, low yields (around 16%) are observed.

Due to the novelty of aldehyde catalysis for epoxidations with hydrogen peroxide, mechanistic studies to probe the reaction pathway of this epoxidation were carried out (Scheme 4). In agreement with analogous studies on the auto-oxidation of aldehydes,<sup>15a,15b,15d,16i,18b</sup> control experiments for the epoxidation of **1a** to **2a** support a mechanism involving an auto-oxidation of aldehyde **3a**. When the reaction was performed under an argon atmosphere (Scheme 4), only 11% product formation was observed, emphasizing the involvement of oxygen in the reaction in agreement with the epoxidation occurring through the Payne intermediate (see Table 1, entry 3). In contrast, almost quantitative yields were obtained by performing the reaction under open air or a balloon of oxygen, verifying the notion that oxygen is a constituent of the mechanism. According to literature,<sup>15a,15b,15d,16i,18b</sup> oxygen is required for the generation of the Criegee intermediate **IV**.<sup>22</sup> Indeed, adding 2 mol% of *m*CPBA in the mechanistic experiment that was performed in the absence of oxygen, the epoxidation reaction can be restored to a good extent (42% yield, see below). Furthermore, if TEMPO or BHT were added to the reaction as a radical trap, no epoxidation product was formed, verifying the generation of radical intermediates in the autoxidation of the catalyst. When the reaction was performed in the dark, the yield of the desired epoxide **2a** remained constant, proving that the reaction mechanism does not follow a photochemical pathway. In order to exclude the




Scheme 3 Substrate Scope. <sup>a</sup> $\text{K}_2\text{CO}_3$ /EDTA buffer (pH = 11). Yields of products after isolation by column chromatography.



Scheme 4 Mechanistic and radical clock experiments.





Scheme 5 Proposed mechanism.

possibility that the epoxidation is promoted by an *in situ* generated peracid, we substituted catalyst **3a** by its acid or alternatively by PhCOOH and observed a very low yield, which again can be assigned to the reactivity of the Payne intermediate. This is in agreement with literature,<sup>10</sup> where an acid in combination with MeCN/H<sub>2</sub>O<sub>2</sub> cannot promote the epoxidation reaction. These results in combination with the almost quantitative recovery of **3a** from the reaction mixture, verify that a peracid is not responsible for the epoxidation, since in that case, **3a** could not have been recovered intact and should have been oxidized to the corresponding acid,<sup>23</sup> which is not capable of performing the epoxidation. In order to test the nature of the active intermediate that enables the epoxidation of the unactivated alkenes, we performed the epoxidation with the radical clock **1p**. Epoxide **2p** was observed in 19% yield, along with 19% of diol **2p'** and 62% of unreacted **1p**, without any ring-opened product detected. This finding supports that the active oxidant is of non-radical nature, and that the radical steps during autoxidation of the aldehyde precede the epoxidation step.

A proposed mechanism consistent with the findings consequently involves the initial autoxidation of a fraction of the aldehyde **3a** under aerobic conditions, forming radical intermediate **I** (Scheme 5).<sup>15a,15b,15d,16i,18b</sup> The peroxy-radical intermediate **II** and peracid **III** is subsequently formed with oxygen.<sup>15</sup> The addition of peracid **III** to aldehyde **3a** provides the Criegee intermediate **IV**.<sup>15b,22</sup> At pH = 11, the Payne intermediate **V** is generated,<sup>4,20b,21</sup> which is in equilibrium with **VI** from a reaction with peracid **III**. The Criegee and Payne intermediates (**IV** + **V**) result in a postulated species **VII**, which breaks up to form the key dioxirane intermediate **VIII**, the acetamide byproduct and regenerates peracid **III**. Another plausible pathway is the

transformation of intermediate **IV** and **V**, to form **III** and **VII'**, which can also give rise to dioxirane **VIII** and acetamide. The dioxirane intermediate **VIII** as the active oxidant then reacts with unactivated alkene **1a** to form epoxide **2a**, regenerating aldehyde **3a** to close the catalytic cycle. The crucial role of the buffer for the generation of intermediate **V** is highlighted in the control experiments, as well as the inability of **V** alone to push the reaction to completion. Oxygen from air is required for the autoxidation of the aldehyde to form intermediate **IV**. A small amount of peracid is necessary to promote the catalytic cycle *via* the generation of **IV** and not to perform the epoxidation, as highlighted by the recovery of **3a** after reaction completion and the reinstatement of the epoxidation yield when catalytic *m*CPBA was added to the control experiment under argon. A finding consistent with the formation of dioxirane **VIII** is that **3a** under Shi-like conditions with Oxone as the oxidant, simulating the formation of the dioxirane-intermediate, provided the desired epoxide **2a** in good yield (65%).<sup>24</sup>

Considering this mechanistic hypothesis, we examined the possible pathways of the reaction by High Resolution Mass Spectrometry (HRMS, see ESI† for details) and thereby detected several of the intermediates of the proposed pathway. Adding TEMPO as the radical scavenger to the reaction mixture, the adducts of **I** and **II** (trapped with TEMPO) were identified, supporting that the autoxidation step is occurring. The Criegee intermediate **IV** and both intermediates **V** and **VI** were also detected, corroborating the formation of the Payne intermediate, when using the appropriate aqueous buffer (pH = 11). Moreover, also the central alkene oxidant dioxirane **VIII** was detected by HRMS. In order to rule out artifacts, the measurements were also repeated using CD<sub>3</sub>CN or 2-bromobenzaldehyde (**3g**) to detect similar intermediates, such as the corresponding Criegee intermediate and the dioxirane by HRMS, thus further underpinning the proposed mechanism.<sup>24</sup> Intermediate **VII** was observed in the case of **3g** with CD<sub>3</sub>CN, while **VII'** was detected with **3a** in the presence of TEMPO. In accord with these findings, both short-lived intermediates provide exploratory support of an aldehyde-catalyzed epoxidation by means of dioxirane formation.

## Conclusions

In conclusion, we have developed a sustainable and efficient method for the epoxidation of unactivated alkenes utilizing an atropisomeric multiaxis aldehyde as the catalyst in combination with hydrogen peroxide as the oxidant. This study reveals the feasibility of aldehyde-catalyzed epoxidations and the mechanistic studies shed first light on this unique reactivity. To our knowledge this is the first example of an epoxidation reaction that employs an aldehyde in catalytic amounts. Ongoing studies aim to rationalize the individual requirements for catalyst design to further increase catalytic efficiency and to accomplish stereoselectivity.

## Data availability

Experimental data are available from the authors upon request.



## Author contributions

I. T., C. S. and C. G. K. conceived the study, designed the experiments, and analyzed the data. I. T. performed the experiments and M. G. K. the HRMS studies. D. L. provided catalysts 3b-3e. The manuscript was written through contributions of all authors.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

C. G. K., I. T. and M. G. K. gratefully acknowledge the Hellenic Foundation for Research and Innovation (HFRI) for financial support through a grant, which is financed by 1<sup>st</sup> Call for H. F. R. I. Research Projects to Support Faculty Members & Researchers and the procurement of high-cost research equipment grant (grant number 655). I. T. and M. G. K. would like to thank State Scholarships Foundation (IKY) for financial support through a post-doctoral research scholarship funded by the “Supporting Post-Doctoral Researchers-Call B” (MIS 5033021), action of the Operational Programme “Human Resources Development, Education and Lifelong Learning” and co-funded by the European Social Fund (ESF) and Greek National Resources. The COST Action C-H Activation in Organic Synthesis CA15106 is acknowledged for supporting the Short-Term Scientific Mission of I. T. to the University of Basel and Dr D. Kaldre for helpful discussions. The Swiss National Science Foundation is gratefully acknowledged for financial support (C. S. and D. L.).

## Notes and references

- For selected reviews, see: (a) O. A. Wong and Y. Shi, *Chem. Rev.*, 2008, **108**, 3958; (b) Y. Zhu, Q. Wang, R. G. Cornwall and Y. Shi, *Chem. Rev.*, 2014, **114**, 8199; For books, see: (c) A. K. Yudin, *Aziridines and Epoxides in Organic Synthesis*, Wiley-VCH, Weinheim, 2006; (d) G. Centi, S. Perathoner and S. Abate, *Modern Heterogeneous Oxidation Catalysis: Design, Reactions and Characterization*, Wiley-VCH, Weinheim, 2009.
- For a recent review, see: I. Triandafyllidi, D. I. Tzaras and C. G. Kokotos, *ChemCatChem*, 2018, **10**, 2521.
- For the initial introduction of *m*-CBPA: N. Prileschajew, *Ber. Dtsch. Chem. Ges.*, 1909, **42**, 4811.
- G. B. Payne, P. H. Deming and P. H. Williams, *J. Org. Chem.*, 1961, **26**, 659.
- For selected examples, see: (a) W. Adam, R. Curci and J. O. Edwards, *Acc. Chem. Res.*, 1989, **22**, 205; (b) W. Adam, P. B. Rao, H.-G. Degen, A. Levai, T. Patonay and C. R. Sahamoller, *J. Org. Chem.*, 2002, **67**, 259.
- For selected examples, see: R. Mello, M. Fiorentino, O. Sciacovelli and R. Curci, *J. Org. Chem.*, 1988, **53**, 3890.
- For selected examples, see: (a) D. Yang, Y.-C. Yip, M.-W. Tang, M.-K. Wong, J.-H. Zheng and K.-K. Cheng, *J. Am. Chem. Soc.*, 1996, **118**, 491; (b) D. Yang, M.-K. Wong, Y.-C. Yip, X.-C. Wang, M.-W. Tang, J.-H. Zheng and K.-K. Cheng, *J. Am. Chem. Soc.*, 1998, **120**, 5943.
- For selected examples, see: (a) S. E. Denmark, Z. Wu, C. M. Crudden and H. Matsuhashi, *J. Org. Chem.*, 1997, **62**, 8288; (b) S. E. Denmark and H. Matsuhashi, *J. Org. Chem.*, 2002, **67**, 3479.
- For selected examples, see: (a) Y. Tu, Z.-X. Wang and Y. Shi, *J. Am. Chem. Soc.*, 1996, **118**, 9806; (b) Z.-X. Wang, Y. Tu, M. Frohn, J.-R. Zhang and Y. Shi, *J. Am. Chem. Soc.*, 1997, **119**, 11224; (c) C. P. Burke and Y. Shi, *J. Org. Chem.*, 2007, **72**, 4093; (d) N. Nieto, I. J. Munslow, H. Fernandez-Perez and A. Vidal-Ferran, *Synlett*, 2008, **18**, 2856.
- For selected examples, see: (a) G. Peris, C. E. Jakobsche and S. J. Miller, *J. Am. Chem. Soc.*, 2007, **129**, 8710; (b) C. E. Jakobsche, G. Peris and S. J. Miller, *Angew. Chem., Int. Ed.*, 2008, **47**, 6707; (c) P. A. Lichitor and S. J. Miller, *Nat. Chem.*, 2012, **4**, 990.
- For selected examples, see: (a) P. C. B. Page, G. A. Rassias, D. Bethell and M. B. Schilling, *J. Org. Chem.*, 1998, **63**, 2774; (b) P. C. B. Page, M. M. Farah, B. R. Buckley and A. J. Blacker, *J. Org. Chem.*, 2007, **72**, 4424.
- The environmental impact of H<sub>2</sub>O<sub>2</sub> is a matter of debate: (a) *Hydrogen Peroxide, Toxicological Overview*, Public Health England, 2009, [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/337708/Hydrogen\\_Peroxide\\_Toxicological\\_Overview\\_phe\\_v1.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/337708/Hydrogen_Peroxide_Toxicological_Overview_phe_v1.pdf); (b) T. Mahaseth and A. Kuzminov, *Mutat. Res.*, 2017, **773**, 274; (c) S. Fukuzumi, Y. -M. Lee and W. Nam, *Chin. J. Catal.*, 2021, **42**, 1241; (d) Y. Xue, Y. Wang, Z. Pan and K. Sayama, *Angew. Chem., Int. Ed.*, 2021, **60**, 10469.
- (a) G. Goor, J. Glenneberg and S. Jacobi, *Hydrogen Peroxide*, Wiley-VCH, Weinheim, 2012. For reviews see: (b) G. Grigoropoulou, J. H. Clark and J. A. Elings, *Green Chem.*, 2003, **5**, 1; (c) G. De Faveri, G. Ilyashenko and M. Watkinson, *Chem. Soc. Rev.*, 2011, **40**, 1722.
- For a recent review, see: Q. Wang, Q. Gu and S. You, *Angew. Chem., Int. Ed.*, 2019, **58**, 6818.
- For selected examples, see: (a) J. R. McNesby and C. A. Heller Jr, *Chem. Rev.*, 1954, **54**, 325; (b) L. Vanoye, A. Favre-Réguillon, A. Aloui, R. Philippe and C. de Bellefon, *RSC Adv.*, 2013, **3**, 18931; (c) K. Miyamoto, J. Yamashita, S. Narita, Y. Sakai, K. Hirano, T. Saito, C. Wang, M. Ochiai and M. Uchiyama, *Chem. Commun.*, 2017, **53**, 9781; (d) A. Maity, S.-M. Hyun and D. C. Powers, *Nat. Chem.*, 2018, **10**, 200.
- (a) T. Yamada, T. Takai, O. Rhode and T. Mukaiyama, *Chem. Lett.*, 1991, **20**, 1; (b) T. Yamada, T. Takai, O. Rhode and T. Mukaiyama, *Bull. Chem. Soc. Jpn.*, 1991, **64**, 2109; (c) W. W. Nam, H. J. Kim, S. H. Kim, R. Y. N. Ho and J. S. Valentine, *Inorg. Chem.*, 1996, **35**, 1045; (d) B. B. Wentzel, P. A. Gosling, M. C. Feiters and R. J. M. Nolte, *J. Chem. Soc., Dalton Trans.*, 1998, 2241; (e) B. B. Wentzel, P. L. Alsters, M. C. Feiters and R. J. M. Nolte, *J. Org. Chem.*, 2004, **69**, 3453; (f) A. C. Serra and A. Gonsalves, *Tetrahedron Lett.*, 2011, **52**, 3489; (g)



Y. Li, X. T. Zhou, S. Y. Chen, R. C. Luo, J. Jiang, Z. X. Liang and H. B. Ji, *RSC Adv.*, 2015, **5**, 30014; (h) J. Wang, M. Yang, W. Dong, Z. Jin, J. Tang, S. Fan, Y. Lu and G. Wang, *Catal. Sci. Technol.*, 2016, **6**, 161; (i) W. Zhou, J. Zhou, Y. Chen, A. Cui, F. Sun, M. He, Z. Xu and Q. Chen, *Appl. Catal., A*, 2017, **542**, 191.

17 (a) K. Kaneda, S. Haruna, T. Imanaka, M. Hamamoto, Y. Nishiyama and Y. Ishii, *Tetrahedron Lett.*, 1992, **33**, 6827; (b) K. R. Lassila, F. J. Waller, S. E. Werkheiser and A. L. Wressell, *Tetrahedron Lett.*, 1994, **35**, 8077; (c) C. Lehtinen and G. Brunow, *Org. Process Res. Dev.*, 1999, **3**, 101; (d) F. Loeker and W. Leitner, *Chem.-Eur. J.*, 2000, **6**, 2011.

18 (a) F. Minisci, C. Gambarotti, M. Pierini, O. Porta, C. Punta, F. Recupero, M. Lucarini and V. Mugnaini, *Tetrahedron Lett.*, 2006, **47**, 1421; (b) R. Spaccini, L. Liguori, C. Punta and H. R. Bjorsvik, *ChemSusChem*, 2012, **5**, 261.

19 D. Lotter, A. Castrogiovanni, M. Neuburger and C. Sparr, *ACS Cent. Sci.*, 2018, **4**, 656.

20 (a) D. Limnios and C. G. Kokotos, *J. Org. Chem.*, 2014, **79**, 4270; (b) E. Voutyritsa, A. Theodorou, M. G. Kokotou and C. G. Kokotos, *Green Chem.*, 2017, **19**, 1291; (c) D. Limnios and C. G. Kokotos, *ACS Catal.*, 2013, **3**, 2239; For examples on the use of trifluoromethyl ketones for oxidation reaction from other groups, see: (d) Q.-Q. Cheng, Z. Zhou, H. Jiang, J. H. Siitonens, D. H. Ess, X. Zhang and L. Kürti, *Nat. Catal.*, 2020, **3**, 386; (e) M. Cai, K. Xu, Y. Li, Z. Nie, L. Zhang and S. Luo, *J. Am. Chem. Soc.*, 2021, **143**, 1078.

21 (a) G. B. Payne and P. H. Williams, *J. Org. Chem.*, 1961, **26**, 651; (b) L. A. Arias, S. Adkins, C. J. Nagel and R. D. Bach, *J. Org. Chem.*, 1983, **48**, 888; (c) S. Ye, J. J. Hu and D. Yang, *Angew. Chem., Int. Ed.*, 2018, **57**, 10173.

22 (a) J. Criegee, *Justus Liebigs Ann. Chem.*, 1948, **560**, 127. This Criegee intermediate is not to be confused with the Criegee zwitterion formed in alkene ozonolysis. (b) Z. Hassan, M. Stahlberger, N. Rosenbaum and S. Bräse, *Angew. Chem., Int. Ed.*, 2021, DOI: 10.1002/anie.202014974.

23 (a) N. Shapiro and A. Vigalok, *Angew. Chem., Int. Ed.*, 2008, **47**, 2849; (b) Z. E. Hamami, L. Vanoye, P. Fongarland, C. de Bellefon and A. Favre-Reguillon, *J. Flow Chem.*, 2016, **6**, 206.

24 For detailed mechanistic studies and results, see ESI.†

