



Cite this: *Chem. Sci.*, 2021, **12**, 11756

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 27th June 2021
 Accepted 31st July 2021

DOI: 10.1039/d1sc03487a
rsc.li/chemical-science

Introduction


Silacycles have attracted increasing attention because they have shown unique physical, optoelectronic, and physiological properties in medicinal chemistry and materials science.¹ In this context, considerable efforts have been devoted to the development of synthetic methods for silacycles, which is a prerequisite for fully discovering their application potentials.^{2,3} Among them, of particular interest to synthetic chemists is the synthesis of silacycles *via* the cleavage of C-Si bonds.³ Typical strategies include direct annulation of silicon-based frameworks^{3a–n} and C-Si/C-C bond exchange reaction of small ring systems based on the existence of a high ring strain.^{3o–r} However, these methods generally require transition-metal catalysts to assist the activation of C-Si bonds. Transition-metal-free catalyzed annulation to assemble silacycles by cleaving C-Si bonds is still elusive so far.

The Brook rearrangement enables an intramolecular migration of a silyl group from carbon to oxygen atoms *via* a hypervalent silicon species,⁴ which was initially introduced by Brook⁵ and was demonstrated to be a reversible process.⁶ Its reverse process, namely retro-Brook rearrangement, can, in turn, be achieved by the transfer of a silyl group from oxygen to carbon atoms (Scheme 1A).⁷ Obviously, the Brook and retro-

Me₃SiSiMe₂(OⁿBu): a disilane reagent for the synthesis of diverse silacycles *via* Brook- and retro-Brook-type rearrangement†

Yankun Xu, Weiwei Xu, Xinyang Chen, Xiai Luo, Haiyan Lu, Minghao Zhang, Xiumei Yang, Guobo Deng, Yun Liang * and Yuan Yang *

Herein, a readily available disilane Me₃SiSiMe₂(OⁿBu) has been developed for the synthesis of diverse silacycles *via* Brook- and retro-Brook-type rearrangement. This protocol enables the incorporation of a silylene into different starting materials, including acrylamides, alkene-tethered 2-(2-iodophenyl)-1H-indoles, and 2-iodobiaryls, *via* the cleavage of Si-Si, Si-C, and Si-O bonds, leading to the formation of spirobenzosiloles, fused benzosiloles, and π -conjugated dibenzosiloles in moderate to good yields. Preliminary mechanistic studies indicate that this transformation is realized by successive palladium-catalyzed bis-silylation and Brook- and retro-Brook-type rearrangement of silane-tethered silanols.



National & Local Joint Engineering Laboratory for New Petro-Chemical Materials and Fine Utilization of Resources, Key Laboratory of Chemical Biology and Traditional Chinese Medicine Research, Ministry of Education, Key Laboratory of the Assembly and Application of Organic Functional Molecules of Hunan Province, Hunan Normal University, Changsha 410081, China. E-mail: yliang@hunnu.edu.cn; yuanyang@hunnu.edu.cn

† Electronic supplementary information (ESI) available. CCDC 2040657–2040659. For ESI and crystallographic data in CIF or other electronic format see DOI: [10.1039/d1sc03487a](https://doi.org/10.1039/d1sc03487a)

Scheme 1 (A) Brook and retro-Brook rearrangement. (B) The Known reaction modes of disilanes. (C) Our work: a disilane reagent Me₃SiSiMe₂(OⁿBu) for the synthesis of diverse silacycles.




Brook rearrangements allow the cleavage of a C-Si bond and the formation of a C-Si bond under transition-metal-free catalysis.⁸ On the other hand, the existing studies are limited to the translocation of a single silyl group from the starting materials. The migration of two different silyl groups in one event has not been reported so far. In this context, we want to explore the synthesis of silacycles *via* the migration of two different silyl groups in the Brook and retro-Brook rearrangement. The design and synthesis of silane-tethered silanols is undoubtedly the primary task and challenge to achieve this hypothesis.

Disilanes have been among the most versatile silylation reagents in organic synthesis. Over the past few decades, a myriad of methods for accessing organosilanes from disilanes have focused on the activation of Si-Si bonds for the development of mono-silylation involving aryl halides or cyanides, alkenes, and C-H bonds and bis-silylation of alkynes, alkenes, carbenes, and palladacycles (Scheme 1B).^{9–13} Particularly, the pioneering work that realizes the bis-silylation of *in situ* generated palladacycles with hexamethyldisilane *via* the cleavage of Si-Si bonds has recently been reported by Zhang, Cheng, and us.^{12,13} These advances have inspired us to modify hexamethyldisilane, namely the replacement of the methyl group with an oxygen-containing group, for the synthesis of silane-tethered silanols, which were further converted into silacycles *via* Brook and retro-Brook-type rearrangement. Herein, we disclose a Brook and retro-Brook-type rearrangement strategy for the synthesis of diverse silacycles, including spirobenzosiloles, fused benzosiloles, and π -conjugated dibenzosiloles, by employing a readily available disilane reagent $\text{Me}_3\text{SiSiMe}_2(\text{O}^{\prime}\text{Bu})$ that could be prepared by a simple treatment of pentamethylchlorodisilane with *n*-butanol in the presence of NEt_3 at room temperature (Scheme 1C). Notably, the cleavage of Si-Si, Si-C, and Si-O bonds is involved in the transformation.

Results and discussion


We initiated the studies by investigating the reaction of acrylamide **1a** with 1-butoxy-1,1,2,2,2-pentamethyldisilane **2a**. To our delight, the anticipated spirobenzosilole **3a** was indeed afforded in 66% yield by using a simple catalytic system composed of $\text{Pd}(\text{OAc})_2$ and K_2CO_3 in DMA at 90 °C. Encouraged by these initial results, various parameters were screened, and the optimized reaction conditions are as follows: **1a** (0.2 mmol),



Scheme 2 Screening of disilanes (2).

2a (0.3 mmol), $\text{Pd}(\text{OAc})_2$ (10 mol%), PPh_3 (20 mol%), and K_2CO_3 (0.6 mmol) in DMA (2 mL) at 90 °C under N_2 for 6 h (see the ESI† for details). Next, several disilanes **2b–h** were tested. As shown in Scheme 2, when the *n*-butyl group of disilane **2a** was replaced by other functional groups such as *n*-hexyl, benzyl, cyclohexyl, and 2-oxopropyl, all of them could afford the desired product **3a**, albeit in a lower yield. Unexpectedly, disiloxane **2g** was unreactive. Finally, disiloxane **2g** was found to produce product **3a** in 58% yield.

With the optimal reaction conditions and disilane reagent confirmed, the scope of acrylamides **1** was subsequently examined. Gratifyingly, this protocol was applicable to a large variety of acrylamides **1** to afford spiro[benzo[*b*]silole-3,3'-indolin]-2'-ones **3a–u** in moderate to good yields (Scheme 3). Note that the replacement of iodine atoms with bromine atoms on the acrylamide showed good reactivity, delivering the product **3a** in 66% yield. Moreover, the three substructures of acrylamides **1** were systematically investigated. Regarding different substituents on the nitrogen atom, methyl or ethyl group substituted acrylamides **1b** and **1c** were competent substrates, while acrylamide **1d** with a Ts group could not give the target product **3d** under the standard conditions. For the 2-iodoaniline fragment, a broad range of functional groups on the




Scheme 3 Variations of the acrylamides (1). ^aReaction conditions: **1** (0.2 mmol), **2a** (0.3 mmol), $\text{Pd}(\text{OAc})_2$ (10 mol%), PPh_3 (20 mol%), K_2CO_3 (3 equiv.), and DMA (2 mL) at 90 °C under a N_2 atmosphere for 6 h. ^b*N*-Benzyl-*N*-(2-bromophenyl)-2-phenylacrylamide. ^c**1a** (1 mmol). ^d70 °C and 1 h.




benzene ring, including electron-donating groups (Me and OMe), modifiable halogen groups (F and Cl), and even strong electron-withdrawing groups (CF_3 , CO_2Me , and NO_2), were well tolerated (**3e–n**). Meanwhile, the structure of **3g** was unambiguously confirmed by X-ray crystallography. Their electronic properties seem to affect the reactivity, since substrates **1l–n** with strong electron-withdrawing groups, especially CO_2Me and NO_2 groups, resulted in a diminished yield. Finally, the compatibility was further demonstrated by testing the key 2-phenylalkene moiety. Both the benzene ring containing *ortho* or *para* substituents and the naphthalene ring could survive, affording spirobenzosilole **3o–u** in moderate to good yields. Nevertheless, a slight modification of the reaction conditions was required when substrates bearing F groups were used (**3q** and **3t**). When the reaction was scaled up to 1 mmol, 70% spirocyclic product **3a** could also be obtained.

To highlight the generality of this domino Heck/silacyclization, we envision that fused benzosiloles can be synthesized by a domino Heck/*ortho* C–H functionalization of aryl iodides. Therefore, 2-(2-iodophenyl)-1-(2-methylallyl)-1*H*-indole was employed to react with disilane **2a** under the above reaction conditions. To our delight, indolo[2,1-*a*]silolo[4,3,2-*de*]isoquinolines **5a** could smoothly be produced in 58% yield by the cleavage of Si–Si and Si–O bonds. Encouraged by these results, the scope of 2-(2-halophenyl)-1-(2-methylallyl)-1*H*-indoles was then explored (Scheme 4). Bromine atoms instead of iodine atoms on the substrate **4a'** were subjected to the standard conditions, which could give product **5a**, albeit in a lower yield. Satisfactorily, substrates **4b–g** being diversely substituted (Me, F, and Cl) on the indole ring were able to undergo this domino Heck/silacyclization with disilane **2a** to provide the desired products **5b–g** in moderate yields. Differently, using substrate **4h** required relatively mild conditions.



Scheme 5 Variations of the 2-iodobiphenyls (**6**). ^aReaction conditions: **6** (0.2 mmol), **2a** (0.24 mmol), $\text{Pd}(\text{OAc})_2$ (10 mol%), K_2CO_3 (3 equiv.), and DMF (2 mL) at $90\text{ }^\circ\text{C}$ under a N_2 atmosphere for 12 h. ^b2-Bromobiphenyl was used. ^c2-Iodo-4'-methyl-1,1'-biphenyl was used. ^d2-Iodo-3'-methyl-1,1'-biphenyl was used.




Scheme 4 Variations of the 2-(2-iodophenyl)-1*H*-indoles chlorides (**4**). ^aReaction conditions: **4** (0.2 mmol), **2a** (0.3 mmol), $\text{Pd}(\text{OAc})_2$ (10 mol%), PPh_3 (20 mol%), K_2CO_3 (3 equiv.), and DMA (2 mL) at $90\text{ }^\circ\text{C}$ under a N_2 atmosphere for 6 h. ^b2-(2-Bromophenyl)-1-(2-methylallyl)-1*H*-indole. ^c $70\text{ }^\circ\text{C}$ and 1 h.

To emphasize the versatility of the disilane reagent $\text{Me}_3\text{-SiSiMe}_2(\text{O}^{\prime}\text{Bu})$, we next attempted to synthesize π -conjugated dibenz[*b,d*]siloles by performing the reaction of 2-iodo-1,1'-biphenyl with $\text{Me}_3\text{SiSiMe}_2(\text{O}^{\prime}\text{Bu})$ **2a**. However, no anticipated product was observed under the above standard conditions. Subsequently, PPh_3 was found to suppress the reaction, since dibenz[*b,d*]siloles **7a** could be afforded in 52% yield by the removal of PPh_3 . Encouraged by these results, the optimal reaction conditions that could furnish 64% of **7a** were established by the screening of various parameters (see the ESI† and Scheme 5). Afterward, a series of substituted 2-iodobiphenyls were examined. Delightfully, the electron-donating group (Me and OMe) on the benzene ring could be tolerated for the silacyclization reaction with disilane **2a**, thus delivering the desired products **7b–k** in moderate yields. Unfortunately, this protocol was not applicable to substrates **6l–o** bearing an electron-withdrawing group. The possible reason is that the protonation of palladacycles formed by substrates **6l–o** is easier than bis-silylation (see the ESI†).

To gain insight into the reaction mechanism, a range of control experiments were performed (Scheme 6). The reaction of acrylamides **1a** with disiloxane **2h** or **2i** could afford product **3a** in 27% and 50% yields, respectively, under the optimal reaction conditions (eqn (1)†). Moreover, hexamethyldisiloxane (TMSOTMS) was detected by gas chromatography in the model reaction of **1a** with **2a** (eqn (2), see the ESI†). These results indicated that the silicon source of product **3a** came from the dimethylsilyl group generated by disilane **2a** via the cleavage of Si–Si and Si–O bonds. Interestingly, two disilylated products **8a** and **8a'** as well as product **3a** were isolated in 43%, 27%, and 16% yields when **1a** and **2a** were reacted under the standard conditions for 20 minutes (eqn (3)†). Therefore, we speculated that two disilylated products **8a** and **8a'** were the reaction intermediates (the structure of **8a** was absolutely confirmed by X-ray crystallography). Finally, a spiropalladacycle¹⁴ that could






Scheme 6 Mechanistic experiments.


be prepared from acrylamides and stoichiometric $\text{Pd}(\text{PPh}_3)_4$ was employed for the reaction with disilane **2a** (eqn (4) \dagger). Unexpectedly, no product **3b** was observed in the absence of $\text{Pd}(\text{OAc})_2$ and PPh_3 . Surprisingly, extra addition of $\text{Pd}(\text{OAc})_2$ and PPh_3 could give product **3b** in 60% yield. These results suggested that the spiropalladacycle as a reaction intermediate underwent transmetalation with intermediate **G**, rather than direct oxidative addition with disilane **2a**, to furnish disilylated products **8a** and **8a'**, which were then converted into the desired product **3a**.

To verify our hypothesis and propose the possible formation process of **3a** from **8a** or **8a'**, we conducted several control experiments (Scheme 7). The silacyclization of disilylated products **8a** and **8a'**, respectively, was conducted in the absence

Scheme 7 Mechanistic experiments. The possible process for the transformation of **8a** and **8a'** into **3a**.

of $\text{Pd}(\text{OAc})_2$ and PPh_3 , and the product **3a** was obtained in 66% and 60% yields as expected (eqn (1) \dagger). Besides, TMSOTMS was also detected (see the ESI \dagger). These results demonstrated that the transformation of **8a** and **8a'** into **3a** involved the cleavage of the $\text{Me}_3\text{Si}-\text{C}$ bond, which did not require the assistance of palladium catalysts. On the basis of these results from eqn (1) \dagger and previous work reported by Smith and Takeda,¹⁵ we speculated that **8a** and **8a'** undergo a Brook- and retro-Brook-type rearrangement to afford **3a** (path c and path d). To capture carbanion species **F** and **F'**, two common electrophilic reagents, such as iodomethane and benzyl bromide, were added for the silacyclization of **8a** and **8a'** (eqn (2) and (3) \dagger). However, the corresponding products **F-1** and **F'-2** were not observed. Therefore, these results are more favorable to this pathway involving synergistic Brook/retro-Brook-type rearrangement (path d).

Based on the results of mechanistic experiments as well as reported work,^{12,13,15} a plausible mechanism for the synthesis of silacycles was proposed (Scheme 8). Initially, oxidative addition followed by intramolecular Heck-cyclization of acrylamides **1a** to $\text{Pd}(0)$ species forms intermediate **A**, which then undergoes a C-H activation to afford spiropalladacycle **B**. Next, spiropalladacycle **B** produces disilylated products **8a/8a'** and regenerates $\text{Pd}(0)$ by sequential transmetalation with intermediate **G** generated by disilane **2a**, reductive elimination and further hydrolysis (path a). **8a** and **8a'** then undergo a synergistic Brook/retro-Brook-type rearrangement to afford **3a** and a trimethylsiloxy anion, which could be converted into TMSOTMS (path d).¹⁶ Notably, another possible pathway that synthesizes disilylated products **8a/8a'** by direct oxidative addition of spiropalladacycle **B** with disilane **2a** is ruled out by the results of eqn (4) \dagger (path b).



Scheme 8 Possible reaction mechanism.

Conclusions

In conclusion, we have disclosed the first example of divergent synthesis of silacycles *via* a Brook- and retro-Brook-type rearrangement strategy by employing a readily accessible disilane reagent $\text{Me}_3\text{SiSiMe}_2(\text{O}^n\text{Bu})$. In this novel transformation, divergent silacycles, such as spirobenzosiloles, fused benzsiloles, and π -conjugated dibenzosiloles, can be produced in moderate to good yields by an unprecedented complex process composed of a bis-silylation of a palladacycle and a Brook- and retro-Brook-type rearrangement. Notably, mechanistic studies reveal that bis-silylation of the palladacycle is completed by a transmetalation process. Further applications of the disilane reagent $\text{Me}_3\text{SiSiMe}_2(\text{O}^n\text{Bu})$ and the rearrangement are still in progress in our laboratory.

Data availability

All experimental procedures and spectroscopic data can be found in the ESI.[†]

Author contributions

Y. X. designed and performed the experiments and analysed the data. W. X., X. C., X. L., H. L., M. Z., X. Y. and G. D. contributed to data analysis and scientific discussion. Y. Y. and Y. L. supervised the project, analyzed the results and wrote the manuscript.

Conflicts of interest

The authors declare no competing interests.

Acknowledgements

The authors thank the National Natural Science Foundation of China (21901071 and 21971061), the Natural Science Foundation of Hunan Province (2020JJ5350), the Scientific Research Fund of Hunan Provincial Education Department (18A002 and 19B359), and the Science and Technology Planning Project of Hunan Province (2018TP1017) for financial support. We thank Prof. Yuanhong Ma (Hunan Normal University) for helpful discussions and Dr Xiaoming Zhu (Hengyang Normal University) for the analysis of X-ray crystallography.

Notes and references

- (a) A. K. Franz and S. O. Wilson, *J. Med. Chem.*, 2013, **56**, 388; (b) J. B. Grimm, T. A. Brown, A. N. Tkachuk and L. D. Lavis, *ACS Cent. Sci.*, 2017, **3**, 975; (c) R. Ramesh and D. S. Reddy, *J. Med. Chem.*, 2018, **61**, 3779; (d) K. Umezawa, M. Kamiya and Y. Urano, *Angew. Chem., Int. Ed.*, 2018, **57**, 9346; (e) S. J. Barraza and S. E. Denmark, *J. Am. Chem. Soc.*, 2018, **140**, 6668; (f) K. L. Chan, M. J. McKiernan, C. R. Towns and A. B. Holmes, *J. Am. Chem. Soc.*, 2005, **127**, 7662; (g) M. Akhtaruzzaman, Y. Seya, N. Asao, A. Islam, E. Kwon,

A. Han, L. El-Shafei and Y. Yamamoto, *J. Mater. Chem.*, 2012, **22**, 10771.

- 2 For recent reviews, see: (a) U. Sharma, R. Sharma, R. Kumar, I. Kumar and B. Singh, *Synthesis*, 2015, **47**, 2347; (b) L. Li, Y. Zhang, L. Gao and Z. Song, *Tetrahedron Lett.*, 2015, **56**, 1466; (c) Y. Yang and C. Wang, *Sci. China: Chem.*, 2015, **58**, 1266; (d) Z. Xu, W.-S. Huang, J. Zhang and L.-W. Xu, *Synthesis*, 2015, **47**, 3645; (e) Q. W. Zhang, K. An and W. He, *Synlett*, 2015, **9**, 1145; (f) T. Komiyama, Y. Minami and T. Hiyama, *ACS Catal.*, 2016, **7**, 631; (g) W.-T. Zhao, F. Gao and D. Zhao, *Synlett*, 2018, **29**, 2595; (h) Q.-C. Mu, J. Chen, C.-G. Xia and L.-W. Xu, *Coord. Chem. Rev.*, 2018, **374**, 93; (i) S. C. Richter and M. Oestreich, *Trends Chem.*, 2020, **2**, 13.

- 3 For selected examples, see: (a) T. Matsuda, S. Kadowaki, Y. Yamaguchi and M. Murakami, *Chem. Commun.*, 2008, 2744; (b) M. Tobisu, M. Onoe, Y. Kita and N. Chatani, *J. Am. Chem. Soc.*, 2009, **131**, 7506; (c) K. Mochida, M. Shimizu and T. Hiyama, *J. Am. Chem. Soc.*, 2009, **131**, 8350; (d) E. Shirakawa, S. Masui, R. Narui, R. Watabe, D. Ikeda and T. Hayashi, *Chem. Commun.*, 2011, **47**, 9714; (e) Y. Liang, S. Zhang and Z. Xi, *J. Am. Chem. Soc.*, 2011, **133**, 9204; (f) R. Shintani, K. Moriya and T. Hayashi, *J. Am. Chem. Soc.*, 2011, **133**, 16440; (g) Y. Liang, W. Geng, J. Wei and Z. Xi, *Angew. Chem., Int. Ed.*, 2012, **51**, 1934; (h) M. Onoe, K. Baba, Y. Kim, Y. Kita, M. Tobisu and N. Chatani, *J. Am. Chem. Soc.*, 2012, **134**, 19477; (i) Q.-W. Zhang, K. An and W. He, *Angew. Chem., Int. Ed.*, 2014, **53**, 5667; (j) N. Ishida, W. Ikemoto and M. Murakami, *J. Am. Chem. Soc.*, 2014, **136**, 5912; (k) Q.-W. Zhang, K. An, L.-C. Liu, S. Guo, C. Jiang, H. Guo and W. He, *Angew. Chem., Int. Ed.*, 2016, **55**, 6319; (l) Q.-W. Zhang, K. An, L.-C. Liu, Q. Zhang, H. Guo and W. He, *Angew. Chem., Int. Ed.*, 2017, **56**, 1125; (m) S. Okumura, F. Sun, N. Ishida and M. Murakami, *J. Am. Chem. Soc.*, 2017, **139**, 12414; (n) Q. Yang, L. Liu, Y. Chi, W. Hao, W.-X. Zhang and Z. Xi, *Org. Chem. Front.*, 2018, **5**, 860; (o) W. T. Zhao, F. Gao and D. Zhao, *Angew. Chem., Int. Ed.*, 2018, **57**, 6329; (p) H. Chen, Y. Chen, X. Tang, S. Liu, R. Wang, T. Hu and Z. Song, *Angew. Chem., Int. Ed.*, 2019, **58**, 4695; (q) X. B. Wang, Z. J. Zheng, J. L. Xie, X. W. Gu, Q. C. Mu, G. W. Yin, F. Ye, Z. Xu and L. W. Xu, *Angew. Chem., Int. Ed.*, 2020, **59**, 790; (r) Y. Qin, J. L. Han, C. W. Ju and D. Zhao, *Angew. Chem., Int. Ed.*, 2020, **59**, 8481.

- 4 For reviews, see: (a) A. G. Brook and A. R. Bassendale, in *Rearrangements in Ground and Excited States*, ed. P. de Mayo, Academic Press, New York, 1980, vol. 2, p. 149; (b) K. Tomooka, in *The Chemistry of Organolithium Compounds*, ed. Z. Rappoport and I. Marck, Wiley, Chichester, 2004, vol. 2, p. 749; (c) A. G. Brook, *Acc. Chem. Res.*, 1974, **7**, 77; (d) W. H. Moser, *Tetrahedron*, 2001, **57**, 2065; (e) E. Schaumann and A. Kirschning, *Synlett*, 2007, **18**, 177; (f) A. B. Smith III and W. M. Wuest, *Chem. Commun.*, 2008, 5883; (g) A. B. Smith III and C. M. Adams, *Acc. Chem. Res.*, 2004, **37**, 365; (h) Y. Deng and A. B. Smith III, *Acc. Chem. Res.*, 2020, **53**, 988.



5 (a) A. G. Brook, *J. Am. Chem. Soc.*, 1958, **80**, 1886; (b) A. G. Brook, C. M. Warner and M. E. McGriskin, *J. Am. Chem. Soc.*, 1959, **81**, 981; (c) A. G. Brook and N. V. Schwartz, *J. Am. Chem. Soc.*, 1960, **82**, 2435; (d) A. G. Brook and B. Iachia, *J. Am. Chem. Soc.*, 1961, **83**, 827.

6 (a) R. West, R. Lowe, H. F. Stewart and A. Wright, *J. Am. Chem. Soc.*, 1971, **93**, 282; (b) R. J. Linderman and A. Ghannam, *J. Am. Chem. Soc.*, 1990, **112**, 2392; (c) X. Jiang and W. F. Bailey, *Organometallics*, 1995, **14**, 5704; (d) T. Kawashima, K. Naganuma and R. Okazaki, *Organometallics*, 1998, **17**, 367; (e) K. Naganuma, T. Kawashima and R. Okazaki, *Chem. Lett.*, 1999, **28**, 1139.

7 For selected examples, see: (a) A. Wright and R. J. West, *J. Am. Chem. Soc.*, 1974, **96**, 3214; (b) A. Wright and R. West, *J. Am. Chem. Soc.*, 1974, **96**, 3227; (c) W. C. Still and T. Macdonald, *J. Am. Chem. Soc.*, 1974, **96**, 5561; (d) R. J. Linderman and A. Ghannam, *J. Am. Chem. Soc.*, 1990, **112**, 2392; (e) X. L. Jiang and W. F. Bailey, *Organometallics*, 1995, **14**, 5704; (f) Y. Mori, Y. Futamura and K. Horisaki, *Angew. Chem., Int. Ed.*, 2008, **47**, 1091.

8 (a) M. E. Jung and C. J. Nichols, *J. Org. Chem.*, 1996, **61**, 9065; (b) E. Bures, P. G. Spinazze, G. Beese, I. R. Hunt, C. Rogers and B. A. Keay, *J. Org. Chem.*, 1997, **62**, 8741; (c) J. Clayden, D. W. Watson and M. Chambers, *Tetrahedron*, 2005, **61**, 3195.

9 For reviews, see: (a) K. A. Horn, *Chem. Rev.*, 1995, **95**, 1317; (b) H. K. Sharma and K. H. Pannell, *Chem. Rev.*, 1995, **95**, 1351; (c) M. Suginome and Y. Ito, *J. Chem. Soc., Dalton Trans.*, 1998, 1925; (d) I. Beletskaya and C. Moberg, *Chem. Rev.*, 1999, **99**, 3435; (e) M. Suginome and Y. Ito, *Chem. Rev.*, 2000, **100**, 3221; (f) I. Beletskaya and C. Moberg, *Chem. Rev.*, 2006, **106**, 2320; (g) C. Cheng and J. F. Hartwig, *Chem. Rev.*, 2015, **115**, 8946; (h) P. Xiao, L. Gao and Z. Song, *Chem.*, 2019, **25**, 2407; (i) Y. Zhang, B. Zhou and A. Lu, *Synlett*, 2018, **30**, 685.

10 For recent representative examples of mono-silylation, see: (a) Y. J. Liu, Y. H. Liu, Z. Z. Zhang, S. Y. Yan, K. Chen and B.-F. Shi, *Angew. Chem., Int. Ed.*, 2016, **55**, 13859; (b) B.-B. Zhan, J. Fan, L. Jin and B.-F. Shi, *ACS Catal.*, 2019, **9**, 3298; (c) A. Maji, S. Guin, S. Feng, A. Dahiya, V. K. Singh, P. Liu and D. Maiti, *Angew. Chem., Int. Ed.*, 2017, **56**, 14903; (d) A. Deb, S. Singh, K. Seth, S. Pimparkar, B. Bhaskararao, S. Guin, R. B. Sunoj and D. Maiti, *ACS Catal.*, 2017, **7**, 8171; (e) H. Saito, K. Nogi and H. Yorimitsu, *Angew. Chem., Int. Ed.*, 2018, **57**, 11030.

11 For recent representative examples of bis-silylation, see: (a) M. B. Ansell, D. E. Roberts, F. G. Cloke, O. Navarro and J. Spencer, *Angew. Chem., Int. Ed.*, 2015, **54**, 5578; (b) Z. Liu, H. Tan, T. Fu, Y. Xia, D. Qiu, Y. Zhang and J. Wang, *J. Am. Chem. Soc.*, 2015, **137**, 12800; (c) M. Ahmad, A. C. Gaumont, M. Durandett and J. Maddaluno, *Angew. Chem., Int. Ed.*, 2017, **56**, 2464; (d) T. He, L. C. Liu, L. Guo, B. Li, Q. W. Zhang and W. He, *Angew. Chem., Int. Ed.*, 2018, **57**, 10868; (e) P. H. Xiao, Y. J. Cao, Y. Y. Gui, L. Gao and Z. L. Song, *Angew. Chem., Int. Ed.*, 2018, **57**, 4769.

12 (a) A. Lu, X. Ji, B. Zhou, Z. Wu and Y. Zhang, *Angew. Chem., Int. Ed.*, 2018, **57**, 3233; (b) X. Ji, F. Wei, B. Wan, C. Cheng and Y. Zhang, *Chem. Commun.*, 2020, **56**, 7801; (c) B. Zhou, A. Lu, C. Shao, X. Liang and Y. Zhang, *Chem. Commun.*, 2018, **54**, 10598; (d) X. Ma, A. Lu, X. Ji, G. Shi and Y. Zhang, *Asian J. Org. Chem.*, 2018, **7**, 1403; (e) W. Lv, S. Wen, J. Yu and G. Cheng, *Org. Lett.*, 2018, **20**, 4984; (f) W. Lv, J. Yu, B. Ge, S. Wen and G. Cheng, *J. Org. Chem.*, 2018, **83**, 12683; (g) W. Lv, S. Liu, Y. Chen, S. Wen, Y. Lan and G. Cheng, *ACS Catal.*, 2020, **10**, 10516; (h) M. Wollenburg, J. Bajohr, A. D. Marchese, A. Whyte, F. Glorius and M. Lautens, *Org. Lett.*, 2020, **22**, 3679.

13 (a) H. Lu, X. Yang, L. Zhou, W. Li, G. Deng, Y. Yang and Y. Liang, *Org. Chem. Front.*, 2020, **7**, 2016; (b) W. Li, C. Zhang, H. Lu, Y. Wang, G. Deng, Y. Liang and Y. Yang, *Org. Chem. Front.*, 2020, **7**, 2075; (c) W. Li, W. Chen, B. Zhou, Y. Xu, G. Deng, Y. Liang and Y. Yang, *Org. Lett.*, 2019, **21**, 2718; (d) Y. Xu, X. Liu, W. Chen, G. Deng, Y. Liang and Y. Yang, *J. Org. Chem.*, 2018, **83**, 13930; (e) G. Xiao, L. Chen, G. Deng, J. Liu and Y. Liang, *Tetrahedron Lett.*, 2018, **59**, 1836; (f) W. Li, G. Xiao, G. Deng and Y. Liang, *Org. Chem. Front.*, 2018, **5**, 1488.

14 H. Yoon, A. Lossouarn, F. Landau and M. Lautens, *Org. Lett.*, 2016, **18**, 6324.

15 (a) A. B. Smith III, R. Tong, W.-S. Kim and W. A. Maio, *Angew. Chem., Int. Ed.*, 2011, **50**, 8904; (b) H. Taguchi, K. Tadaki, M. Ghoroku, A. Tsubouchi and T. Takeda, *J. Org. Chem.*, 2002, **67**, 8450; (c) H. Taguchi, K. Takami, A. Tsubouchi and T. Takeda, *Tetrahedron Lett.*, 2004, **45**, 429.

16 (a) K. J. Shea and D. A. Loy, *Acc. Chem. Res.*, 2001, **34**, 707; (b) F. Neumeyer and N. Auner, *Chem.-Eur. J.*, 2016, **22**, 1; (c) L. V. Zhilitskaya, N. O. Yarosh, L. G. Shagun, I. A. Dorofeev and L. I. Larina, *Mendeleev Commun.*, 2017, **27**, 352.

