Chemical Science



EDGE ARTICLE

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2021, 12, 15727

dll publication charges for this article have been paid for by the Royal Society of Chemistry

Received 30th September 2021 Accepted 8th November 2021

DOI: 10.1039/d1sc05404j

rsc.li/chemical-science

Dibenzocycloheptanones construction through a removable *P*-centered radical: synthesis of allocolchicine analogues†

Yan Zhang, (1)** Zhenzhi Cai, a Julia Struwe, b Chanchan Ma, a Wangyu Zeng, a Xinyi Liao, a Min Xua and Lutz Ackermann (1)** **

Dibenzocycloheptanones containing a tricyclic 6–7–6-system are present in numerous biologically active natural molecules. However, the simple and efficient preparation of derivatives containing a dibenzocycloheptanone scaffold remains difficult to date. Herein, we report a versatile strategy for the construction of these challenging seven-membered rings using a 7-endo-trig cyclization which is initiated by a phosphorus-centered radical. This approach provides a step-economical regime for the facile assembly of a wide range of phosphorylated dibenzocycloheptanones. Remarkably, we also have devised a traceless addition/exchange strategy for the preparation of dephosphorylated products at room temperature with excellent yields. Therefore, this protocol allows for the concise synthesis of biorelevant allocochicine derivatives.

Introduction

Seven-membered rings represent a class of structurally special compounds with a synthetically difficult scaffold.¹ In particular, the 6–7–6 ring system has been found to be a privileged structure which is present in a number of marketed natural products and bioactive molecules, such as *N*-acetylcolchicinol methyl ether (NSC 51046), ZD6126 and other pharmaceutical compounds (Fig. 1, bottom).² However, because of inherent transannular and entropic factors,⁴ the synthesis of functionalized 6–7–6 ring systems still represents an ongoing challenge compared to numerous efficient and modular procedures to access common six- or five-membered cyclic compounds through traditional substitution, condensation and coupling reactions.6

During the last decades, various synthetic methods producing allocolchicine analogues have been developed. These examples include enyne ring-closing metathesis/Diels-Alder reaction⁷ or Diels-Alder reaction/aromatization approach, palladium-catalyzed intramolecular direct arylation, intramolecular Nicholas reaction, oxidative coupling, then penanthrol ring expansion, electrophilic iodocyclization and

Representative privileged scaffolds of allocochicine analogues

Fig. 1 Dibenzocycloheptanones construction with reaction design, and selected examples of bioactive 6–7–6 tricyclic compounds.

Suzuki–Miyaura coupling/Aldol condensation. ¹⁴ Although those procedures definitely represent attractive strategies to access the challenging 6–7–6 ring system, the continuing investigation of direct and efficient approaches towards this core structure is still highly valuable. The construction of new chemical bonds by radical addition reactions represents a powerful tool for chemical transformations and has attracted considerable recent attention, ¹⁵ as this approach usually features mild reaction conditions, predictable reaction outcome, a high functional group tolerance and the use of diverse radical precursors

This Work

P(O)(OEt)2
7-endo-process
(R = aryl or alkyl)

7-endo-trig process
excellent regioselectivity

> 60 new six-seven-six tricylic products

[&]quot;Key Laboratory of the Ministry of Education for Advanced Catalysis Materials, Drug Discovery & Innovation Center, College of Chemistry and Life Sciences, Zhejiang Normal University, China. E-mail: zhangyan001@zjnu.edu.cn

^bInstitut für Organische und Biomolekulare Chemie, Georg-August-Universität Göttingen, Germany. E-mail: Lutz.Ackermann@chemie.uni-goettingen.de

[†] Electronic supplementary information (ESI) available: Experimental details and characterization of all new compounds and details for DFT calculations. CCDC 2103940 and 2103941. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1sc05404j

and initiation strategies. As 7-endo-cyclization reaction are considered to be favorable following the Baldwin rules and the extended ones from Alabugin, ¹⁶ we questioned whether the efficiency noted in a removable radical formation could be translated into a practical formation strategy for 6-7-6 scaffolds. Although this direct radical annulation approach seems to be straightforward for the construction of the tricyclic core, key challenges remained to be addressed (Fig. 1, top).

First, the regioselectivity between 5-exo, 6-exo and 7-endo cyclization needs to be well controlled by adjusting the influence of substituents on the Ar'-moiety. Compared to five- or sixmembered ring formation, the use of the 7-endo-trig approach is rare to construct seven-membered rings via radical cyclization. 17 Second, similar to common directing groups in C-H activation, almost all radical sources will also leave behind undesired chemical "foot prints",18 thereby requiring further synthetic derivatization and restricting the method to the desired skeleton of the products. Third, only few radical cascade reactions initiated by the addition of a phosphonyl radical to alkynes were reported, presumably due to slow kinetics of radical [(RO)₂P(O) '] addition step. 19,20 Thus, the development of efficient P-radical tandem procedures and step-economical routes for the construction of various seven-membered rings is still desirable. Within our ongoing interest in radical addition reactions to the alkynones, 20a,21 we herein report an unprecedented cascade cyclization of biaylynones, which can be performed with commercially available, inexpensive reagents in water-tolerant radical reaction conditions. Notable features of our strategy include (a) the first phosphonyl radical addition to internal alkynes resulting in a 7-endo-trig process, (b) the P-centered radical motif can be removed traceless under operationallysimple conditions, (c) excellent regioselectivity, and (d) ample substrate scope.

Results and discussion

Optimization of reaction condition

We initiated our studies by probing various reaction conditions for the envisioned cascade cyclization of 1-(3',5'-dimethoxy-[1,1'-biphenyl]-2-yl)-3-phenylprop-2-yn-1-one (1a) as the model substrate (Tables 1 and S2 in the ESI†), using the inexpensive HPO(OEt)₂ (2) as the phosphonyl radical source. Gratifyingly, the desired 6-7-6 tricyclic product 3 was isolated in 60% yield, when the reaction was performed in the presence of AgNO₃/ K₂S₂O₈ in a 1,4-dioxane/H₂O solvent system at 70 °C for 8 h. Different reaction media were explored, but fell short in efficiently delivering the desired products (2-8). Furthermore, a series of chemical oxidants was tested but showed not to be beneficial (entries 9-12). Either decreasing or increasing the reaction temperature failed to improve the yield of 3 (entry 13). Not surprisingly, substituents on the Ar'-ring have a decisive influence on the formation of the seven-membered ring. Therefore, a substitution effect was first examined under the optimized conditions (entry 11 in Table S2†). Indeed, the type and position of substituents on the arene moiety determined the efficacy of the 7-endo-trig cyclization process. We were pleased to find that 4-(tert-butyl)phenyl alkynone 1i gave the

Table 1 Optimization of the cascade cyclization/phosphorylation^a

Entry	Deviation from standard conditions	Yield/%
1	No change	60
2	MeCN	41
3	1,4-Dioxane	27
4	$MeCN/H_2O(1:1)$	Trace
5	$DMSO/H_2O(1:1)$	Trace
6	$DMF/H_2O(1:1)$	31
7	$MeOH/H_2O(1:1)$	Trace
8	$THF/H_2O(1:1)$	40
9	Cu(OTf) ₂ and THBP in MeCN	Trace
10	Mn(OAc) ₃ (2.5 equiv.) in MeCN	15
11	$AgNO_3$ (2.0 equiv.)	48
12	Na ₂ S ₂ O ₈ instead of K ₂ S ₂ O ₈	54
13	Reaction at 60 °C	52

 a Standard conditions: 1a (0.2 mmol, 1.0 equiv), 2 (0.5 mmol, 2.5 equiv.), $K_2S_2O_8$ (0.5 mmol, 2.5 equiv.), AgNO₃ (20 mol%), 1,4-dioxane/H₂O (1 : 1, 2 mL), under N₂, 70 °C, 10 h. Yield of the isolated product.

best result and the desired product **11** was thereby obtained in 67% yield (Fig. 2).

Robustness

With the optimized conditions in hand, we became interested in investigating the substrate scope of this radical cyclization and we tested a diverse range of biarylyone substrates with different aromatic rings Ar and R (Scheme 1). The reaction tolerated a variety of H-phosphonates and substituents with diverse electronic properties in all positions of the arene (12–





Fig. 2 Substituent impact on 7-endo radical cyclization process.^a Standard conditions: **1a** (0.2 mmol, 1.0 equiv.), **2** (0.5 mmol, 2.5 equiv.), $K_2S_2O_8$ (0.5 mmol, 2.5 equiv.), $AgNO_3$ (20 mol%), 1,4-dioxane/ H_2O (1:1,2 mL), under N_2 , 70 °C, 10 h. Yield of the isolated products. ^b6-Exo product **5** formed in 60% yield.



Scheme 1 Scope of 7-endo-trig cyclization process. Reaction conditions from Table 1, entry 1, alkynone (1, 0.3 mmol), 6-10 h. ^aAfter being stirred at 70 °C for 10 h, the mixture was neutralized with NaHCO₃ until pH 7-8 and extracted with ethyl acetate.

21), with the mass balance largely accounting for unreacted starting material. We were pleased to find that ortho- and metasubstituted arenes R underwent this transformation efficiently despite a possible steric repulsion (18-21), even for polysubstituted substrates. Furthermore, heterocyclic substrates bearing thiophene, pyridine and benzothiophene as well as naphthyl substituents were also tolerated in this transformation (22-25). After having demonstrated the broad applicability with respect to the arene R, substitutions on the phenone scaffold were examined as well. Substrates decorated with both electronwithdrawing and electron-donating groups on the aryl ring Ar did not significantly alter the reaction efficacy and the tricyclic compounds were efficiently obtained (26-38). As depicted, the 7-endo- approach was compatible with valuable sensitive functional groups, such as chloro-, bromo-, ether-, nitro-, cyano- and ester-functionalities. A heterocyclic substrate was also applicable in this transformation to afford the corresponding product 39 in good yield. Noteworthy, also alkyl alkynes were applicable and gave minor amounts of the corresponding products (40-41). The connectivity of product 31 was

unambiguously confirmed by single-crystal X-ray analysis (Scheme 1).²²

In order to increase the potential synthetic elaboration, we also explored an access to other classes of allocolchicine analogues. To our delight, we found that the incorporated phosphonyl motif could be easily removed in the presence of alkoxide or hydroxide bases. Next, the substrate scope of this traceless bond-constructing strategy was explored (Scheme 2). As expected, this mild anion addition/exchange approach was found to be generally applicable for the dephosphorylation (43–61). The structure of product 59 was unambiguously verified by X-ray crystallographic analysis. ^{22b}

To further demonstrate the synthetic utility, the developed strategy was exploited for the preparation of bioactive NSC

Scheme 2 Scope of dephosphorylation process.

61: 80%

51046 analogues (Scheme 3). NSC 51046 is a natural product which displays potent anticancer activity by inhibition of the tubulin polymerization.2c Due to the modularity of the our synthesis, this approach could be beneficial for preparing diverse NSC 51046 analogues from 3-phenyl-1-(3',4,5'-trimethoxy-[1,1'-biphenyl]-2-yl)prop-2-yn-1-one 1n' which could be easily prepared from 2-bromo-5-methoxybenzaldehyde via cross-coupling reaction. Under the standard reaction conditions, the phosphorylation product 62 was obtained in gramscale. Subsequently, phosphonate 62 was converted to the dephosphorylation product enone 63 following our defunctionalization procedure. Reduction of alkene 63 led to the formation of intermediate 64, which provided the allocolchicine 66 after reduction followed by acetylation. Thus, this strategy opens a new avenue to a divergent synthesis of allocolchicine analogues with readily available starting materials and high efficacy.

Considering that this conversion process includes a radical addition step, a control experiment was conducted thereafter. The key radical intermediate A could be trapped by 2,2,6,6tetramethyl-1-piperidinyloxy (TEMPO) to form the TEMPOadduct 67 (Fig. 3a). Based on these experimental result and related findings,23 a plausible mechanism involving a radicaltype catalytic cycle is proposed as depicted in Fig. 3. First, the P-radical is generated from diethyl H-phosphonate (2) initiated by AgNO₃. Selective radical addition of the P-radical to the α position of the carbonyl in 1i affords the vinyl radical A, which undergoes 7-endo-trig cyclization resulting in intermediate B. Then, intermediate B undergoes further SET oxidation to form the product 11. In addition, we also found by pH measurement that after the reaction between 1i and 2 was performed under standard conditions, the reaction mixture became clearly acidic. Finally, enone 43 could be obtained after dephosphorylation involving an alkoxy anion addition/exchange process,

Scheme 3 Gram-scale reaction and NSC 51046 analogue synthesis.

60: 78%

Edge Article Chemical Science

Fig. 3 Mechanistic investigation on the radical annulation and dephosphorylation process (a) radical trapping experiment, (b) proposed mechanism.

generating **42** as a byproduct that can be detected by HRMS and ³¹P-NMR (see the ESI†).

Conclusions

In summary, we have developed an efficient approach for the construction of dibenzocycloheptanones through a cascade radical addition of biarylynones with diethylphosphite as the radical source. This versatile strategy showed high regioselectivity, ample scope and good functional group compatibility. It is worth mentioning that the introduced radical fragment of the phosphorylated products was easily removed in the presence of bases to give the corresponding allocolchicine analogues. The developed approach represents one of the rare 7-endo-trig radical cyclization processes and is a useful method for the concise construction of a variety of novel and potentially useful drug-type, and natural product-like molecules bearing valuable 6-7-6-scaffolds.

Data availability

All experimental data, procedures for data analysis and pertinent data sets are provided in the ESI. \dagger

Author contributions

Y. Z. and L. A. conceived the project. Y. Z., Z. C., C. M. W. Z., X. L. and M. X. performed the experiments, Y. Z., Z.C. and C. M. and J. S. analyzed and interpreted the experimental data. Y. Z. and J. S. drafted the paper. Y. Z. and L. A. supervised the project. All of

the authors discussed the results and contributed to the preparation of the final manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Generous support by the National Natural Science Foundation of China (No. 21702188) and by the DFG (Gottfried-Wilhelm-Leibniz award to LA) is gratefully acknowledged.

Notes and references

- 1 (*a*) I. Shiina, *Chem. Rev.*, 2007, **107**, 239–273; (*b*) T. P. Majhi, B. Achari and P. Chattopadhyay, *Heterocycles*, 2007, **71**, 1011–1052; (*c*) M. Inoue, *Chem. Rev.*, 2005, **105**, 4379–4405.
- 2 (a) T. Graening and H.-G. Schmalz, Angew. Chem., Int. Ed.,
 2004, 43, 3230–3256; (b) B. Pérez-Ramírez, M. J. Gorbunoff and S. N. Timasheff, Biochemistry, 1998, 37, 1646–1661; (c)
 A. Brossi, J. Med. Chem., 1990, 33, 2311–2319.
- 3 (a) F. Büttner, S. Bergemann, D. Guénard, R. Gust, G. Seitz and S. Thoret, *Bioorg. Med. Chem.*, 2005, 13, 3497–3511; (b) K. Nakagawa-Goto, M. K. Jung, E. Hamel, C.-C. Wu, K. F. Bastow, A. Brossi, S. Ohta and K.-H. Lee, *Heterocycles*, 2005, 65, 541–550; (c) Q. Shi, K. Chen, X. Chen, A. Brossi, P. Verdier-Pinard, E. Hamel, A. T. McPhail, A. Tropsha and K.-H. Lee, *J. Org. Chem.*, 1998, 63, 4018–4025.
- 4 (a) R. L. Reyes, T. Iwai and M. Sawamura, *Chem. Rev.*, 2021, **121**, 8926–8947; (b) L. Yet, *Chem. Rev.*, 2000, **100**, 2963–3007.
- 5 G. Illuminati and L. Mandolini, *Acc. Chem. Res.*, 1981, 14, 95– 102.
- 6 (a) R. S. Doerksen, T. Hodík, G. Hu, N. O. Huynh,
 W. G. Shuler and M. J. Krische, *Chem. Rev.*, 2021, 121,
 4045–4083; (b) J.-R. Chen, X.-Q. Hu, L.-Q. Lu and
 W.-J. Xiao, *Chem. Rev.*, 2015, 115, 5301–5365.
- 7 F.-D. Boyer and I. Hanna, Org. Lett., 2009, 9, 715-718.
- 8 A. V. Vorogushin, A. V. Predeus, W. D. Wulff and H.-J. Hansen, *J. Org. Chem.*, 2003, **68**, 5826–5831.
- 9 M. Leblanc and K. Fagnou, Org. Lett., 2005, 7, 2849-2852.
- 10 (a) S. Djurdjevic, F. Yang and J. R. Green, J. Org. Chem., 2010, 75, 8241–8251; (b) S. Djurdjevic and J. R. Green, Org. Lett., 2007, 9, 5505–5508.
- 11 G. Besong, K. Jarowicki, P. J. Kocienski, E. Sliwinski and F. T. Boyle, *Org. Biomol. Chem.*, 2006, 4, 2193–2207.
- 12 W. M. Seganish and P. DeShong, *Org. Lett.*, 2006, **8**, 3951–3954.
- 13 Y. Chen, C. Huang, X. Liu, E. Perl, Z. Chen, J. Namgung, G. Subramaniam, G. Zhang and W. H. Hersh, *J. Org. Chem.*, 2014, 79, 3452–3464.
- 14 Y. L. Choi, C.-M. Yu, B. T. Kim and J.-N. Heo, *J. Org. Chem.*, 2009, 74, 3948–3951.
- 15 For reviews on radical chemistry, see: (*a*) H. Xiao, Z. Zhang, Y. Fang, L. Zhu and C. Li, *Chem. Soc. Rev.*, 2021, **50**, 6308–6319; (*b*) H. Zhou, Z.-L. Li, Q.-S. Gu and X.-Y. Liu, *ACS Catal.*, 2021, **11**, 7978–7986; (*c*) M. Latrache and

N. Hoffmann, Chem. Soc. Rev., 2021, 50, 7418–7435; (d) D. Leifert and A. Studer, Angew. Chem., Int. Ed., 2020, 59, 74–108; (e) Q.-Q. Zhou, Y.-Q. Zou, L.-Q. Lu and W.-J. Xiao, Angew. Chem., Int. Ed., 2019, 58, 1586–1604; (f) Y. Wei, P. Hu, M. Zhang and W. Su, Chem. Rev., 2017, 117, 8864–8907; (g) A. Studer and D. P. Curran, Angew. Chem., Int. Ed., 2016, 55, 58–102; (h) M.-C. Belhomme, T. Besset, T. Poisson and X. Pannecoucke, Chem. - Eur. J., 2015, 21, 12836–12865; (i) E. Merino and C. Nevado, Chem. Soc. Rev., 2014, 114, 2587–2693; (j) F. Dénès, M. Pichowicz, G. Povie and P. Renaud, Chem. Rev., 2014, 114, 2587–2693; (k) U. Wille, Chem. Rev., 2013, 113, 813–853; (l) P. Chen and G. Liu, Synthesis, 2013, 2919–2939.

- 16 K. Gilmore and I. V. Alabugin, *Chem. Rev.*, 2011, **111**, 6513–6556
- 17 (a) P. Xiong, H.-H. Xu, J. Song and H.-C. Xu, *J. Am. Chem. Soc.*, 2018, **140**, 2460–2464; (b) Y. Li and J.-H. Li, *Org. Lett.*, 2018, **20**, 5323–5326.
- 18 Y. Lu, D.-H. Wang, K. M. Engle and J.-Q. Yu, *J. Am. Chem. Soc.*, 2010, **132**, 5916–5921.
- 19 For reviews on phosphorous-centered radical reactions, see: (a) X.-Q. Pan, J.-P. Zou, W.-B. Yi and W. Zhang, *Tetrahedron*, 2015, 71, 7481–7529; (b) D. Leca, L. Fensterbank, E. Lacôte

- and M. Malacria, *Chem. Soc. Rev.*, 2005, **34**, 858–865; (c) W. G. Bentrude, *Acc. Chem. Res.*, 1982, **15**, 117–125.
- 20 For phosphonyl radical addition to alkynes, see: (a) Y. Zhang, J. Zhang, B. Hu, M. Ji, S. Ye and G. Zhu, *Org. Lett.*, 2018, **20**, 2988–2992; (b) Z.-D. Liu, W.-Z. Bi, Y.-Y. Xia, H.-T. Wu and Y.-F. Zhao, *Chem. Commun.*, 2015, **51**, 3846–3849; (c) L.-J. Wang, A.-Q. Wang, Y. Xia, X.-X. Wu, X.-Y. Liu and Y.-M. Liang, *Chem. Commun.*, 2014, **50**, 13998–14001; (d) X. Mi, C. Wang, M. Huang, J. Zhang, Y. Wu and Y. Wu, *Org. Lett.*, 2014, **16**, 3356–3359; (e) X.-Q. Pan, J.-P. Zou, G.-L. Zhang and W. Zhang, *Chem. Commun.*, 2010, **46**, 1721–1723.
- 21 (a) Y. Zhang, C. Ma, J. Struwe, J. Feng, G. Zhu and L. Ackermann, *Chem. Sci.*, 2021, 12, 10092–10096; (b) Y. Zhang, D. Guo, S. Ye, Z. Liu and G. Zhu, *Org. Lett.*, 2017, 19, 1302–1305; (c) W. Ma and L. Ackermann, *Synthesis*, 2014, 2297–2304.
- 22 (a) CCDC 2103940;†; (b) CCDC 2103941.†
- 23 (a) X. Tan, Z. Liu, H. Shen, P. Zhang, Z. Zhang and C. Li, *J. Am. Chem. Soc.*, 2017, **139**, 12430–12433; (b) Z.-Z. Zhou, X.-B. Yan, X.-Y. Liu and Y.-M. Liang, *Org. Lett.*, 2014, **16**, 5616–5619.