

Cite this: *Food Funct.*, 2021, 12, 2389

Marama bean [*Tylosema esculentum* (Burch.) A. Schreib.]: an indigenous plant with potential for food, nutrition, and economic sustainability

Abiodun Olusola Omotayo ^a and Adeyemi Oladapo Aremu ^{*a,b}

Developing countries need to explore undervalued indigenous plants to fully enhance their food and nutrition security, health, and economic viability. This review explores the nutritional, phytochemical, and economic potential of marama bean (*Tylosema esculentum*, Fabaceae), a non-nodulating indigenous legume that can be cultivated in and is well-adapted to dry or low moisture conditions. Marama bean is popularly referred to as 'green gold' due to the considerable value derived from its above ground and underground organs. The seeds have nutritional value comparable to legumes such as groundnut and soybean. In addition, the seeds are a rich source of phytochemicals such as phenolic acids, phytosterols, flavonoids, behenic acid and griffonilide while carbohydrates are abundant in the tubers. Based on the existing literature, marama bean remains poorly explored, mainly anecdotal with limited scientific evidence available to support its nutritional and medicinal uses as well as economic benefits. This has been ascribed to a shortage of clear research goals and limited resources specifically directed to this underutilized indigenous plant. From an economic and commercial perspective, the high phytochemical content suggests the possibility of developing a functional health drink and associated value-added products. However, efficient cultivation protocols for marama bean, especially to ensure the sustainable supply of the plant material, remain crucial. Furthermore, novel approaches, especially the use of molecular techniques that can facilitate rapid selection of desired traits in marama, are recommended. These anticipated improved agronomical traits will enhance the commercial and economical potential of marama and also contribute to rural–urban food–nutrition sustainability globally.

Received 24th July 2020,
Accepted 2nd February 2021
DOI: 10.1039/d0fo01937b

rsc.li/food-function

1. Introduction

Legumes are excellent crops for global food and nutrition security, serving as protein, fibre, vitamin and mineral sources.^{1,2} In developing countries where protein–energy malnutrition is often a major nutritional problem, the importance of consuming legumes as a less expensive alternative cannot be over-emphasized.^{3,4} Indigenous legumes have the potential to increase the protein content of diets in developing countries.^{5,6} However, many underutilized indigenous legumes remain undomesticated thereby limiting their potential nutritional and economic value.^{7,8} Exploring the domestication and cultivation of “underutilized plants” including

marama beans (*Tylosema esculentum* (Burch.) A. Schreib) can ameliorate the present dependence on a few legumes that can pose agronomic, nutritional, economic and ecological threats. The success of exploring indigenous plants for nutritional and commercial value lies in their effective domestication, cultivation, preservation and conservation.^{9–11}

Agronomic conditions, nutritional composition, and molecular characteristics need to be determined for marama breeding and domestication.¹² Furthermore, exploring the underutilized marama bean for nutritional, health and economic benefits may contribute to food security and sustainable livelihoods in Africa.¹³ Therefore, a detailed scientific study of marama bean, its nutrients and phytochemical constituents, and breeding approaches for plant improvement as well as environmental sustainability is urgently needed. This review provides a critical appraisal of the botany, taxonomic description, nutritional and phytochemical characteristics and biological properties of marama bean. In addition, the economic prospects of marama bean for household food and nutrition security, income and value addition opportunities are explored.

^aFood Security and Safety Niche Area, Faculty of Natural and Agricultural Sciences, North-West University, Private Bag X2046, Mmabatho 2790, North West Province, South Africa. E-mail: Oladapo.Aremu@nwu.ac.za; Tel: +27 18 389 2573

^bIndigenous Knowledge Systems (IKS) Centre, Faculty of Natural and Agricultural Sciences, North-West University, Private Bag X2046, Mmabatho 2790, North West Province, South Africa



2. Botanical, phenotypic and taxonomic description of marama bean

Marama bean is a perennial legume, a member of the Fabaceae, and its common names include Braiiboontjie Marumama, Gembok beans, Gembokboontjies, Muraki, Tamani berry and Morama. The plant grows sporadically during summer and flourishes in soils with little organic matter, nitrogen, and phosphorus at temperatures of 28–37 °C. In addition, it needs sunshine for 3 h on either side of midday, *ca.* 2000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ light intensity. As a “drought tolerant” plant, it has a unique role in the agroecology where it is cultivated.¹² The stem grows as a vine up to 3 m in length at maturity (Fig. 1a and b). The leaves are bi-lobed and soft/succulent when young but become thick or leathery and greyish at maturity (Fig. 1a). The flowers are approximately 25 mm long, borne in a raceme and yellowish in colour. It has pods that are hardy in nature with 1–3 seeds per pod (Fig. 1b). The seeds in the pods are oval-shaped and tough, and have reddish-brown-black coloration (Fig. 1c).

3. Origin, geographical distribution and the major growing areas of marama bean

Marama bean, also known as “the green gold” is considered a native of the Kalahari Desert of South Africa.¹⁴ It grows natu-

rally in countries such as Namibia and Botswana. The marama bean is undergoing experimental cultivation in Australia, Israel, Kenya and the United States of America (Fig. 2A).¹⁵ This plant is also sparingly cultivated in Mozambique and Zambia. Marama bean is found in the North West, Northern Cape, Limpopo and Gauteng Provinces of South Africa (Fig. 2B), according to the National Department of Agriculture, Forestry and Fisheries.¹⁶

4. Nutritional and phytochemical content in marama bean

Marama is a major source of diverse phytochemical and nutritional compounds. The chemical profile of marama bean and its comparative analysis with other protein rich legumes have been recently reported in the literature.¹⁷ Its nutritional and phytochemical content has a diverse range of uses in industries involved in functional foods, pharmaceuticals and nutraceuticals.¹⁸

4.1. Nutritional composition of marama bean

The matured seeds of marama bean are prepared in different ways such as roasting, boiling or grinding.¹⁹ In addition, the seeds are used for producing oil, butter and porridge. Generally, the seeds are well-preserved when they remain unopened in their pods due to their hard outer cover. Heated and unheated




Fig. 1 Morphology of marama bean (*Tylosema esculentum*): (A) a typical growing marama bean vine, (B) immature pods and (C) harvested dry seeds.






Fig. 2 Geographical locations for marama bean cultivation (A) globally and (B) in South Africa.^{15,16}



marama beans can be made into flour for food.²⁰ Similarly, marama bean milk, a nutritious beverage, resembles soymilk or dairy milk and has other refreshing possibilities.^{21,22} Marama bean has macronutrients that are required in human diet, particularly protein, dietary fibre and healthy fats. The micronutrient (e.g. vitamins and minerals) content is highlighted in Table 1. Marama beans are rich in calcium (241 mg per 100 g), which is essential for human bones and teeth.²³

The beans are a source of magnesium (274.5 mg per 100 g), which serves as a catalyst in biochemical and physiological processes.²⁵ According to Mùseler and Schönfeldt,²³ marama bean contains iron (3.95 mg per 100 g), which is essential for the prevention of anaemia. In addition, marama bean contains zinc (6.2 mg per 100 g), which is useful for protein metabolism and growth. The deficiencies of these micronutrients are common among the rural populations of developing countries and are often caused by the lack of financial means to acquire and consume sufficient quantities of red meat, chicken, fish, fruit, and vegetables.²⁶

The underground part of marama produces a tuber that is larger in size and more nutritious than common potatoes, yams and sugar beets which are common staple tubers.²⁷ The tuber is edible raw, boiled, or roasted but needs to be harvested at a younger stage of the plant's growth (i.e. 1 or 2 years old), when it is nutritious and succulent for consumption.²⁵ The tuber becomes fibrous and bitter to taste, but could serve as an important water reservoir for farmers, hunters and animals in times of emergency. This tuber within a few years of maturity can attain a mass of 12 kg with 90% water content on average.²⁵ Furthermore, the tubers are rich in carbohydrate (11.85 g). Young tubers of marama have a good, succulent, and sweet taste and their texture has been described as similar to that of artichoke (*Cynara cardunculus* var. *scolymus*). Marama tubers are reddish when dried and edible.²⁸ In addition, the immature and mature stems are edible and used for making soup. It is also regarded as particularly good fodder for livestock.²⁵

Table 1 Comparison of the nutrient composition of marama bean (per 100 g dry matter) with that of soybean and groundnut '—' = not done

Class	Nutrient	Marama bean ²³	Soybean ²⁴	Groundnut ²⁴
Proximate	Ash (%)	3.19	4.50	3.80
	Dry matter (%)	96.22	92	95
	Fat (%)	40.06	25	50
	Moisture (%)	2.67	7	9
	Non-structured carbohydrates (g)	11.85	15	20
	Protein (%)	34.71	45	25
Minerals	Calcium (mg)	241	—	—
	Chromium (mg)	0.06	—	—
	Copper (mg)	1.04	—	—
	Iodine (mg)	0.06	—	—
	Iron (mg)	3.95	—	—
	Magnesium (mg)	274.5	—	—
	Zinc (mg)	6.2	—	—
Vitamins	Vitamin B ₁₂ (mg)	0.043	—	—
	Vitamin B ₆ (mg)	1.56	—	—

4.2. Amino acids in marama bean and their functions

Marama beans contain amino acids such as alanine, arginine, aspartic, cysteine, isoleucine and leucine (Table 2). These compounds are building blocks of proteins and play many critical roles in the human body.²³ Amino acids regulate multiple processes with regard to gene expression and modulate proteins that mediate messenger RNA (mRNA) translation.²⁹ Essential amino acid deficiencies lead to a lack of protein synthesis. As shown in Table 3, the major amino acids (composition per 100 g) in marama seed are glutamic acid, aspartic acid and tyrosine acid.^{20,30,31}

4.3. Fatty acids in marama bean and their functions

Marama bean oil is yellow (i.e. golden yellow). The principal fatty acids are: arachidic (3%), essential for synthesis of various hormones, such as prostaglandins, thromboxanes, and leukotrienes;³³ linoleic (19–26%), an important constituent of neuronal membrane phospholipids and a precursor of prostaglandins;³⁴ oleic (48–49%), modifies plasma lipid and lipoprotein concentrations, inhibits coagulation, improves glucose homeostasis, and attenuates inflammation;³⁵ stearic (7–10%), does not increase atherosclerosis risk and is associated with reduced blood pressure and improved heart function;³⁶ and palmitic acid (12–14%), has shown anti-inflammatory and lipid lowering effects linked to the prevention of metabolic syndrome, cardiovascular disease and insulin resistance associated with diabetes and obesity.³⁷ As reported by van der Maesen,²⁸ 100 g dry weight of defatted marama bean seed meal contains 194 kJ (i.e. 46 kcal) of energy, 55 g of protein, 13 g of available starch and 1.6 g of fibre. As shown in Table 3, marama bean contains high levels of oleic acid (42.16% of total fatty acids) and linoleic acid (31.11% of total fatty acids). Furthermore, long-chain unsaturated fats are healthier than saturated fats which have been reported to cause astrocyte activation and brain inflammation associated with neuropathy in metabolic disease such as obesity.³⁸ The fatty acids present in marama bean demonstrate several health benefits.^{23,31}

4.4. Phytochemicals in marama bean and their functions

Marama bean is a potential source of diverse phytochemicals including phenolics, flavonoids, saponins and phytosterols (Table 4 and Fig. 3). These phytochemicals are capable of enhancing the immune system and protecting the body against pathogens.^{42,43} The biological activities of different parts of marama in mitigating the risk of non-communicable diseases, diabetes and some cancers have been explored.³⁹ According to Jackson *et al.* and Bousquet,^{22,44} marama is effective against gastroenteritis by inhibiting enteric bacterial and viral pathogens, fights drug resistant infections and reduces inflammation. Furthermore, Chingwaru *et al.*⁴⁵ identified that marama bean has anti-microbial (anti-bacterial) activities. Marama bean seed coat and cotyledon polyphenolic fractions have relatively high anti-bacterial and anti-fungal activities, implying their possible use as new and potent anti-microbial drugs.⁴⁶ In addition, the cotyledon and seed



Table 2 Overview of amino acids and their functions in nutrition and metabolism as well as the composition in marama bean (per 100 g dry matter) '—' = not available

Amino acid	IUPAC name	Function in nutrition and precursor	Composition per 100 g dry matter ^{20,31}	
Arginine	(2S)-2-Amino-5-(diaminomethylideneamino) pentanoic acid	Antioxidant; regulates hormone secretion; ammonia detoxification; gene expression; immune function; N ₂ reservoir; methylation of proteins; deamination (formation of citrulline) of proteins. Its metabolism produces nitric oxide (NO) a signaling molecule; regulates nutrient metabolism, hemodynamics, angiogenesis, spermatogenesis, embryogenesis, fertility, immune function, hormone secretion, wound healing, neurotransmission, mitochondrial biogenesis ³²	6.3	7.3
Cysteine	(2R)-2-Amino-3-sulfanylpropanoic acid	Disulfide linkage in protein; transport of sulfur, its metabolite taurine: antioxidant; regulates cellular redox state; osmolyte, cellular metabolism and nutrition ³²	0.8	0.1
Histidine	(2S)-2-Amino-3-(1H-imidazol-5-yl) propanoic acid	Hemoproteins (e.g., hemoglobin, myoglobin, catalase, and cytochrome c); production of CO (a signaling molecule); protein methylation; hemoglobin structure and function; antioxidative dipeptides; one-carbon unit metabolism; allergic reaction; vasodilator; central acetylcholine secretion; regulation of gut function; modulation of the immune response in skin ³²	2.4	2.8
Leucine	(2S)-2-Amino-4-methylpentanoic acid	Regulates protein turnover through cellular mTOR signaling and gene expression; activates glutamate dehydrogenase; branched-chain amino acids (BCAA) balance; helps regulate blood sugar levels, stimulates wound healing and produces growth hormones ³²	5.9	6.6
Lysine	(2S)-2,6-Diaminohexanoic acid	Major role in protein synthesis, hormone and enzyme production and calcium absorption. Important for energy production, immune function. ³² Regulates NO synthesis; antiviral activity (treatment of Herpes simplex); protein methylation (e.g., trimethyllysine in calmodulin), acetylation, ubiquitination, and O-linked glycosylation ³²	5.5	5.7
Isoleucine	(2S,3S)-2-Amino-3-methylpentanoic acid	Involved in muscle metabolism; heavily concentrated in muscle tissue. Important for immune function, hemoglobin production and energy regulation ³²	4.0	4.5
Methionine	(2S)-2-Amino-4-methylsulfanylbutanoic acid	Important role in metabolism and detoxification. Necessary for tissue growth and Zn and Se absorption ³²	0.8	0.8
Phenylalanine	(2S)-2-Amino-3-phenylpropanoic acid	Activates tetrahydrobiopterin (BH ₄) (a cofactor for NOS synthesis); tyrosine synthesis; neurological development and function. Precursor for neurotransmitters tyrosine, dopamine, epinephrine and norepinephrine ³²	4.8	4.8
Threonine	(2S,3R)-2-Amino-3-hydroxybutanoic acid	Principal part of structural proteins such as collagen and elastin, important components of skin and connective tissue. Plays a role in fat metabolism and immune function, synthesis of mucin protein required for maintaining intestinal integrity and function; immune function; protein phosphorylation and O-linked glycosylation; glycine synthesis ³²	3.0	3.1
Tryptophan	(2S)-2-Amino-3-(1H-indol-3-yl)propanoic acid	Neurotransmitter; inhibiting production of inflammatory cytokines and superoxide; inhibitor of BH ₄ synthesis; antioxidant; inhibition of the production of inflammatory cytokines and superoxide; antioxidant; inhibition of the production of inflammatory cytokines and superoxide; inhibiting the production of pro inflammatory T-helper-1 cytokines; preventing autoimmune neuro inflammation; enhancing immune function; a component of NAD and NADP, coenzymes for many oxidoreductases ³²	1.7	—
Tyrosine	(2S)-2-Amino-3-(4-hydroxyphenyl) propanoic acid	Protein phosphorylation, nitrosation, and sulfation. Its metabolites include (dopamine, epinephrine (EPN) and norepinephrine (NEPN), melanin) Neurotransmitter; regulates immune response, neurotransmitters; cell metabolism, antioxidant; inhibits the production of inflammatory cytokines and superoxide ³²	3.0	3.1
Valine	(2S)-2-Amino-3-methylbutanoic acid	Important for building and repairing muscle cells and involved in energy production; synthesis of glutamine and alanine; balance among BCAA ³²	4.4	4.9

coat^{39,47} have phenolic acids with anti-inflammatory, anti-hyperglycemic and pro-apoptotic activities.⁴⁸ Marama seed coat extracts have potent anti-oxidant activity and protective effects against free radical induced erythrocyte haemolysis.⁴⁹ The polyphenol rich seed coat and cotyledons of marama

bean interfere with the replicative pathways of rotavirus by reducing virus induced inflammation and the over-expression of nitric oxide.⁵⁰ Therefore, marama has major potential as an anti-microbial, anti-inflammatory, and anti-cancer agent (Table 4).



Table 3 Fatty acids in marama bean^{22,39–41} ‘—’ = not available

Fatty acid	Content (%)	IUPAC name	Chemical structure
Alpha-linolenic acid	—	(9Z,12Z,15Z)-Octadeca 9,12,15-trienoic acid	
Arachidic acid	3.00	Eicosanoic acid (or icosanoic acid)	
Arachidonic acid	0.368	(5Z,8Z,11Z,14Z)-Icosa-5,8,11,14-tetraenoic acid	
Behenic acid	1.28	Docosanoic acid	
Erucic acid	—	(Z)-Docos-13-enoic acid	
Linoleic acid	19–26	(9Z,12Z)-Octadeca-9,12-dienoic acid	
Myristic acid	—	Tetradecanoic acid	
Oleic acid	48–49	(9Z)-Octadec-9-enoic acid	
Palmitic acid	12–14	Hexadecanoic acid	
Palmitoleic acid	0.47	(9Z)-Hexadec-9-enoic acid	
Stearic acid	7–10	Octadecanoic acid	

5. Propagation of marama beans: towards attainment of food and nutrition security

The marama bean grows well in soil with low organic matter and low nutrient content, especially nitrogen. According to the Department of Agriculture, Forestry and Fisheries (South Africa),¹⁶ marama bean prefers a neutral pH (7) soil and does not grow well under waterlogged conditions. The plant grows as a legume scrub, with long stems

that can reach 3 m in length arising from a big underground tuber. It grows well in dry areas with less than 100 mm annual rain and with 250–600 mm optimum rainfall. It can be propagated through seed, in well prepared soil that is soft, well-drained and free of weeds. The seeds are planted (October–November) after being scarified to ensure proper germination.¹⁶

Scarification involves low temperature seed exposure to bring about prompt and uniform germination. Seed treatment is also important to prevent seed, soil and air borne diseases. Scarifying with boiling water or by soaking in water



Table 4 Phytochemicals in different parts of marama bean and their associated biological activities

Class of phytochemical	Compound	IUPAC	Plant part	Biological activity of compound in literature
Phenolic acids	Gallic acid	3,4,5-Trihydroxybenzoic acid	Cotyledon and seed coat ^{22, 39, 47}	Anti-carcinogenic, antimicrobial, anti-mutagenic, anti-angiogenic and anti-inflammatory agents ⁵¹
	Protocatechuic acid	3,4-Dihydroxybenzoic acid	Seed coat, cotyledon and seed coat ^{39, 47}	Anti-inflammatory, anti-hyperglycemic, pro-apoptotic agent ⁴⁸
	<i>p</i> -Hydroxybenzoic acid	4-Hydroxybenzoic acid	Seed coat and cotyledon ^{39, 47}	Anti-oxidant ⁵²
	Vanillic acid	4-Hydroxy-3-methoxybenzoic acid	Seed coat and cotyledon ^{39, 47}	Anti-oxidant ⁵³
	Syringic acid	4-Hydroxy-3,5-dimethoxy-benzoic acid	Seed coat ³⁹	Anti-inflammatory activity ⁵⁴
	Caffeic acid	3,4-Dihydroxycinnamic acid	Seed coat and cotyledon ^{39, 47}	Anti-oxidant ⁵⁵
	<i>p</i> -Coumaric acid	(2 <i>E</i>)-3-(4-Hydroxyphenyl)prop-2-enoic acid	Seed coat ^{39, 47}	Anti-microbial activity ⁵⁶
	Sinapic acid	3-(4-Hydroxy-3,5-dimethoxyphenyl)prop-2-enoic acid	Cotyledon, seed coat ^{39, 47}	Anti-oxidant, anti-inflammatory and anti-mutagenic agent ^{57, 58}
	Ferulic acid	(2 <i>E</i>)-3-(4-Hydroxy-3-methoxyphenyl)prop-2-enoic acid	Cotyledon and seed coat ^{39, 47}	Anti-oxidant, anti-microbial activities ⁵⁹
	Tannic acid	[2,3-Dihydroxy-5-[[[3,4-dihydroxy-5-(3,4,5-trihydroxybenzoyl)oxybenzoyl]oxy]oxan-2-yl]methoxycarbonyl]phenyl]3,4,5-trihydroxybenzoate	Seed ³⁹	Anti-oxidant ⁶⁰
Flavonoids	Fisetin	2-(3,4-Dihydroxyphenyl)-3,7-dihydroxychromen-4-one	Cotyledon and seed coat ^{39, 47}	Neurotrophic, anti-carcinogenic, anti-inflammatory, and other health beneficial effects ⁶¹
	Myricetin	3,5,7-Trihydroxy-2-(3,4,5-trihydroxyphenyl)-4-chromenone	Seed coat ⁴⁷	Anti-oxidative and pro-oxidative properties. It is a potent anti-carcinogen and anti-mutagen ⁶²
	Quercetin	2-(3,4-Dihydroxyphenyl)-3,5,7-trihydroxy-4 <i>H</i> -chromen-4-one	Seed coat ^{39, 47}	Anti-oxidant, anti-inflammatory ⁶³
	Kaempferol	3,5,7-Trihydroxy-2-(4-hydroxyphenyl)-4 <i>H</i> -1-benzopyran-4-one	Seed coat and cotyledon ^{39, 47}	Antioxidant, anti-inflammatory, anticancer ⁶⁴
	Astragalin	5,7-Dihydroxy-2-(4-hydroxyphenyl)-3-[[2 <i>S</i> ,3 <i>R</i> ,4 <i>S</i> ,5 <i>S</i> ,6 <i>R</i>]-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxychromen-4-one	Seed coat, tuber ³⁹	Anti-oxidant, anti-bacterial activities ⁶⁵
	Rutin	2-(3,4-Dihydroxyphenyl)-5,7-dihydroxy-3-[[2 <i>S</i> ,3 <i>R</i> ,4 <i>S</i> ,5 <i>S</i> ,6 <i>R</i>]-3,4,5-trihydroxy-6-[[[2 <i>R</i> ,3 <i>R</i> ,4 <i>R</i> ,5 <i>R</i> ,6 <i>S</i>]-3,4,5-trihydroxy-6-methyloxan-2-yl]oxymethyl]oxan-2-yl]oxychromen-4-one	Seed coat, tuber and cotyledon ^{39, 47}	Antimicrobial, anti-inflammatory, anticancer, and antidiabetic ⁶⁶
	Naringin	(2 <i>S</i>)-7-[[2 <i>S</i> ,3 <i>R</i> ,4 <i>S</i> ,5 <i>S</i> ,6 <i>R</i>]-4,5-dihydroxy-6-(hydroxymethyl)-3-[[2 <i>S</i> ,3 <i>R</i> ,4 <i>R</i> ,5 <i>R</i> ,6 <i>S</i>]-3,4,5-trihydroxy-6-methyloxan-2-yl]oxyoxan-2-yl]oxy-5-hydroxy-2-(4-hydroxyphenyl)-2,3-dihydrochromen-4-one	Tuber, cotyledon, seed coat ^{39, 47}	Anti-oxidant, antitumor, anti-inflammatory ⁶⁷
	Hesperidin	(2 <i>S</i>)-5-Hydroxy-2-(3-hydroxy-4-methoxyphenyl)-7-[[2 <i>S</i> ,3 <i>R</i> ,4 <i>S</i> ,5 <i>S</i> ,6 <i>R</i>]-3,4,5-trihydroxy-6-[[[2 <i>R</i> ,3 <i>R</i> ,4 <i>R</i> ,5 <i>R</i> ,6 <i>S</i>]-3,4,5-trihydroxy-6-methyloxan-2-yl]oxymethyl]oxan-2-yl]oxy-2,3-dihydrochromen-4-one	Tuber, cotyledon, seed coat ^{39, 47}	Anti-inflammatory and analgesic ⁶⁸
	Catechin	(2 <i>R</i> ,3 <i>S</i>)-2-(3,4-Dihydroxyphenyl)-3,4-dihydro-2 <i>H</i> -chromene-3,5,7-triol	Cotyledon ^{39, 47}	Anti-oxidation and free-radical scavenging ⁶⁹
	Benzofuranone	Griffonilide	(6 <i>R</i> ,7 <i>S</i> ,7 <i>aS</i>)-6,7-Dihydroxy-7,7 <i>a</i> -dihydro-6 <i>H</i> -1-benzofuran 2-one	Tuber ⁵⁰
Cyclohexene acetonitrile glycoside	Griffonin (Lithospermoside)	(2 <i>Z</i>)-2-[[[4 <i>R</i> ,5 <i>S</i> ,6 <i>S</i>]-4,5-Dihydroxy-6-[[2 <i>R</i> ,3 <i>R</i> ,4 <i>S</i> ,5 <i>S</i> ,6 <i>R</i>]-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxycyclohex-2-en-1-ylidene]acetonitrile	Tuber ⁵⁰	Antimicrobial ⁵⁰
Phytosterols	<i>Beta</i> -sitosterol	(3 <i>S</i> ,8 <i>S</i> ,9 <i>S</i> ,10 <i>R</i> ,13 <i>R</i> ,14 <i>S</i> ,17 <i>R</i>)-17-[[2 <i>R</i> ,5 <i>R</i>]-5-Ethyl-6-methylheptan-2-yl]-10,13 dimethyl-2,3,4,7,8,9,11,12,14,15,16,17-dodecahydro-1 <i>H</i> -cyclopenta[<i>a</i>]phenanthren-3-ol	Seed ³⁹	Anti-oxidant, anti-inflammatory ⁷¹






Fig. 3 Examples of the chemical structures of some selected phenolic acids, flavonoids and phytosterols in marama bean (*Tylosema esculentum*).

should be avoided as these methods kill the seeds.³⁹ For planting, the seeds are arranged in layers of sand in shallow boxes with pits and trenches to aid rapid germination, a method commonly used for peaches, cherries, oats and grapes.⁷²

The plant needs 5–15 cm planting depth but 20 cm when the soils are wet, with one seed per hole being ideal.¹⁶ Once germinated and growing, it does not need fertilizer or irrigation. However, it needs weeding, as well as pest and disease control from leaf eaters and pod borers which are the most problematic pests of marama beans and can be treated with standard sprays.¹⁶ Harvesting and marketing (April–June) occurs when its pods are brownish in color, usually at the beginning of winter when the seeds are handpicked and the tubers are carefully dug and removed by hand to avoid scarring.¹⁶

6. Novel approaches for selection and improvement of marama bean

There is a need for conscious, efficient and directional improvement of the existing varieties of marama plants to increase the yield. Improvement of its quality, especially in micronutrients, should also be considered. Novel approaches such as next-generation sequencing and *de novo* domestication of marama bean may be pertinent.

6.1. Genetic diversity and next-generation sequencing of marama bean

As described by Takundwa *et al.*,⁷³ marama bean has a relatively small genome of about 1 Gb according to Feulgen staining. However, coverage of the genome in next-generation



sequencing still requires confirmation by flow cytometry.¹² Although the chromosome number of marama bean is 42, there is no proper timing record of the genome duplications.¹² The genomic data were used to identify genes related to various important proteins including cysteine proteases and their inhibitors. It was also used to identify possible genes underlying self-incompatibility (heterostyly) by comparative analysis with the incompatibility genes from other plant species.^{12,74}

Furthermore, genomic resources form the groundwork for developing genetic maps and useful DNA markers for marker-assisted selection (MAS) in marama bean.^{75,76} Early studies have demonstrated a wide diversity among geographically distinct populations of marama bean.^{7,77} This is to be expected for an outcrossing species that is self-incompatible and heterostylous. However, this may be difficult given the heterogeneity in plant varieties and the prevailing environmental situations of the source material available for breeding. Therefore, breeding of marama varieties is also likely to be difficult because the pollinators involved in fertilization and the production of plants and their range remain undefined.¹²

Better elucidation of the reproductive biology of marama beans is of scientific value and essential to accelerate breeding and improvement of the germplasm. The self-incompatibility of marama bean affects agronomic strategies. Compounding this characteristic, marama bean is a hexaploid (containing six homologous sets of chromosomes). Hence, employing molecular marker approaches will be a more complex strategy for identification of not only useful allelic polymorphisms, but also differences in duplicate loci.⁷⁸

As reported by Takundwa *et al.*,⁷⁹ many polymorphisms were identified at the first screening with the aid of simple sequence repeats (SSRs) within a limited germplasm and clear polymorphisms within an individual. The SSRs data confirm earlier studies on wide variability within populations of marama bean. However, there may be specific attributes such as seed number per pod that has a more restricted distribution. There is a need to document the phenotypic as well as molecular diversity in order to select appropriate germplasm for an initial marama bean breeding population.

6.2. Application of *de novo* domestication for marama bean

Technological advancements in wild plant domestication have recently paved the way for *de novo* domestication of indigenous plants as a route for designing ideal crops for food and nutrition security.^{80,81} *De novo* domestication is the introduction of domestication genes into non-domesticated plants; such re-domestication of wild relatives represents an important opportunity for fitting cultivated species to the climatic niche which they habit.⁸² Given the current uncertainties due to climate change, increasing population, and deterioration of arable lands, it is important to consider the production of local food for human sustainability.⁸³

Existing studies have established that in developing countries, about 10–15% of the land is cultivated by traditional methods while about 25 billion individuals cultivate food in

smallholder farms for subsistence.⁸² This indicates a large dependency on small scale farming methods which are unpredictable in terms of sustainable food supply. At the moment, *de novo* domestication application is still scanty but there are examples of how domestication genes were targeted in biotechnological methods that predate the development and widespread adoption of genome-editing techniques.⁸⁰ Two parallel approaches have been suggested for the *de novo* domestication of wild plants: traditional breeding approaches^{84,85} and gene editing.⁸⁶

For gene editing, the CRISPR/Cas9 approach is the method of choice in recent times.^{86–88} This genome-editing tool, which is modified from a prokaryotic immune system, induces double-stranded DNA breaks by the action of Cas9 nuclease at a genome location corresponding to a designed guide RNA.^{82,86} Intriguingly, studies^{82,83,89} have suggested that restarting cultivation of lost plant species may be particularly valuable to the farmers in developing nations since the local species are often already adapted to the local environment. Interestingly, marama bean⁸⁹ has been considered for genome sequencing by the African Orphan Crops Consortium (<http://africanorphancrops.org/>).

7. Challenges and possible ways to enhance the economic potential of marama beans

A number of challenges need to be addressed to foster the economics of marama bean; some of these are highlighted below:

7.1. Demand and supply dichotomy

Most developed countries use leguminous varieties in the production of different foods as increasingly health-conscious populations are looking for more nutritious foods.⁹⁰ In Africa, the quest for low cost but nutritious food (such as marama beans) is urgently needed, especially with increased expectation and demand for existing legumes. Breeding, domestication and commercial cultivation of marama bean could assist in closing the protein food supply gap as well as malnutrition. The lifestyle, tradition and culture of individuals can equally affect the demand and supply chain for the underutilized marama bean.

7.2. Affordability

Currently, legumes are some of the cheapest protein sources in the world.⁹¹ In terms of the environment, indigenous legumes including marama beans are naturally less demanding to cultivate and are economical natural resources. Marama bean can be used in crop rotation, resulting in good yields for subsequent crops, which can in turn improve national food security. However, the affordability of marama bean may depend on the financial capability and economic status of individuals in any particular place.



7.3. Marama domestication, availability and accessibility

The breeding and domestication of marama bean must be encouraged in low and middle income nations, especially in countries where marama bean has comparative advantages

over other legumes. Currently, marama is still mainly from the wild and research is needed on breeding (genome, genetic diversity and next-generation crop) to improve its yield. This will lead to increased availability and accessibility to potential consumers.




Fig. 4 Strategic framework of potential priority areas for policy and intervention on marama bean.



7.4. Awareness and education

Appropriate awareness and education about the benefits associated with marama will ultimately boost its productivity. This can be achieved by researchers, extension agents and other organizations through bulletins, newspapers, magazines, workshops, and training seminars. The cultivation of this plant can uplift rural areas and low-income groups. As most developed nations use legumes for food and because people are becoming increasingly health-conscious, proper awareness and education about marama bean are needed in developing nations for sustainability.⁹⁰

7.5. Marketing opportunities and economic potential of marama bean and its products

Domestication and commercial cultivation of marama beans has the potential to address the shortage in protein food sources in Southern Africa. Increased plant production should be associated with distribution systems and markets for marama products. The expectations of producers, distributors and consumers of marama beans need to be established and strengthened. The South African region has a comparative advantage for producing this underutilized plant, and intensifying efforts in breeding, domesticating and producing more marama beans to alleviate food and nutrition insecurity would lead to significant economic prosperity. Developing countries could gain a competitive advantage by cultivating marama beans for local consumption and export markets. South Africa and other countries in the arid and semi-arid regions should put incentives in place to support local farmers and their markets and encourage the cultivation and marketing of marama beans.

8. Policy and intervention priority areas for the ideal marama bean

In response to the identified challenges associated with the economic potential of marama bean, multidisciplinary research and extensive collaboration among different stakeholders remain critical. As depicted in Fig. 4, a strategic framework that entails a wide range of stakeholders with coordinated efforts is essential to deal with many parts of marama development. Potential and priority areas for marama bean as an alternative protein rich food, as well as a source of nutrition security, were proposed through five (5) interventions in this review and are listed below.

(1) New knowledge generation through genetic mapping of marama bean and scientific research to expand the general knowledge.

(2) Effective transfer of generated data to raise awareness and build capacity amongst stakeholders. This can be facilitated through the use of demonstration sites, targeted campaigns, development of school curricula and training.

(3) Production of convincing evidence-based data that have the potential to influence policy at all levels and remove barriers to marama bean production and marketing.

(4) Improved market development through practical interventions, farmer training, and fostering public-private partnerships at all stages of the value chain to improve the supply and demand for marama and its products. These intervention areas are supported by improved interaction and partnerships amongst all stakeholders

(5) The use of biotechnology to speed up marama bean development through genomics and genome mapping activities, genetic diversity and next-generation sequencing.

In addition, the proposed framework highlights the links between these intervention areas and their expected effects. The following are the key effects:

(a) The worldwide knowledge base will be enhanced by scientific research, mapping the undervalued marama bean and documentation of existing information that are currently inaccessible.

(b) Through targeted communication in a variety of formats, this information will raise the awareness of the general public about the value of underutilized marama bean; relevant information will be fed into curriculum development to inform the present and next generation which will also contribute to capacity building amongst the primary stakeholders.

(c) Two main cases of 'communicating knowledge' have been highlighted. First, food policy has to remove existing barriers such as trade to arrive at appropriate policies with increased public awareness. Second, market development to improve the supply of quality marama bean.

(d) The prerequisites for continuing targeted research and development activities that will use and further increase the global knowledge base include increased capacity among primary stakeholders and a better educated younger generation. This will produce an iterative process that will enable sustainable growth in the production of and demand for marama bean and its products.

(e) Finally, when appropriate policies, increased demand, sustainable production and increased/better marama bean supply coincide, the goal of increased consumption/use of marama bean to address the overall development and food nutrition and economic sustainability problem will be reached.

9. Conclusion and recommendations

Presently, marama bean is mostly restricted to wild population which has resulted in high agronomic and nutritional variability. This review highlighted the nutritional, biological (anti-carcinogenic, antimicrobial, anti-mutagenic, anti-angiogenic and anti-inflammatory properties) and phytochemical potential as well as the economic potential of marama bean. The plant has great potential to contribute to the livelihood of rural communities as well as resource-poor farmers, because the local environmental conditions are appropriate for its cultivation. The challenges confronting marama plant development were discussed with some novel approaches for selection and improvement of nutrition, health and economic sustainability. Particularly, there is a possibility of *de novo* domesti-



cation of marama bean as a viable solution for designing ideal crops while maintaining food security and a more sustainable low-input agriculture. In addition, there is a need for a multidisciplinary research approach that is aimed at efficient breeding and domestication for commercialization. Exploration of the value chain to enhance the commercial and economic value of marama bean is needed as well as the suggested interventions to improve the challenges associated with marama bean consumption. Overall, the review proposed priority areas for policy and intervention and recommends an all-inclusive and sustainable development approach as marama bean could support the attainment of the United Nations Sustainable Development Goals (UN SDG, 2030), in the areas of malnutrition, hunger eradication, economic prosperity and sustainability.

Author contributions

A. O. O. and A. O. A. conceptualized the project. A. O. O. sourced literature and prepared the manuscript with assistance from A. O. A.

Conflicts of interest

The authors of this work declare that there is no conflict of interest.

Acknowledgements

Abiodun Olusola Omotayo is a recipient of a Postdoctoral Fellowship from the North-West University, Mmabatho, South Africa. Adeyemi Oladapo Aremu received research funding from the National Research Foundation (NRF, Incentive Funding for Rated Researchers, Grant UID: 109508). We also thank the Food Security and Safety Niche Area as well as the Higher Degree Committee of the Faculty of Natural and Agricultural Sciences (FNAS), North-West University, Mmabatho, South Africa. We sincerely acknowledge the contribution of Ms Moloko Mojapelo (Department of Agriculture, Forestry and Fisheries Directorate Plant Production: Indigenous Food Crops Division) for her support in providing the photographs of marama bean used in this review article. We appreciate Dr Jerry Adeyemi (Department of Chemistry, North-West University) and Dr Simeon Isaiah (Centre for Human Metabolomics, Department of Biochemistry, North-West University) for assisting in validating the chemical structures and the biochemistry sections of the review. Finally, the English language editing service rendered by Carrin Martin is sincerely appreciated.

References

- O. S. Ijarotimi and O. O. Keshinro, Formulation and nutritional quality of infant formula produced from germinated popcorn, Bambara groundnut and African locust bean flour, *J. Microbiol., Biotechnol. Food Sci.*, 2020, **9**, 1358–1388.
- A. W. Ebert, Potential of underutilized traditional vegetables and legume crops to contribute to food and nutritional security, income and more sustainable production systems, *Sustainability*, 2014, **6**, 319–335.
- N. J. Tonukari and D. G. Omotor, Biotechnology and food security in developing countries, *Biotechnol. Mol. Biol. Rev.*, 2010, **4**, 13–23.
- E. A. Shimelis and S. K. Rakshit, Proximate composition and physico-chemical properties of improved dry bean (*Phaseolus vulgaris* L.) varieties grown in Ethiopia, *LWT-Food Sci. Technol.*, 2005, **38**, 331–338.
- C. Cullis and K. J. Kunert, Unlocking the potential of orphan legumes, *J. Exp. Bot.*, 2017, **68**, 1895–1903.
- M. J. Considine, K. H. Siddique and C. H. Foyer, Nature's pulse power: legumes, food security and climate change, *J. Exp. Bot.*, 2017, **68**, 1815–1818.
- M. L. Hartley, E. Tshamekeng and S. M. Thomas, Functional heterostyly in *Tylosema esculentum* (Caesalpinioideae), *Ann. Bot.*, 2002, **89**, 67–76.
- E. O. Amonsou, J. R. Taylor and A. Minnaar, Adhesive potential of marama bean protein, *Int. J. Adhes. Adhes.*, 2013, **41**, 171–176.
- V. E. Rubatzky and M. Yamaguchi, *World vegetables: principles, production, and nutritive values*, Springer Science & Business Media, 2012.
- M. Faria, E. Mabaya and D. Jordaan, Markets for marama beans in southern Africa: Linking sustainable products with sustainable livelihoods, *Dev. South. Afr.*, 2011, **28**, 477–492.
- A. Nirmalakumari and V. Ulaganathan, Evaluation and identification of suitable finger millet genotypes for higher productivity in combating climate change, in *3rd International conference on neglected*, 2016, pp. 108.
- C. Cullis, D. W. Lawlor, P. Chimwamurombe, N. Bbebe, K. Kunert and J. Vorster, Development of marama bean, an orphan legume, as a crop, *Food Energy Secur.*, 2019, **8**, e00164.
- V. H. Heywood, Overview of agricultural biodiversity and its contribution to nutrition and health, in *Diversifying Food and Diets*, Routledge, 2013, pp. 67–99.
- J. D. Uzabakiriho, Diversity and distribution of *Tylosema esculentum* (Marama bean) endophytic bacteria communities in Omitara, Harnas and Otjinene, eastern Namibia, Doctoral dissertation, University of Namibia, 2016. <http://repository.unam.edu.na/handle/11070/1686>.
- L. Van der Maesen, *Tylosema esculentum* (Burch.) A. Schreib, in *PROTA*, 2006, vol. 1.
- Department of Agriculture Forestry and Fisheries (DAFF), *Production guidelines-Mara bean*, DAFF, Pretoria, South Africa, 2014.
- A. Kuvare, M. Kandawa-Schulz and P. T. Kapewangolo, Comparative study of antioxidant properties, polyphenols and flavonoid contents of the tuber and seed extracts of



- Marama bean (*Tylosema esculentum*), *Int. Sci. Technol. J. Namibia*, 2015, **6**, 21–31.
- 18 M. Holse, S. Husted and Å. Hansen, Chemical composition of marama bean (*Tylosema esculentum*)—A wild African bean with unexploited potential, *J. Food Compos. Anal.*, 2010, **23**, 648–657.
- 19 B. Phuthogo, *Physico-Functional Properties of Wheat-Morama Bean Composite Flour and its Performance in Food Systems*, Master of philosophy degree in food science, University of Ghana, 2014.
- 20 G. N. Maruatona, K. G. Duodu and A. Minnaar, Physicochemical, nutritional and functional properties of marama bean flour, *Food Chem.*, 2010, **121**, 400–405.
- 21 J. C. Jackson, Technology and Nutrition Opportunities for Healthful Foods from Morama Beans, an Emerging Crop in Botswana, in *Global Food Security and Wellness*, Springer, New York, NY, 2017, pp. 125–140.
- 22 J. C. Jackson, K. G. Duodu, M. Holse, M. D. L. de Faria, D. Jordaan, W. Chingwaru, A. Hansen, A. Cencic, M. Kandawa-Schultz, S. M. Mpotokwane, P. Chimwamurombe, H. L. de Kock and A. Minnaar, The Morama Bean (*Tylosema esculentum*): A Potential Crop for Southern Africa, in *Advances in Food and Nutrition Research*, ed. S. L. Taylor, Academic Press, Elsevier, 2010, vol. 61, pp. 187–246.
- 23 D. L. Musesler and H. C. Schönfeldt, The nutrient content of the marama bean (*Tylosema esculentum*), an underutilised legume from Southern Africa, *Agricola*, 2006, **16**, 7–13.
- 24 M. Lokuruka, Soybean nutritional properties: The good and the bad about soy foods consumption—A review, *Afr. J. Food, Agric., Nutr. Dev.*, 2010, **10**, 2439–2459.
- 25 D. L. Musesler, Evaluation of the quality characteristics of the marama bean (*Tylosema esculentum*), an underutilized grain and tuber producing legume in Southern Africa, University of Namibia, 2005.
- 26 A. Iqbal, I. A. Khalil, N. Ateeq and M. S. Khan, Nutritional quality of important food legumes, *Food Chem.*, 2006, **97**, 331–335.
- 27 M. H. Hamunyela, E. Nepolo and M. N. Emmambux, Proximate and starch composition of marama (*Tylosema esculentum*) storage roots during an annual growth period, *S. Afr. J. Sci.*, 2020, **116**, DOI: 10.17159/sajs.2020/6782.
- 28 L. J. G. van der Maesen, *Tylosema esculentum* (Burch.) A. Schreib, in *PROTA 1, Cereals and Pulses*, 2006.
- 29 S. R. Kimball and L. S. Jefferson, New functions for amino acids: effects on gene transcription and translation, *Am. J. Clin. Nutr.*, 2006, **83**, 500S–507S.
- 30 E. Kayitesi, H. L. de Kock, A. Minnaar and K. G. Duodu, Nutritional quality and antioxidant activity of marama-sorghum composite flours and porridges, *Food Chem.*, 2012, **131**, 837–842.
- 31 N. Bower, K. Hertel, J. Oh and R. Storey, Nutritional evaluation of marama bean (*Tylosema esculentum*, Fabaceae): Analysis of the seed, *Econ. Bot.*, 1988, **42**, 533–540.
- 32 G. Wu, Amino acids: metabolism, functions, and nutrition, *Amino Acids*, 2009, **37**, 1–17.
- 33 T. Brody, *Nutritional Biochemistry*, Academic Press, San Diego, 2nd edn, 1998, [https://books.google.co.za/books?hl=en&lr=&id=n2fgyhDUaTEC&oi=fnd&pg=PP1&dq=T.+O.+M.+Brody,+in+Nutritional+Biochemistry+\(Second+Edition\)&ots=Cv4QOXOGZ&sig=q_zUIGjZT6gnuSSgj_wstDOWLTI#v=onepage&q=T.%20O.%20M.%20Brody%2C%20in%20Nutritional%20Biochemistry%20\(Second%20Edition\)&f=false](https://books.google.co.za/books?hl=en&lr=&id=n2fgyhDUaTEC&oi=fnd&pg=PP1&dq=T.+O.+M.+Brody,+in+Nutritional+Biochemistry+(Second+Edition)&ots=Cv4QOXOGZ&sig=q_zUIGjZT6gnuSSgj_wstDOWLTI#v=onepage&q=T.%20O.%20M.%20Brody%2C%20in%20Nutritional%20Biochemistry%20(Second%20Edition)&f=false).
- 34 M. Brownlee, L. P. Aiello, M. E. Cooper, A. I. Vinik, J. Plutzky and A. J. M. Boulton, Complications of diabetes mellitus, in *Williams Textbook of Endocrinology*, ed. S. Melmed, K. S. Polonsky, P. R. Larsen and H. M. Kronenberg, Elsevier, Philadelphia, 13th edn, 2016, pp. 1484–1581, DOI: 10.1016/B978-0-323-29738-7.00033-2.
- 35 S. Lopez, B. Bermudez, Y. M. Pacheco, A. Ortega, L. M. Varela, R. Abia and F. J. G. Muriana, Oleic Acid: The Main Component of Olive Oil on Postprandial Metabolic Processes, in *Olives and Olive Oil in Health and Disease Prevention*, ed. V. R. Preedy and R. R. Watson, Academic Press, San Diego, 2010, pp. 1385–1393. DOI: 10.1016/B978-0-12-374420-3.00154-6.
- 36 D. Senyilmaz-Tiebe, D. H. Pfaff, S. Virtue, K. V. Schwarz, T. Fleming, S. Altamura, M. U. Muckenthaler, J. G. Okun, A. Vidal-Puig, P. Nawroth and A. A. Teleman, Dietary stearic acid regulates mitochondria in vivo in humans, *Nat. Commun.*, 2018, **9**, 3129.
- 37 N. Morse, Are some health benefits of palmitoleic acid supplementation due to its effects on 5' adenosine monophosphate-activated protein kinase (AMPK)?, *Lipid Technol.*, 2015, **27**, 278–281.
- 38 S. Tokuyama and K. Nakamoto, Pain as modified by polyunsaturated fatty acids, in *Omega-3 Fatty Acids in Brain and Neurological Health*, Academic Press, Boston, 2014, pp. 131–146. DOI: 10.1016/B978-0-12-410527-0.00012-0.
- 39 W. Chingwaru, J. Vidmar, P. T. Kapewangolo, O. Mazimba and J. Jackson, Therapeutic and prophylactic potential of morama (*Tylosema esculentum*): a review, *Phytother. Res.*, 2015, **29**, 1423–1438.
- 40 D. L. Musesler and H. Schönfeldt, The nutrient content of the Marama bean (*Tylosema esculentum*), an underutilised legume from Southern Africa, *Agricola*, 2006, **16**, 7–13.
- 41 M. Biesele and R. Murray, in *Alternative food plants for arid regions: Marama and other plant foods of Kalahari foragers, An applied ethno-botanical study. Report to the Center for the Study of Human Adaptation, Division of Graduate Nutrition*, University of Texas, Austin, 1983.
- 42 A. Khan, M. Suleman, S. Abdul Baqi and M. Ayub, 1. Phytochemicals and their role in curing fatal diseases: A Review, *Pure Appl. Biol.*, 2019, **8**, 343–354.
- 43 M. Saxena, J. Saxena, R. Nema, D. Singh and A. Gupta, Phytochemistry of medicinal plants, *J. Pharmacogn. Phytochem.*, 2013, **1**, 168–182.
- 44 J. Bousquet, The morama bean of the Kalahari desert as a potential food crop, with a summary of current research in Texas, *Desert Plants*, 1981, **3**, 213–215.



- 45 W. Chingwaru, G. Duodu, Y. Van Zyl, C. Schoeman, R. T. Majinda, S. O. Yeboah, J. Jackson, P. T. Kapewangolo, M. Kandawa-Schulz and A. Minnaar, Antibacterial and anticandidal activity of *Tylosema esculentum* (marama) extracts, *S. Afr. J. Sci.*, 2011, **107**, DOI: 10.4102/sajs.v107i3/4.366.
- 46 S. O. Yeboah, W. Chingwaru, J. Jackson, Y. Van Zyl, A. Cencic, M. Kandawa-Schulz, R. T. Majinda, G. Duodu, A. Minnaar and P. T. Kapewangolo, Antibacterial and anticandidal activity of *Tylosema esculentum*, (marama) extracts, *S. Afr. J. Sci.*, 2011, **107**, DOI: 10.4102/sajs.v107i3/4.366.
- 47 J. Van Zyl, *Marama bean cotyledon and seed coats: Phenolic composition and antioxidant activity*, University of Pretoria, Pretoria, South Africa B. Sc. Honours Project Report, 2007.
- 48 R. Masella, C. Santangelo, M. D'archivio, G. LiVolti, C. Giovannini and F. Galvano, Protocatechuic acid and human disease prevention: biological activities and molecular mechanisms, *Curr. Med. Chem.*, 2012, **19**, 2901–2917.
- 49 J. S. Shelembe, D. Cromarty, M. J. Bester, A. Minnaar and K. G. Duodu, Characterisation of phenolic acids, flavonoids, proanthocyanidins and antioxidant activity of water extracts from seed coats of marama bean [*Tylosema esculentum*] – an underutilised food legume, *Int. J. Food Sci. Technol.*, 2012, **47**, 648–655.
- 50 O. Mazimba, R. R. Majinda, C. Modibedi, I. B. Masesane, A. Cencič and W. Chingwaru, *Tylosema esculentum* extractives and their bioactivity, *Bioorg. Med. Chem.*, 2011, **19**, 5225–5230.
- 51 S. Choubey, L. R. Varughese, V. Kumar and V. Beniwal, Medicinal importance of gallic acid and its ester derivatives: a patent review, *Pharm. Pat. Anal.*, 2015, **4**, 305–315.
- 52 A. N. Li, S. Li, Y. J. Zhang, X. R. Xu, Y. M. Chen and H. B. Li, Resources and biological activities of natural polyphenols, *Nutrients*, 2014, **6**, 6020–6047.
- 53 P. S. M. Prince, K. Dhanasekar and S. Rajakumar, Vanillic acid prevents altered ion pumps, ions, inhibits Fas-receptor and caspase mediated apoptosis-signaling pathway and cardiomyocyte death in myocardial infarcted rats, *Chem.-Biol. Interact.*, 2015, **232**, 68–76.
- 54 J. R. Ham, H. I. Lee, R. Y. Choi, M. O. Sim, K. I. Seo and M. K. Lee, Anti-steatotic and anti-inflammatory roles of syringic acid in high-fat diet-induced obese mice, *Food Funct.*, 2016, **7**, 689–697.
- 55 J. H. Chen and C.-T. Ho, Antioxidant activities of caffeic acid and its related hydroxycinnamic acid compounds, *J. Agric. Food Chem.*, 1997, **45**, 2374–2378.
- 56 D. Stojković, J. Petrović, M. Soković, J. Glamočlija, J. Kukić-Marković and S. Petrović, In situ antioxidant and antimicrobial activities of naturally occurring caffeic acid, p-coumaric acid and rutin, using food systems, *J. Sci. Food Agric.*, 2013, **93**, 3205–3208.
- 57 N. Ničiforović and H. Abramovič, Sinapic acid and its derivatives: natural sources and bioactivity, *Compr. Rev. Food Sci. Food Saf.*, 2014, **13**, 34–51.
- 58 G. M. Ramolemana, Influence of soil chemical characteristics on the nutritional value of morama (*Tylosema esculentum*) bean seed a potential crop in Botswana, *J. Agric. Sci.*, 2013, **6**, 185–189.
- 59 A. Borges, C. Ferreira, M. J. Saavedra and M. Simoes, Antibacterial activity and mode of action of ferulic and gallic acids against pathogenic bacteria, *Microb. Drug Resist.*, 2013, **19**, 256–265.
- 60 N. S. Khan, A. Ahmad and S. Hadi, Anti-oxidant, pro-oxidant properties of tannic acid and its binding to DNA, *Chem.-Biol. Interact.*, 2000, **125**, 177–189.
- 61 N. Khan, D. N. Syed, N. Ahmad and H. Mukhtar, Fisetin: a dietary antioxidant for health promotion, *Antioxid. Redox Signaling*, 2013, **19**, 151–162.
- 62 K. C. Ong and H. Khoo, Biological effects of myricetin, *Gen. Pharmacol.: Vascul. Syst.*, 1997, **29**, 121–126.
- 63 K. Horvathova, L. Novotny, D. Tothova and A. Vachalkova, Determination of free radical scavenging activity of quercetin, rutin, luteolin and apigenin in H₂O₂-treated human ML cells K562, *Neoplasma*, 2004, **51**, 395–399.
- 64 J. M. Calderon-Montano, E. Burgos-Morón, C. Pérez-Guerrero and M. López-Lázaro, A review on the dietary flavonoid kaempferol, *Mini-Rev. Med. Chem.*, 2011, **11**, 298–344.
- 65 A. Riaz, A. Rasul, G. Hussain, M. K. Zahoor, F. Jabeen, Z. Subhani, T. Younis, M. Ali, I. Sarfraz and Z. Selamoglu, Astragalin: a bioactive phytochemical with potential therapeutic activities, *Adv. Pharmacol. Sci.*, 2018, **2018**, DOI: 10.1155/2018/9794625.
- 66 B. Gullón, T. A. Lú-Chau, M. T. Moreira, J. M. Lema and G. Eibes, Rutin: A review on extraction, identification and purification methods, biological activities and approaches to enhance its bioavailability, *Trends Food Sci. Technol.*, 2017, **67**, 220–235.
- 67 M. Cavia-Saiz, M. D. Busto, M. C. Pilar-Izquierdo, N. Ortega, M. Perez-Mateos and P. Muñiz, Antioxidant properties, radical scavenging activity and biomolecule protection capacity of flavonoid naringenin and its glycoside naringin: a comparative study, *J. Sci. Food Agric.*, 2010, **90**, 1238–1244.
- 68 E. Galati, M. Monforte, S. Kirjavainen, A. Forestieri, A. Trovato and M. Tripodo, Biological effects of hesperidin, a citrus flavonoid.(Note I): antiinflammatory and analgesic activity, *Farmaco*, 1994, **40**, 709–712.
- 69 A. M. Haque, M. Hashimoto, M. Katakura, Y. Tanabe, Y. Hara and O. Shido, Long-term administration of green tea catechins improves spatial cognition learning ability in rats, *J. Nutr.*, 2006, **136**, 1043–1047.
- 70 S. Okwute, G. Ndukwe, K. Watanabe and N. Ohno, Isolation of griffonilide from the stem bark of *Bauhinia thonningii*, *J. Nat. Prod.*, 1986, **49**, 716–717.
- 71 N.-R. Han, H.-M. Kim and H. Jeong, The β -sitosterol attenuates atopic dermatitis-like skin lesions through down-regulation of TSLP, *Exp. Biol. Med.*, 2014, **239**, 454–464.
- 72 M. G. Kains, *Propagation Of Plants-A Complete Guide For Professional And Amateur Growers Of Plants By Seeds, Layers, Grafting And Budding, With Chapters On Nursery And Greenhouse Management*, Read Books Ltd, 2020.



- 73 M. Takundwa, P. Chimwamurombe and C. Cullis, A chromosome count in marama bean (*Tylosema esculentum*) by Feulgen staining using garden pea (*Pisum sativum*, L.) as a standard, *Res. J. Biol.*, 2012, **2**, 177–181.
- 74 S. C. Barrett and M. B. Cruzan, Incompatibility in heterostylous plants, in *Genetic control of self-incompatibility and reproductive development in flowering plants*, Springer, 1994, pp. 189–219.
- 75 K.-J. Kim, K.-S. Choi and R. K. Jansen, Two chloroplast DNA inversions originated simultaneously during the early evolution of the sunflower family (Asteraceae), *Mol. Biol. Evol.*, 2005, **22**, 1783–1792.
- 76 Y. Kim and C. Cullis, A novel inversion in the chloroplast genome of marama (*Tylosema esculentum*), *J. Exp. Bot.*, 2017, **68**, 2065–2072.
- 77 M. Takundwa, P. Chimwamurombe, K. Kunert and C. Cullis, Developing DNA barcoding (matK) primers for marama bean [*Tylosema esculentum* (Burchell) Schreiber], *Afr. J. Biotechnol.*, 2012, **11**, 16305–16312.
- 78 D. X. Zhang and G. M. Hewitt, Nuclear DNA analyses in genetic studies of populations: practice, problems and prospects, *Mol. Ecol.*, 2003, **12**, 563–584.
- 79 M. Takundwa, P. M. Chimwamurombe, K. J. Kunert and C. A. Cullis, Isolation and characterization of microsatellite repeats in Marama bean (*Tylosema esculentum*), *Afr. J. Agric. Res.*, 2010, **5**, 561–566.
- 80 J. T. Østerberg, W. Xiang, L. I. Olsen, A. K. Edenbrandt, S. E. Vedel, A. Christiansen, X. Landes, M. M. Andersen, P. Pagh and P. Sandøe, Accelerating the domestication of new crops: feasibility and approaches, *Trends Plant Sci.*, 2017, **22**, 373–384.
- 81 I. K. Dawson, W. Powell, P. Hendre, J. Bančič, J. M. Hickey, R. Kindt, S. Hoad, I. Hale and R. Jamnadass, The role of genetics in mainstreaming the production of new and orphan crops to diversify food systems and support human nutrition, *New Phytol.*, 2019, **224**, 37–54.
- 82 A. R. Fernie and J. Yan, *De novo* domestication: an alternative route toward new crops for the future, *Mol. Plant*, 2019, **12**, 615–631.
- 83 O. Shelef, P. J. Weisberg and F. D. Provenza, The value of native plants and local production in an era of global agriculture, *Front. Plant Sci.*, 2017, **8**, DOI: 10.3389/fpls.2017.02069.
- 84 B. C. Runck, M. B. Kantar, N. R. Jordan, J. A. Anderson, D. L. Wyse, J. O. Eckberg, R. J. Barnes, C. L. Lehman, L. R. DeHaan and R. M. Stupar, The reflective plant breeding paradigm: A robust system of germplasm development to support strategic diversification of agroecosystems, *Crop Sci.*, 2014, **54**, 1939–1948.
- 85 L. R. DeHaan, D. L. Van Tassel, J. A. Anderson, S. R. Asselin, R. Barnes, G. J. Baute, D. J. Cattani, S. W. Culman, K. M. Dorn and B. S. Hulke, A pipeline strategy for grain crop domestication, *Crop Sci.*, 2016, **56**, 917–930.
- 86 F. Altpeter, N. M. Springer, L. E. Bartley, A. E. Blechl, T. P. Brutnell, V. Citovsky, L. J. Conrad, S. B. Gelvin, D. P. Jackson and A. P. Kausch, Advancing crop transformation in the era of genome editing, *Plant Cell*, 2016, **28**, 1510–1520.
- 87 M. Pacher and H. Puchta, From classical mutagenesis to nuclease-based breeding—directing natural DNA repair for a natural end-product, *Plant J.*, 2017, **90**, 819–833.
- 88 A. Scheben, F. Wolter, J. Batley, H. Puchta and D. Edwards, Towards CRISPR/Cas crops—bringing together genomics and genome editing, *New Phytol.*, 2017, **216**, 682–698.
- 89 P. Smýkal, M. N. Nelson, J. D. Berger and E. J. Von Wettberg, The impact of genetic changes during crop domestication, *Agronomy*, 2018, **8**, DOI: 10.3390/agronomy8070119.
- 90 A. Lartey, J. Meerman and R. Wijesinha-Bettoni, Why food system transformation is essential and how nutrition scientists can contribute, *Ann. Nutr. Metab.*, 2018, **72**, 193–201.
- 91 N. Alexandratos and J. Bruinsma, *World agriculture towards 2030/2050: the 2012 revision*, ESA Working paper FAO, Rome, 2012.

