

CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#)



Cite this: *Anal. Methods*, 2022, **14**, 3355

DOI: 10.1039/d2ay90110b

rsc.li/methods

Correction: Novel ^{90}Sr analysis of environmental samples by Ion-Laser InterAction Mass Spectrometry

Maki Honda,^{*ab} Martin Martschini,^a Oscar Marchhart,^a Alfred Priller,^a Peter Steier,^a Robin Golser,^a Tetsuya K. Sato,^b Tsukada Kazuaki^b and Aya Sakaguchi^c

Correction for 'Novel ^{90}Sr analysis of environmental samples by Ion-Laser InterAction Mass Spectrometry' by Maki Honda *et al.*, *Anal. Methods*, 2022, **14**, 2732–2738, <https://doi.org/10.1039/D2AY00604A>.

In the original article, in the sentence in the introduction beginning “Previous studies have reported LODs of <1 Bq...”, the number of atoms approximately equal to 1 Bq was incorrect. The corrected sentence is given below.

Previous studies have reported LODs of <1 Bq ($1 \text{ Bq} \approx 1.3 \times 10^9$ atoms) by ICP-MS/MS,^{8,9} 4 mBq by RIMS,¹¹ and <1 mBq by TIMS.^{12,13}

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aFaculty of Physics, Isotope Physics, University of Vienna, Währinger Strasse 17, Vienna 1090, Austria. E-mail: honda.maki@jaea.go.jp

^bJapan Atomic Energy Agency, 2-4 Shirakata, Tokai, Ibaraki 319-1195, Japan

^cCenter for Research in Isotopes and Environmental Dynamics, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8577, Japan

