


 Cite this: *Chem. Commun.*, 2022, 58, 1546

 Received 3rd December 2021,  
 Accepted 23rd December 2021

DOI: 10.1039/d1cc06800h

rsc.li/chemcomm

# Wavelength dependent photoextrusion and tandem photo-extrusion reactions of ninhydrin bis-acetals for the synthesis of 8-ring lactones, benzocyclobutenes and orthoanhydrides†


 Wei Sun,<sup>a</sup> Surajit Kayal,<sup>b</sup> William A. T. Raimbach,<sup>a</sup> Xue-Zhong Sun,<sup>b</sup> Mark E. Light,<sup>a</sup> Magnus W. D. Hanson-Heine,<sup>b</sup> Michael W. George<sup>b,c</sup> and David C. Harrowven<sup>b,\*a</sup>

**Ninhydrin bis-acetals give access to 8-ring lactones, benzocyclobutenes and spirocyclic orthoanhydrides through photoextrusion and tandem photoextrusion reactions. Syntheses of fimbriallyxlactone B, isosihunine and numerous biologically-relevant heterocycles show the value of the methods, while TA-spectroscopy and TD-DFT studies provide mechanistic insights on their wavelength dependence.**

The photodecarbonylation of ninhydrin bis-acetal **2** to benzocyclobutene **3** was developed by Butenschön *et al.* to provide low cost access to benzocyclobutenedione (Scheme 1).<sup>1</sup> Indeed, it remains the most convenient method for its synthesis in spite of the modest yield attained for the photoextrusion of carbon monoxide from **2**. Consequently, when a need for benzocyclobutenedione arose, we decided to develop the reaction as a flow photochemical process in order to facilitate its optimisation with respect to wavelength, residence time, solvent and acetal protecting group.<sup>2,3</sup> In the event, we uncovered conditions leading to a modest yield improvement for benzocyclobutene **3** and discovered a wavelength dependence for the reaction giving high-yielding access to 8-ring lactone **5a** and orthoanhydride **6**. Herein we describe our development of those methods, together with demonstrations of their usefulness in heterocyclic and natural products synthesis, and a detailed mechanistic study by transient absorption (TA) spectroscopy and TD-DFT analysis.

Our investigation began with an examination of the wavelength dependence on the photoextrusion of carbon monoxide

from bis-acetal **2**. While all wavelengths studied led to the evolution of gas, which was vented from the system using a circulating flow technique (see ESI†), only UVA irradiation gave benzocyclobutene **3** as a significant product.<sup>3,4</sup> An optimum yield of 57% was realised when a 0.05 M solution of **2** in acetonitrile was irradiated with  $30 \times 1.5$  W UVA LEDs for residence time of 3 h. By contrast, irradiation of bis-acetal **2** with UVB or UVC light gave rise to a complex mixture of products on work-up that varied from experiment to experiment and contained no benzocyclobutene **3**, *vide infra*. Curiously, when the same reaction was performed on the homologous ninhydrin bis-acetal **4a**, 8-ring lactone **5a** was given in high yield (Scheme 1). Various analogues of **5** carrying substituents on the acetals and arene also gave the reaction in high yield (Table 1). In general, bis-1,3-dioxanes with a symmetrical arene and/or acetal substitution pattern gave a single product in 80–90% yield, *e.g.* **5a–c**. Substrates lacking those symmetry elements also gave high yields but led to isomeric mixtures, *e.g.* lactones **5d–g**. Thought this limited the yield attained for individual products to ~40%, in the context of medium ring synthesis such an outcome remains competitive.<sup>5</sup>



**Scheme 1** The discovery of wavelength and acetal dependence on the outcome of photoextrusion reactions involving ninhydrin bis-acetals.

<sup>a</sup> Chemistry, University of Southampton, Highfield, Southampton, SO17 1BJ, UK.  
 E-mail: dch2@soton.ac.uk


<sup>b</sup> School of Chemistry, University of Nottingham, Nottingham NG7 2RD, UK

<sup>c</sup> Department of Chemical and Environmental Engineering, University of Nottingham Ningbo China, 199 Taikang East Road, Ningbo 315100, China


† Electronic supplementary information (ESI) available: Experimental accounts with spectral details and copies of NMR spectra are available as supplementary information. CCDC 1908866. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1cc06800h



Table 1 Extensions to other ninhydrin bis-acetals



To demonstrate the utility of the method, we next applied it in a total synthesis of fimbriallyxlactone **B 11**, a natural product from the roots of *Strophoblachia fimbriallyx* Boerl. (Euphorbiaceae) (Scheme 2).<sup>6</sup> Thus, bis-acetal **4h** was formed from the corresponding 1-indanone **7** in 76% yield through sequential SeO<sub>2</sub> oxidation and acetalisation with 2,2-dimethylpropanediol.<sup>7</sup> Exposure of an acetonitrile solution of **4h** to UVC light under circulating flow next gave a 3 : 2 mixture of lactones **5h** and **5h'** that was readily separated by column chromatography. Methanolysis of the major isomer **5h** then gave ester **9** which was advanced to fimbriallyxlactone **B 11** by ruthenium tetroxide oxidation to carboxylic acid **10** and acetal hydrolysis with HCl.<sup>8</sup> Importantly, the spectral and physical data recorded on our sample matched those reported for the natural product. Its identity was also confirmed by X-ray crystallographic analysis (Scheme 2, CCDC 1908866).<sup>†,6,9</sup>





Scheme 2 Total synthesis and X-ray crystal structure of fimbriallyxlactone B.

The potential of these 8-ring lactones to serve as intermediates in heterocyclic synthesis was next demonstrated with the conversion of **5a** to spirolactone **12** on acetal hydrolysis (Scheme 3).<sup>9</sup> In turn, its treatment with methylamine gave isoshihunine **13** in 77% yield, from which the alkaloids shihunidine **14** and shihunine **15** can each be derived.<sup>10,11</sup> Furthermore, reaction with hydrazines gave high yielding access to phthalazones (e.g. **16**, **17**, **21**), which are important pharmacophores present in the chemotherapeutic agent olaparib (Lynparza<sup>®</sup>),<sup>12</sup> and azelastine (Astelin, Optivar, Allergodil), a frontline drug treatment for mild rhinitis.<sup>13</sup> By contrast, *o*-phenylenediamine, ethanolamine and ethylenediamine gave lactols **18** – **20** respectively.

The mechanistic course of the photoextrusion reaction was next studied by transient absorption (TA) spectroscopy following excitation of bis-acetal **4a** in CH<sub>3</sub>CN at 266 nm (Scheme 4 and Fig. S1, ESI<sup>†</sup>). Analysis of the ps-TA spectra (Fig. S1a, ESI<sup>†</sup>) allowed us to extract the time constants associated with spectral evolution using global kinetic analysis.<sup>14</sup> The data could be fitted using two exponential decays with time constants  $\tau_1 = 1.8 \pm 0.4$  ps and  $\tau_2 = 1.8 \pm 0.2$  ns (Fig. S1b and c, ESI<sup>†</sup>). The observed transient spectrum at 1 ps time delay showed an absorption band at 585 nm which was assigned to the  $n\pi^*$  ( $S_1$ ) excited state <sup>1</sup>[**4a**] following comparison of the extracted spectrum with that computed for the  $S_1$  state (Fig. S7, ESI<sup>†</sup>). Decay of the  $S_1$  state was associated with the formation of a new transient species with an absorption band at 454 nm. This was attributed to biradical <sup>1</sup>[**22**] based on comparison with results reported for the photoexcitation of 1,1,3-triphenyl-3-hydroxy-2-indanone.<sup>15</sup> Additionally, the result aligns with a recent TA study of the C–C bond cleavage of cyclopentanone in cyclohexane following 255 nm photoexcitation, which Kao *et al.* found occurred in  $\leq 1$  ps.<sup>16</sup> Likewise the  $\tau_2$  component [ $1.8 \pm 0.2$  ns] for the rate of formation of <sup>3</sup>[**22**] from <sup>1</sup>[**22**] is consistent with previously reported ISC timescales for acyl-alkyl diradicals.<sup>17</sup>

Spectral evolution by ns-TA spectroscopy was also monitored (Fig. S1d, ESI<sup>†</sup>). Three exponentials are required to describe these data, with time constants  $\tau_1 = 4.8 \pm 0.4$  ns,  $\tau_2 = 100 \pm 20$  ns, and

Scheme 3 Heterocycles derived from 8-ring lactone **5a**.




**Scheme 4** Proposed mechanistic course of the photoextrusion reactions of bis-acetal **4a** [UVA (broad), 355–390 nm; UVB (narrow), 310 nm; UVC (narrow), 254 nm]

$\tau_3$  = long-lived (Fig. S1e and f, ESI<sup>†</sup>).<sup>17</sup> The  $\tau_1$  component was associated with a decay of the 460 nm band for  $^3[22]$  with simultaneous formation of a new transient at 550 nm, attributed to the decarbonylation product  $^3[23]$ . A similar transient spectral feature was reported for the decarbonylation of diphenylmethylacyl radical ( $\text{Ph}_2\text{MeCCO}^\bullet$ ) in isooctane, which had a rate constant  $k = 1.5 \times 10^8 \text{ s}^{-1}$ .<sup>15,18</sup> Decay of the 550 nm transient species  $^3[23]$ , with  $\tau_2 = 100 \pm 20 \text{ ns}$ , was associated with the formation of a long-lived transient at 440 nm assigned to quinodimethane **26**. This is in line with the rate constant measured by Fujiwara *et al.* for the formation of *o*-xylylene from 2-indanone following photoexcitation at 266 nm in cyclohexane ( $k = 1.4 \pm 0.1 \times 10^7 \text{ s}^{-1}$ ).<sup>19</sup> Thus, the wavelength dependence of the reaction centres on the fate of long-lived intermediate **26** (Scheme 4). Irradiation of **26** with UVA leads to a singlet excited state  $^1[26]$  that can relax to benzocyclobutene **24**. By contrast, its irradiation with UVB or UVC light gives rise to higher singlet excited states of **26** from which ISC to  $^3[23]$  is allowed. Collapse of  $^3[23]$  to biradical **25** then provides access to 8-ring lactone **5a**.

Applying this mechanistic understanding to ninhydrin bis-acetal **2** (Scheme 5) led us to conclude that the course of that reaction was also dictated by the nature of the excitation of quinodimethane intermediate **28**. As before, its irradiation with UVA gives a singlet excited state  $^1[28]$  from which benzocyclobutene **3** is derived. Similarly, irradiation with UVB or UVC light leads to biradical  $^3[29]$ , which in turn collapses to biradical **30**. However, in this case, ring closure to  $\epsilon$ -lactone **27** is outpaced by the extrusion of ethene to biradical **31**, a precursor of orthoanhydride **6**. Indeed, the near quantitative formation of orthoanhydride **6** on photolysis of bis-acetal **2** with UVB or UVC light was confirmed by analysis of a concentrated product mixture before aqueous work-up (see ESI<sup>†</sup> for characterisation data).

A TD-DFT analysis of quinodimethane **28** provided additional support for the mechanism (Fig. 1).<sup>20</sup> In particular, it indicated that the energy provided by UVA irradiation ( $3.18 \text{ eV} \equiv \lambda_{\text{max}} 390 \text{ nm}$ ) mirrored that required for the  $\pi \rightarrow \pi^*$  transition  $S_0 \rightarrow S_1$ , and that ISC between  $S_1$  of  $^1[28]$  and  $T_1$  of  $^3[29]$  was forbidden as both had  $\pi\pi^*$  character.<sup>21</sup> Consequently,




**Scheme 5** Proposed mechanism for the photoextrusion reactions of bis-acetal **2** [UVA (broad), 355–390 nm; UVB (narrow), 310 nm; UVC (narrow), 254 nm].

irradiation of **28** with UVA induces cyclisation to benzocyclobutene **3**. By contrast, the greater energy provided by UVB and UVC irradiation facilitates ISC from  $^1[28]$  to  $^3[29]$  via higher excited states where both the orbital symmetry and energy requirements can each be fulfilled.<sup>21</sup> As a result, irradiation of **28** with UVB or UVC gives orthoanhydride **6**.


Finally, the use of orthoanhydride **6** as a phthalic anhydride surrogate (Scheme 6) was demonstrated with the phthalimide protection of various 1°-amines under mild conditions leading to **30a–f**.<sup>22</sup> Similarly, reactions with 2°-amines gave the expected amides, *e.g.* **31** and **37**, while bis-nucleophiles provided high yielding access to an array of biologically relevant nitrogen heterocycles **32–36**, **38** and **39**.

In conclusion, we have shown that the photodecarbonylation of ninhydrin bis-acetals is a wavelength dependent reaction that can be conducted on a gram scale to give benzocyclobutenes on UVA irradiation (60 W, ~5 h), and orthoanhydrides or 8-ring lactones on UVB or UVC irradiation (36 W, ~2 h). Notably, they formally constitute ring contraction, ring correlative, and ring expansion reactions, respectively. These versatile intermediates give access to a host of heterocycles, natural products, and pharmacophores. Mechanistic details have been delineated using computational and spectroscopic methods, including picosecond and nanosecond



**Fig. 1** TD-DFT calculated energies for the singlet and triplet excited states of **28** and **29** respectively, and the mechanistic implications.





Scheme 6 Illustrations of the synthetic utility of orthoanhydride 1.

transient absorption spectroscopy. We are currently exploring further applications of the chemistry in target-oriented and natural products total synthesis.

Wei Sun and William Raimbach performed the synthetic chemistry under the supervision of David Harrowven. Wei Sun and Magnus Hanson-Heine performed the DFT analyses while Mark Light conducted the X-ray analysis. Transient absorption spectroscopic analyses were by Surajit Kayal and Xue-Zhong Sun under the supervision of Michael George.

We gratefully acknowledge financial support from EPSRC [EP/P013341/1, EP/L003325/1 and EP/K039466/1], the European Regional Development Fund [ERDF Interreg Va programme (Project 121)], and the University of Nottingham Green Chemicals Beacon.

## Conflicts of interest

There are no conflicts of interest to declare.

## Notes and references

- D. Leinweber, M. Schnebel, R. Wartchow, H. G. Wey and H. Butenschoen, *Eur. J. Org. Chem.*, 2002, 2385; D. Leinweber and H. Butenschoen, *Tetrahedron Lett.*, 1997, 38, 6385; D. Ng, Z. Yang and M. A. Garcia-Garibay, *Tetrahedron Lett.*, 2002, 43, 7063.
- J. P. Knowles, L. D. Elliott and K. I. Booker-Milburn, *Beilstein J. Org. Chem.*, 2012, 8, 2025; K. Gilmore and P. H. Seeberger, *Chem. Rec.*, 2014, 14, 410; C. Sambiagio and T. Noël, *Trends Chem.*, 2020, 2, 92; M. B. Plutschack, B. Pieber, K. Gilmore and P. H. Seeberger, *Chem. Rev.*, 2017, 117, 11796; D. Cambié, C. Bottecchia, N. J. W. Straathof, V. Hessel and T. Noël, *Chem. Rev.*, 2016, 116, 10276; F. Politano and G. Oksdath-Mansilla, *Org. Process Res. Dev.*, 2018, 22, 1045; T. H. Rehm, *ChemPhotoChem*, 2019, 3, 1;

- N. Hoffmann, *Chem. Rev.*, 2008, 108, 1052; M. Oelgemoeller, *Chem. Eng. Technol.*, 2012, 35, 1144.
- D. C. Harrowven, M. Mohamed, T. P. Gonçalves, R. J. Whitby, D. Bolien and H. F. Sneddon, *Angew. Chem., Int. Ed.*, 2012, 51, 4405; T. P. Gonçalves, M. Mohamed, R. J. Whitby, H. F. Sneddon and D. C. Harrowven, *Angew. Chem., Int. Ed.*, 2015, 54, 4531; D. E. Collin, E. H. Jackman, N. Jouandon, W. Sun, D. C. Harrowven and B. Linclau, *Synthesis*, 2021, 1307; M. A. Manning, W. Sun, M. E. Light and D. C. Harrowven, *Chem. Commun.*, 2021, 57, 4556; W. Sun, W. A. T. Raimbach, L. D. Elliott, K. I. Booker-Milburn and D. C. Harrowven, *Chem. Commun.*, 2022, 58, 383.
- B. D. Hook, W. Dohle, P. R. Hirst, M. Pickworth, M. B. Berry and K. I. Booker-Milburn, *J. Org. Chem.*, 2005, 70, 7558; L. D. Elliott, J. P. Knowles, P. J. Koovits, K. G. Maskill, M. J. Robertson-Ralph, G. Lejeune, L. J. Edwards, R. I. Robinson, I. R. Clemens, B. Cox, D. D. Pascoe, G. Koch, M. Eberle, M. B. Berry and K. I. Booker-Milburn, *Chem. – Eur. J.*, 2014, 20, 15226.
- For our previous work on medium-ring synthesis see: W. Sun, D. C. Wilson, M. E. Light and D. C. Harrowven, *Org. Lett.*, 2018, 20, 4346; D. C. Harrowven, N. L'Helias, J. D. Moseley, N. J. Blumire and S. R. Flanagan, *Chem. Commun.*, 2003, 2658.
- P. Seephonkai, S. G. Pyne, A. C. Willis and W. Lie, *J. Nat. Prod.*, 2013, 76, 1358.
- F. Johnson and N. Oberlender, *J. Med. Chem.*, 1991, 34, 1662; L. F. Fieser and W. C. Lothrop, *J. Am. Chem. Soc.*, 1936, 58, 2050.
- H. Niwa, M. Nisiwaki, I. Tsukada, T. Ishigaki, S. Ito, K. Wakamatsu, T. Mori, M. Ishigaki and K. Yamada, *J. Am. Chem. Soc.*, 1990, 112, 9001; B. S. Lucas, L. M. Luther and S. D. Burke, *Org. Lett.*, 2004, 6, 2965.
- Some discrepancies were noted in the  $^{13}\text{C}$  NMR data. These data were reported tentatively in the isolation paper as they had been attained from a spectrum recorded on a 9:1 mixture of fimbriallyxylactones A and B.
- P. M. Hare, C. T. Middleton, K. I. Mertel, J. M. Herbert and B. Kohler, *Chem. Phys.*, 2008, 347, 383; H. Sun and H. Frei, *J. Phys. Chem. B*, 1997, 101, 205.
- V. G. Ore, M. D. Hordia and N. S. Arasimhan, *Tetrahedron*, 1990, 46, 2483; G. A. Flynn, *Chem. Commun.*, 1980, 862; E. Breuer and S. Zbaida, *Tetrahedron*, 1975, 31, 499.
- X.-W. Li, M. Huang, K. Lo, W. L. Chen, Y.-Y. He, Y. Xu, H. Zheng, H. Hu and J. Wang, *Molecules*, 2019, 24, 2673; H. Chen, X. Li, Y. Xu, K. Lo, H. Zheng, H. Hu, J. Wang and Y. Lin, *Molecules*, 2018, 23, 1185; J. Bastida, M. Selles, C. Codina, F. Viladomat and J. L. Leon de la Luz, *Planta Med.*, 1996, 62, 575; G. B. Bodem and E. Leete, *J. Org. Chem.*, 1979, 44, 4696; E. Leete and G. B. Bodem, *J. Am. Chem. Soc.*, 1976, 98, 6321; E. Breuer and S. Zbaida, *Tetrahedron*, 1975, 31, 499; Y. Inubushi, Y. Tsuda, T. Konita and S. Matsumoto, *Chem. Pharm. Bull.*, 1964, 12, 749.
- P. S. Fong, D. S. Boss, T. A. Yap, A. Tutt, P. Wu, M. Mergui-Roelvink, P. Mortimer, H. Swaisland, A. Lau, M. J. O'Connor, A. Ashworth, J. Carmichael, S. B. Kaye, J. H. Schellens and J. S. de Bono, *N. Engl. J. Med.*, 2009, 361, 123.
- The results were analysed with Glotaran (<http://glotaran.org>).
- G. Kuzmanich, J. Xue, J. C. Netto-Ferreira, J. C. Scaiano, M. Platz and M. A. Garcia-Garibay, *Chem. Sci.*, 2011, 2, 1497.
- M. H. Kao, R. K. Venkatraman, M. N. Ashfold and A. J. Orr-Ewing, *Chem. Sci.*, 2020, 11, 1991.
- Y. P. Tsentlovich, O. B. Morozova, N. I. Avdievich, G. S. Ananchenko, A. V. Yurkovskaya, J. D. Ball and M. D. Forbes, *J. Phys. Chem. A*, 1997, 101, 8809.
- C. Chatgililoglu, D. Crich, M. Komatsu and I. Ryu, *Chem. Rev.*, 1999, 99, 1991.
- M. Fujiwara, K. Mishima, K. Tamai, Y. Tanimoto, K. Mizuno and Y. Ishii, *J. Phys. Chem. A*, 1997, 101, 4912.
- A. Drew and M. Head-Gordon, *Chem. Rev.*, 2005, 105, 4009.
- M. A. El-Sayed, *J. Chem. Phys.*, 1963, 38, 2834.
- V. V. Sureshbabu, N. Narendra, in *Amino Acids, Peptides and Proteins in Organic Chemistry: Protection Reactions, Medicinal Chemistry, Combinatorial Synthesis*, ed.: A. B. Hughes, Wiley-VCH, Weinheim, 2011, vol. 4, pp. 1–97.

