



Cite this: *Chem. Commun.*, 2022, **58**, 1546

Received 3rd December 2021,
Accepted 23rd December 2021

DOI: 10.1039/d1cc06800h
rsc.li/chemcomm

Wavelength dependent photoextrusion and tandem photo-extrusion reactions of ninhydrin bis-acetals for the synthesis of 8-ring lactones, benzocyclobutenes and orthoanhydrides†


Wei Sun,^a Surajit Kayal,^b William A. T. Rimbach,^a Xue-Zhong Sun,^b Mark E. Light,^a Magnus W. D. Hanson-Heine,^b Michael W. George^b and David C. Harrowven^a  

Ninhydrin bis-acetals give access to 8-ring lactones, benzocyclobutenes and spirocyclic orthoanhydrides through photoextrusion and tandem photoextrusion reactions. Syntheses of fimbrialcalyx-lactone B, isoshishunine and numerous biologically-relevant heterocycles show the value of the methods, while TA-spectroscopy and TD-DFT studies provide mechanistic insights on their wavelength dependence.

The photodecarbonylation of ninhydrin bis-acetal **2** to benzocyclobutene **3** was developed by Butenschön *et al.* to provide low cost access to benzocyclobutenedione (Scheme 1).¹ Indeed, it remains the most convenient method for its synthesis in spite of the modest yield attained for the photoextrusion of carbon monoxide from **2**. Consequently, when a need for benzocyclobutenedione arose, we decided to develop the reaction as a flow photochemical process in order to facilitate its optimisation with respect to wavelength, residence time, solvent and acetal protecting group.^{2,3} In the event, we uncovered conditions leading to a modest yield improvement for benzocyclobutene **3** and discovered a wavelength dependence for the reaction giving high-yielding access to 8-ring lactone **5a** and orthoanhydride **6**. Herein we describe our development of those methods, together with demonstrations of their usefulness in heterocyclic and natural products synthesis, and a detailed mechanistic study by transient absorption (TA) spectroscopy and TD-DFT analysis.

Our investigation began with an examination of the wavelength dependence on the photoextrusion of carbon monoxide

from bis-acetal **2**. While all wavelengths studied led to the evolution of gas, which was vented from the system using a circulating flow technique (see ESI†), only UVA irradiation gave benzocyclobutene **3** as a significant product.^{3,4} An optimum yield of 57% was realised when a 0.05 M solution of **2** in acetonitrile was irradiated with 30 × 1.5 W UVA LEDs for residence time of 3 h. By contrast, irradiation of bis-acetal **2** with UVB or UVC light gave rise to a complex mixture of products on work-up that varied from experiment to experiment and contained no benzocyclobutene **3**, *vide infra*. Curiously, when the same reaction was performed on the homologous ninhydrin bis-acetal **4a**, 8-ring lactone **5a** was given in high yield (Scheme 1). Various analogues of **5** carrying substituents on the acetals and arene also gave the reaction in high yield (Table 1). In general, bis-1,3-dioxanes with a symmetrical arene and/or acetal substitution pattern gave a single product in 80–90% yield, *e.g.* **5a–c**. Substrates lacking those symmetry elements also gave high yields but led to isomeric mixtures, *e.g.* lactones **5d–g**. Thought this limited the yield attained for individual products to ~40%, in the context of medium ring synthesis such an outcome remains competitive.⁵



Scheme 1 The discovery of wavelength and acetal dependence on the outcome of photoextrusion reactions involving ninhydrin bis-acetals.

^a Chemistry, University of Southampton, Highfield, Southampton, SO17 1BJ, UK.
 E-mail: dch2@soton.ac.uk


^b School of Chemistry, University of Nottingham, Nottingham NG7 2RD, UK

^c Department of Chemical and Environmental Engineering,
 University of Nottingham Ningbo China, 199 Taikang East Road, Ningbo 315100,
 China


† Electronic supplementary information (ESI) available: Experimental accounts with spectral details and copies of NMR spectra are available as supplementary information. CCDC 1908866. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1cc06800h



Table 1 Extensions to other ninhydrin bis-acetals



To demonstrate the utility of the method, we next applied it in a total synthesis of fimbrialylactone B **11**, a natural product from the roots of *Strophioblastia fimbrialyx* Boerl. (Euphorbiaceae) (Scheme 2).⁶ Thus, bis-acetal **4h** was formed from the corresponding 1-indanone **7** in 76% yield through sequential SeO_2 oxidation and acetalisation with 2,2-dimethylpropandiol.⁷ Exposure of an acetonitrile solution of **4h** to UVC light under circulating flow next gave a 3:2 mixture of lactones **5h** and **5h'** that was readily separated by column chromatography. Methanolysis of the major isomer **5h** then gave ester **9** which was advanced to fimbrialylactone B **11** by ruthenium tetroxide oxidation to carboxylic acid **10** and acetal hydrolysis with HCl.⁸ Importantly, the spectral and physical data recorded on our sample matched those reported for the natural product. Its identity was also confirmed by X-ray crystallographic analysis (Scheme 2, CCDC 1908866).^{†,9}





Scheme 2 Total synthesis and X-ray crystal structure of fimbrialylactone B.

The potential of these 8-ring lactones to serve as intermediates in heterocyclic synthesis was next demonstrated with the conversion of **5a** to spirolactone **12** on acetal hydrolysis (Scheme 3).⁹ In turn, its treatment with methylamine gave isoshihunine **13** in 77% yield, from which the alkaloids shihunidin **14** and shihunine **15** can each be derived.^{10,11} Furthermore, reaction with hydrazines gave high yielding access to phthalazones (e.g. **16**, **17**, **21**), which are important pharmacophores present in the chemotherapeutic agent olaparib (Lynparza[®]),¹² and azelastine (Astellin, Optivar, Allergodil), a frontline drug treatment for mild rhinitis.¹³ By contrast, *o*-phenylenediamine, ethanolamine and ethylenediamine gave lactols **18**–**20** respectively.

The mechanistic course of the photoextrusion reaction was next studied by transient absorption (TA) spectroscopy following excitation of bis-acetal **4a** in CH_3CN at 266 nm (Scheme 4 and Fig. S1, ESI[†]). Analysis of the ps-TA spectra (Fig. S1a, ESI[†]) allowed us to extract the time constants associated with spectral evolution using global kinetic analysis.¹⁴ The data could be fitted using two exponential decays with time constants $\tau_1 = 1.8 \pm 0.4$ ps and $\tau_2 = 1.8 \pm 0.2$ ns (Fig. S1b and c, ESI[†]). The observed transient spectrum at 1 ps time delay showed an absorption band at 585 nm which was assigned to the $\text{n}\pi^*$ (S_1) excited state ${}^1[4a]$ following comparison of the extracted spectrum with that computed for the S_1 state (Fig. S7, ESI[†]). Decay of the S_1 state was associated with the formation of a new transient species with an absorption band at 454 nm. This was attributed to biradical ${}^1[22]$ based on comparison with results reported for the photoexcitation of 1,1,3-triphenyl-3-hydroxy-2-indanone.¹⁵ Additionally, the result aligns with a recent TA study of the C–C bond cleavage of cyclopentanone in cyclohexane following 255 nm photoexcitation, which Kao *et al.* found occurred in ≤ 1 ps.¹⁶ Likewise the τ_2 component [1.8 ± 0.2 ns] for the rate of formation of ${}^3[22]$ from ${}^1[22]$ is consistent with previously reported ISC timescales for acyl-alkyl diradicals.¹⁷

Spectral evolution by ns-TA spectroscopy was also monitored (Fig. S1d, ESI[†]). Three exponentials are required to describe these data, with time constants $\tau_1 = 4.8 \pm 0.4$ ns, $\tau_2 = 100 \pm 20$ ns, and

Scheme 3 Heterocycles derived from 8-ring lactone **5a**.




Scheme 4 Proposed mechanistic course of the photoextrusion reactions of bis-acetal **4a** [UVA (broad), 355–390 nm; UVB (narrow), 310 nm; UVC (narrow), 254 nm]

τ_3 = long-lived (Fig. S1e and f, ESI[†]).¹⁷ The τ_1 component was associated with a decay of the 460 nm band for $^3[22]$ with simultaneous formation of a new transient at 550 nm, attributed to the decarbonylation product $^3[23]$. A similar transient spectral feature was reported for the decarbonylation of diphenylmethylacyl radical ($\text{Ph}_2\text{MeCCO}^\bullet$) in isooctane, which had a rate constant $k = 1.5 \times 10^8 \text{ s}^{-1}$.^{15,18} Decay of the 550 nm transient species $^3[23]$, with $\tau_2 = 100 \pm 20 \text{ ns}$, was associated with the formation of a long-lived transient at 440 nm assigned to quinodimethane **26**. This is in line with the rate constant measured by Fujiwara *et al.* for the formation of *o*-xylene from 2-indanone following photoexcitation at 266 nm in cyclohexane ($k = 1.4 \pm 0.1 \times 10^7 \text{ s}^{-1}$).¹⁹ Thus, the wavelength dependence of the reaction centres on the fate of long-lived intermediate **26** (Scheme 4). Irradiation of **26** with UVA leads to a singlet excited state $^1[26]$ that can relax to benzocyclobutene **24**. By contrast, its irradiation with UVB or UVC light gives rise to higher singlet excited states of **26** from which ISC to $^3[23]$ is allowed. Collapse of $^3[23]$ to biradical **25** then provides access to 8-ring lactone **5a**.

Applying this mechanistic understanding to ninhydrin bis-acetal **2** (Scheme 5) led us to conclude that the course of that reaction was also dictated by the nature of the excitation of quinodimethane intermediate **28**. As before, its irradiation with UVA gives a singlet excited state $^1[28]$ from which benzocyclobutene **3** is derived. Similarly, irradiation with UVB or UVC light leads to biradical $^3[29]$, which in turn collapses to biradical **30**. However, in this case, ring closure to ϵ -lactone **27** is outpaced by the extrusion of ethene to biradical **31**, a precursor of orthoanhydride **6**. Indeed, the near quantitative formation of orthoanhydride **6** on photolysis of bis-acetal **2** with UVB or UVC light was confirmed by analysis of a concentrated product mixture before aqueous work-up (see ESI[†] for characterisation data).

A TD-DFT analysis of quinodimethane **28** provided additional support for the mechanism (Fig. 1).²⁰ In particular, it indicated that the energy provided by UVA irradiation (3.18 eV $\equiv \lambda_{\text{max}} 390 \text{ nm}$) mirrored that required for the $\pi \rightarrow \pi^*$ transition $S_0 \rightarrow S_1$, and that ISC between S_1 of $^1[28]$ and T_1 of $^3[29]$ was forbidden as both had $\pi\pi^*$ character.²¹ Consequently,



Scheme 5 Proposed mechanism for the photoextrusion reactions of bis-acetal **2** [UVA (broad), 355–390 nm; UVB (narrow), 310 nm; UVC (narrow), 254 nm].

irradiation of **28** with UVA induces cyclisation to benzocyclobutene **3**. By contrast, the greater energy provided by UVB and UVC irradiation facilitates ISC from $^1[28]$ to $^3[29]$ *via* higher excited states where both the orbital symmetry and energy requirements can each be fulfilled.²¹ As a result, irradiation of **28** with UVB or UVC gives orthoanhydride **6**.

Finally, the use of orthoanhydride **6** as a phthalic anhydride surrogate (Scheme 6) was demonstrated with the phthalimide protection of various 1° -amines under mild conditions leading to **30a–f**.²² Similarly, reactions with 2° -amines gave the expected amides, *e.g.* **31** and **37**, while bis-nucleophiles provided high yielding access to an array of biologically relevant nitrogen heterocycles **32–36**, **38** and **39**.

In conclusion, we have shown that the photodecarbonylation of ninhydrin bis-acetals is a wavelength dependent reaction that can be conducted on a gram scale to give benzocyclobutenes on UVA irradiation (60 W, $\sim 5 \text{ h}$), and orthoanhydrides or 8-ring lactones on UVB or UVC irradiation (36 W, $\sim 2 \text{ h}$). Notably, they formally constitute ring contraction, ring correlative, and ring expansion reactions, respectively. These versatile intermediates give access to a host of heterocycles, natural products, and pharmacophores. Mechanistic details have been delineated using computational and spectroscopic methods, including picosecond and nanosecond





Fig. 1 TD-DFT calculated energies for the singlet and triplet excited states of **28** and **29** respectively, and the mechanistic implications.





Scheme 6 Illustrations of the synthetic utility of orthoanhydride 1.

transient absorption spectroscopy. We are currently exploring further applications of the chemistry in target-oriented and natural products total synthesis.

Wei Sun and William Rimbach performed the synthetic chemistry under the supervision of David Harrowven. Wei Sun and Magnus Hanson-Heine performed the DFT analyses while Mark Light conducted the X-ray analysis. Transient absorption spectroscopic analyses were by Surajit Kayal and Xue-Zhong Sun under the supervision of Michael George.

We gratefully acknowledge financial support from EPSRC [EP/P013341/1, EP/L003325/1 and EP/K039466/1], the European Regional Development Fund [ERDF Interreg Va programme (Project 121)], and the University of Nottingham Green Chemicals Beacon.

Conflicts of interest

There are no conflicts of interest to declare.

Notes and references

- 1 D. Leinweber, M. Schnebel, R. Wartchow, H. G. Wey and H. Butenschoen, *Eur. J. Org. Chem.*, 2002, 2385; D. Leinweber and H. Butenschoen, *Tetrahedron Lett.*, 1997, **38**, 6385; D. Ng, Z. Yang and M. A. Garcia-Garibay, *Tetrahedron Lett.*, 2002, **43**, 7063.
- 2 J. P. Knowles, L. D. Elliott and K. I. Booker-Milburn, *Beilstein J. Org. Chem.*, 2012, **8**, 2025; K. Gilmore and P. H. Seeberger, *Chem. Rec.*, 2014, **14**, 410; C. Sambiagio and T. Noël, *Trends Chem.*, 2020, **2**, 92; M. B. Plutschack, B. Pieber, K. Gilmore and P. H. Seeberger, *Chem. Rev.*, 2017, **117**, 11796; D. Cambié, C. Bottecchia, N. J. W. Straathof, V. Hessel and T. Noël, *Chem. Rev.*, 2016, **116**, 10276; F. Politano and G. Oksdath-Mansilla, *Org. Process Res. Dev.*, 2018, **22**, 1045; T. H. Rehm, *ChemPhotoChem*, 2019, **3**, 1;
- 3 N. Hoffmann, *Chem. Rev.*, 2008, **108**, 1052; M. Oelgemoeller, *Chem. Eng. Technol.*, 2012, **35**, 1144.
- 4 D. C. Harrowven, M. Mohamed, T. P. Gonçalves, R. J. Whitby, D. Bolien and H. F. Sneddon, *Angew. Chem., Int. Ed.*, 2012, **51**, 4405; T. P. Gonçalves, M. Mohamed, R. J. Whitby, H. F. Sneddon and D. C. Harrowven, *Angew. Chem., Int. Ed.*, 2015, **54**, 4531; D. E. Collin, E. H. Jackman, N. Jouandon, W. Sun, D. C. Harrowven and B. Linclau, *Synthesis*, 2021, 1307; M. A. Manning, W. Sun, M. E. Light and D. C. Harrowven, *Chem. Commun.*, 2021, **57**, 4556; W. Sun, W. A. T. Rimbach, L. D. Elliott, K. I. Booker-Milburn and D. C. Harrowven, *Chem. Commun.*, 2022, **58**, 383.
- 5 For our previous work on medium-ring synthesis see: W. Sun, D. C. Wilson, M. E. Light and D. C. Harrowven, *Org. Lett.*, 2018, **20**, 4346; D. C. Harrowven, N. L'Helias, J. D. Moseley, N. J. Blumire and S. R. Flanagan, *Chem. Commun.*, 2003, 2658.
- 6 P. Seephonkai, S. G. Pyne, A. C. Willis and W. Lie, *J. Nat. Prod.*, 2013, **76**, 1358.
- 7 F. Johnson and N. Oberleender, *J. Med. Chem.*, 1991, **34**, 1662; L. F. Fieser and W. C. Lothrop, *J. Am. Chem. Soc.*, 1936, **58**, 2050.
- 8 H. Niwa, M. Nisiwaki, I. Tsukada, T. Ishigaki, S. Ito, K. Wakamatsu, T. Mori, M. Ishigaki and K. Yamada, *J. Am. Chem. Soc.*, 1990, **112**, 9001; B. S. Lucas, L. M. Luther and S. D. Burke, *Org. Lett.*, 2004, **6**, 2965.
- 9 Some discrepancies were noted in the ¹³C NMR data. These data were reported tentatively in the isolation paper as they had been attained from a spectrum recorded on a 9:1 mixture of fimbrialcalyxketones A and B.
- 10 P. M. Hare, C. T. Middleton, K. I. Mertel, J. M. Herbert and B. Kohler, *Chem. Phys.*, 2008, **347**, 383; H. Sun and H. Frei, *J. Phys. Chem. B*, 1997, **101**, 205.
- 11 V. G. Ore, M. D. Hordia and N. S. Arasimhan, *Tetrahedron*, 1990, **46**, 2483; G. A. Flynn, *Chem. Commun.*, 1980, 862; E. Breuer and S. Zbaida, *Tetrahedron*, 1975, **31**, 499.
- 12 X.-W. Li, M. Huang, K. Lo, W. L. Chen, Y.-Y. He, Y. Xu, H. Zheng, H. Hu and J. Wang, *Molecules*, 2019, **24**, 2673; H. Chen, X. Li, Y. Xu, K. Lo, H. Zheng, H. Hu, J. Wang and Y. Lin, *Molecules*, 2018, **23**, 1185; J. Bastida, M. Selles, C. Codina, F. Viladomat and J. L. Leon de la Luz, *Planta Med.*, 1996, **62**, 575; G. B. Bodem and E. Leete, *J. Org. Chem.*, 1979, **44**, 4696; E. Leete and G. B. Bodem, *J. Am. Chem. Soc.*, 1976, **98**, 6321; E. Breuer and S. Zbaida, *Tetrahedron*, 1975, **31**, 499; Y. Inubushi, Y. Tsuda, T. Konita and S. Matsumoto, *Chem. Pharm. Bull.*, 1964, **12**, 749.
- 13 P. S. Fong, D. S. Boss, T. A. Yap, A. Tutt, P. Wu, M. Mergui-Roelvink, P. Mortimer, H. Swaisland, A. Lau, M. J. O'Connor, A. Ashworth, J. Carmichael, S. B. Kaye, J. H. Schellens and J. S. de Bono, *N. Engl. J. Med.*, 2009, **361**, 123.
- 14 The results were analysed with Glotaran (<http://glotaran.org>).
- 15 G. Kuzmanich, J. Xue, J. C. Netto-Ferreira, J. C. Scaiano, M. Platz and M. A. Garcia-Garibay, *Chem. Sci.*, 2011, **2**, 1497.
- 16 M. H. Kao, R. K. Venkatraman, M. N. Ashfold and A. J. Orr-Ewing, *Chem. Sci.*, 2020, **11**, 1991.
- 17 Y. P. Tsentalovich, O. B. Morozova, N. I. Avdievich, G. S. Ananchenko, A. V. Yurkovskaya, J. D. Ball and M. D. Forbes, *J. Phys. Chem. A*, 1997, **101**, 8809.
- 18 C. Chatgilialoglu, D. Crich, M. Komatsu and I. Ryu, *Chem. Rev.*, 1999, **99**, 1991.
- 19 M. Fujiwara, K. Mishima, K. Tamai, Y. Tanimoto, K. Mizuno and Y. Ishii, *J. Phys. Chem. A*, 1997, **101**, 4912.
- 20 A. Dreuw and M. Head-Gordon, *Chem. Rev.*, 2005, **105**, 4009.
- 21 M. A. El-Sayed, *J. Chem. Phys.*, 1963, **38**, 2834.
- 22 V. V. Sureshbabu, N. Narendra, in *Amino Acids, Peptides and Proteins in Organic Chemistry: Protection Reactions, Medicinal Chemistry, Combinatorial Synthesis*, ed.: A. B. Hughes, Wiley-VCH, Weinheim, 2011, vol. 4, pp. 1–97.

