

### Showcasing research from Professor Christian Müller's laboratory, Institute of Chemistry and Biochemistry, Freie Universität Berlin, Berlin, Germany

A new access to diazaphospholes *via* cycloaddition-cycloreversion reactions on triazaphospholes

The image highlights the [4+2] cycloaddition reaction between an unsaturated 5-membered phosphorus heterocycle and hexafluoro-2-butyne, affording a  $CF_3$ -substituted diazaphosphole after cycloreversion and elimination of 'Bu-C $\equiv$ N.



See László Nyulászi, Christian Müller *et al.*, *Chem. Commun.*, 2022, **58**, 7745.



rsc.li/chemcomm Registered charity number: 207890

# ChemComm



**View Article Online** 

# This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence. Open Access Article. Published on 15 June 2022. Downloaded on 10/5/2024 9:01:00 PM.

# COMMUNICATION

Check for updates

Cite this: Chem. Commun., 2022, 58, 7745

Received 21st April 2022, Accepted 15th June 2022

DOI: 10.1039/d2cc02269a

rsc.li/chemcomm

# A new access to diazaphospholes *via* cycloaddition-cycloreversion reactions on triazaphospholes<sup>†</sup>

Lea Dettling, ២ ‡ª Martin Papke, ‡ª Julian A. W. Sklorz, ª Dániel Buzsáki, ២ b Zsolt Kelemen, b Manuela Weber, ª László Nyulászi 🕩 \*b and Christian Müller 🕩 \*ª

A novel bis-CF<sub>3</sub>-substituted diazaphosphole was synthesized selectively from hexafluoro-2-butyne and a 3H-1,2,3,4-triazaphosphole derivative. The [4+2] cycloaddition and subsequent cycloreversion reaction under elimination of pivaloyl nitrile affords the product in high yield. The heterocycle coordinates *via* the phosphorus atom to a W(CO)<sub>5</sub>-fragment and shows stronger  $\pi$ -accepting properties than the triazaphosphole.

3,5-Disubstituted 3*H*-1,2,3,4-triazaphospholes (**B**) are the phosphorus congeners of the well-studied 1,2,3-triazoles (**A**), according to the isolobal relationship between a trivalent P-atom and a C-H fragment (Chart 1).

These  $\lambda^3$ ,  $\sigma^2$  phosphorus heterocycles have a conjugated  $\pi$ -system with a high degree of aromaticity.<sup>1</sup> They can easily be prepared regioselectively by a modular [3 + 2] cycloaddition reaction, starting from various aryl/alkyl-azides and phosphaalkynes.<sup>2,3</sup> Despite the fact that *3H*-1,2,3,4-triazaphosphole derivatives have been synthesized independently by Carreé and Regitz already in 1984, the first reports on their coordination chemistry have not appeared in literature before 2010.<sup>2,4</sup> As ambidentate ligands the coordination of the heterocycle to a metal center might proceed either *via* the phosphorus atom or the nitrogen donors N<sup>1</sup> or N<sup>2</sup> (Chart 1, C). However, the  $\eta^1(P)$ -coordination mode has so far only been observed in a Pt(0)-complex.<sup>4b</sup>

Even less is known about the chemical reactivity of 3H-1,2,3,4-triazaphosphole derivatives. We could demonstrate that the cationic phosphorus analogues **D** of neutral mesoionic carbenes (1,2,3-triazolylidenes) can be obtained by quaternization of the N1 atom in **B** with Meerwein salts.<sup>5</sup> Moreover, we noticed that the introduction of electron-withdrawing *N*-sulfonyl groups at the N<sup>3</sup>-atom changes the reactivity of the corresponding triazaphosphole considerably. In the presence of stoichiometric amounts of AuCl-S(CH<sub>3</sub>)<sub>2</sub>, loss of N<sub>2</sub> and the formation of *cyclo*-1,3-diphospha(m)-2,4-diazane-Au(I) complexes of type **E** were observed.<sup>6</sup> Inspired by the fact that 6-membered azaphosphinines and 5-membered azaphospholes can undergo [4 + 2] cycloaddition reactions with various alkynes under subsequent nitrile elimination, we decided to investigate the reactivity of **B** towards alkynes in more detail with the aim to synthesize 2*H*-1,2,3-diazaphosphole derivatives (**G**) directly in one step (Chart 2).<sup>7</sup>

These heterocycles are otherwise only accessible by multistep synthetic procedures.<sup>8</sup> In fact, similar reactions with  $RC \equiv P$  elimination from oxadiphospholes and selenadiphospholes *via* a concerted mechanism have been reported.<sup>9</sup> Moreover, an imino-substituted diazaphosphole biradicaloid showed facile isonitrile cycloaddition, but no subsequent cycloreversion.<sup>10</sup>

The 3,5-disubstituted triazaphosphole **1** was prepared according to literature procedures from  $PhN_3$  and <sup>*t*</sup>Bu-C $\equiv P$ .<sup>2*a*</sup> Triazaphosphole **3** does not react with dimethyl



Chart 1 Triazole A, triazaphosphole B and possible coordination modes C. Selected examples (D and E) for the reactivity of B.

<sup>&</sup>lt;sup>a</sup> Freie Universität Berlin, Institute of Chemistry and Biochemistry, Fabeckstr. 34/36, 14195, Berlin, Germany. E-mail: c.mueller@fu-berlin.de

<sup>&</sup>lt;sup>b</sup> Department of Inorganic and Analytical Chemistry, Budapest University of Technology and MTA-BME Computation Driven Chemistry Research Group, Műegyetem rkp. 3, H-1111, Budapest, Hungary.

E-mail: nyulaszi.laszlo@vbk.bme.hu

<sup>†</sup> Electronic supplementary information (ESI) available. CCDC 2163856–2163858. For ESI and crystallographic data in CIF or other electronic format see DOI: https://doi.org/10.1039/d2cc02269a

<sup>‡</sup> These authors have contributed equally.



acetylenedicarboxylate (DMAD) to diazaphosphole 2 (Scheme 1a). Using the stronger dienophile hexafluoro-2-butyne, however, elimination of <sup>t</sup>Bu–C≡N and, according to <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy, quantitative formation of diazaphosphole **3** was observed (Scheme 1b). We could not detect the apparent intermediate **F** (Chart 2) during the course of the reaction. Interestingly, triazoles, such as **4**, did not react with CF<sub>3</sub>C≡CCF<sub>3</sub> to the CF<sub>3</sub>-substituted pyrazole **5**, although cycloaddition/cycloreversion reactions on 1,2,3-triazoles with DMAD have been reported in the literature (Scheme 1c).<sup>11</sup> This is particularly intriguing as 1-aryl-3,4-bis(trifluoromethyl)-substituted pyrazole motifs (5), are present in numerous pharmacologically relevant and bioactive nitrogen heterocycles and have to be prepared *via* a multistep synthesis.<sup>12</sup> Our novel diazaphosphole **3** thus represents a phosphorus derivative of this compound class.

Diazaphosphole **3** was obtained as an off-white solid in 87% isolated yield and shows a signal at  $\delta(\text{ppm}) = 234.4$  (q,  ${}^{3}J_{\text{P-F}} = 25.5$  Hz) in the  ${}^{31}\text{P}{}^{1}\text{H}$  NMR spectrum (starting material **1**:  $\delta(\text{ppm}) = 174.3$ ). For the CF<sub>3</sub>-groups, resonances at  $\delta(\text{ppm}) = -53.3$  (dq,  ${}^{3}J_{\text{F-P}} = 25.5$ ,  ${}^{5}J_{\text{F-F}} = 7.4$  Hz) and  $\delta(\text{ppm}) = -61.8$  (qd,  ${}^{5}J_{\text{F-F}} = 7.4$  Hz,  ${}^{4}J_{\text{F-P}} = 1.2$  Hz) were observed in the  ${}^{19}\text{F}{}^{1}\text{H}$  NMR spectrum. Single crystals of **3** suitable for X-ray diffraction were obtained by slow evaporation of a dichloromethane solution and the molecular structure of **3** in the crystal is depicted in Fig. 1 along with selected bond lengths and distances.

Fig. 1 represents the first crystallographically characterized CF<sub>3</sub>-substituted diazaphosphole. From the X-ray data it is evident that the heterocycle is fully planar and that the P(1)–C(8) and N(1)–N(2) bond distances in 3 are very similar to the ones observed in the starting material 1,<sup>13</sup> with P–C and C–C bond lengths characteristic for aromatic compounds. The significantly negative NICS(1) values (see Table S1 in the ESI<sup>†</sup>)



Scheme 1 Reactivity of 1 and 4 towards electron-withdrawing alkynes.



**Fig. 1** Molecular structure of **3** in the crystal. Displacement ellipsoids are shown at the 50% probability level. Selected bond lengths (Å) and angles (°): P(1)-N(2): 1.693(2), P(1)-C(8): 1.712(2), C(8)-C(7): 1.406(3), C(7)-N(1): 1.323(3), N(1)-N(2): 1.349(2), N(2)-C(1): 1.436(3). N(1)-N(2)-C(1)-C(2): 145.9(2).

are in accordance with aromaticity. Apparently, exchanging pivaloyl nitrile by a perfluorobutyne-moiety does not cause a significant structural change within the heterocycle. The same holds for the inter-ring N(2)-C(1) distance. Also, the N(2)-P(1)-C(8) and P(1)-N(2)-N(1) angles as well as the torsion angle N(1)-N(2)-C(1)-C(2) in 3 are very similar compared to the data found for triazaphosphole **1**.

In order to understand the reaction mechanism,  $\omega$ B97X-D/6-311 + G\*\* DFT calculations (see ESI<sup>†</sup>) were performed after validating the optimized geometries with the X-ray data of **3** (see Table S2, ESI<sup>†</sup>). This level of theory was used successfully for cycloaddtion reactions before.<sup>14</sup>

The concerted cycloaddition-cycloreversion process (Fig. 2, Chart 2 and ESI<sup>†</sup>) is in full agreement with all experimental observations. The cycloaddition step  $\mathbf{B} \rightarrow \mathbf{F}$  (Chart 2) is nearly thermoneutral, while the 'BuC  $\equiv$  N eliminating cycloreversion (forming G) is highly exergonic. Accordingly (see Hammond principle), the rate determining step of the overall reaction is **TS1**, that allows the formation of 3 (27.3 kcal mol<sup>-1</sup> activation Gibbs free energy) but not of 2 and 5 (barriers 31.9 kcal mol<sup>-1</sup>,



**Fig. 2**  $\omega$ B97X-D/6-311 + G\*\*(PCM = toluene) Gibbs free energy (*T* = 130 °C) profiles for the reactions in Scheme 1. Relative energies (in kcal mol<sup>-1</sup>) are compared to the initial van der Waals complex of the reactants.

36.9 kcal mol<sup>-1</sup>, respectively). It is noteworthy that IRC calculations reveal, that the Ph substituent at the nitrogen atom should be in endo position with respect to the approaching/leaving group for any [4+2] cycloaddition step. The interconversion  $\mathbf{F} \rightarrow \mathbf{F}'$  is needed prior to the retro cycloaddition step, by flattening the pyramidal nitrogen atom *via* a small barrier. The fact that no intermediate  $\mathbf{F}/\mathbf{F}'$  could be detected is in accordance with the very small barrier for the cycloreversion step.

A comparison of the Kohn-Sham orbitals of the parent CF<sub>3</sub>substituted diazaphosphole, the parent 2H-1,2,3-diazaphosphole and the parent 3H-1,2,3,4-triazaphosphole (Fig. S1, ESI<sup>+</sup>) shows, that in all three heterocycles, the  $\pi$ -type LUMO has a large coefficient at the phosphorus atom, indicating good  $\pi$ -acceptor properties when coordinated via the phosphorus atom to a metal center. While the orbital energies of the unsubstituted diazaphosphole are generally destabilized with respect to the triazaphosphole (Fig. S1, ESI<sup>+</sup>), in accordance with the observed ionization energies,<sup>1</sup> CF<sub>3</sub>-substitution acts strongly stabilizing (Fig. S1, ESI<sup>†</sup>). Altogether, 3 should be a stronger  $\pi$ -acceptor than 1. In all three compounds, the lone pair at the phosphorus atom (mixed with the nitrogen in-plane lone-pair) is represented by the HOMO-2(CF<sub>3</sub>-diazaphosphole: E = -11.24 eV; 3H-1,2,3,4-triazaphosphole: E = -10.92 eV; 2H-1,2,3-diazaphosphole: E = -10.21 eV). Consequently, triazaphospholes and diazaphospholes are expected to be rather weak  $\sigma$ -donors, as anticipated for low-coordinate phosphorus compounds. The  $\pi$ -donor properties of triazaphospholes and diazaphospholes are evident from the HOMOs, each having a large  $\pi$ -coefficient at the phosphorus atom, as it is known for other electron-rich phosphorus heterocycles.<sup>15</sup> Again, due to the energetically higher HOMO, triazaphosphole 1 should show stronger  $\pi$ -donor properties than the CF<sub>3</sub>-substituted diazaphosphole 3.

The interplay between the above described effects makes the coordination behavior of compound 3 highly interesting, also with respect to triazaphospholes. As a matter of fact, the coordination chemistry of 2H-1,2,3-diazaphospholes is largely unknown and only a few examples can be found in the literature. Chart 3 shows the possible coordination modes for this class of compounds. Analogous to triazaphospholes, diazaphospholes are ambidentate ligands and can coordinate to a metal center either *via* the phosphorus lone pair (H) or the nitrogen donor (I). This has been demonstrated in a few cases by van Koten, Schmidpeter and co-workers by using suitable  $Pt(\pi)$  and  $Pd(\pi)$  complexes as metal precursor.<sup>16</sup> The simultaneous coordination of a diazaphosphole to two metal fragments (J) has so far not been observed. Only recently, Erben and co-workers have investigated the synthesis and coordination chemistry of Si-bridged, chelating diazaphospholes.<sup>17</sup>

We decided to focus on the synthesis of a tungsten carbonyl complex of 3, as it can provide valuable information on the electronic ligand properties via IR spectroscopy. Moreover, <sup>31</sup>P NMR spectroscopy would immediately reveal, whether the coordination of the ligand to the W(CO)<sub>5</sub> fragment occurs via the phosphorus or the nitrogen donor. 3 was reacted with one equivalent of W(CO)<sub>6</sub> in THF at room temperature and under UV irradiation (Scheme 2). After only a short time, the formation of a single new resonance at  $\delta(ppm) = 217.3$  was observed in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum, which corresponds to a coordination shift of  $\Delta\delta(\text{ppm}) = 17.1$  compared to the starting material.<sup>18</sup> The selective reaction towards product 6 was complete within 68h. Interestingly, the signal of the product at  $\delta(\text{ppm}) = 217.3$  shows tungsten satellites with a coupling constant of  ${}^{1}J_{W-P}$  = 326.5 Hz (Fig. S9, ESI<sup>†</sup>). This indicates that coordination of the ligand to the metal center occurs via the phosphorus atom, in agreement with the calculated 10.6 kcal  $mol^{-1}$  preference of the coordination at phosphorus over nitrogen. For comparison reasons, we also reacted triazaphospholes 1 and 8 (Ar = 2,5-diisopropylphenyl, Dipp) with  $W(CO)_6$  in THF at room temperature and under irradiation with UV light. The course of the reaction was again followed by means of NMR spectroscopy, which revealed a selective and quantitative formation of a new species within 5d. The new compounds (7, 9) show a signal at  $\delta$ (ppm) = 136.1, respectively  $\delta(\text{ppm}) = 160.6 \text{ in the } {}^{31}\text{P}{}^{1}\text{H} \text{NMR spectrum } (\Delta\delta(\text{ppm}) = 38.2,$ 39.7). Much to our surprise, these signals also show tungsten satellites ( ${}^{1}J_{P-W}$  = 262.1 Hz; 285.6 Hz), which verifies that also 1 and 8 coordinate via the phosphorus atom to the metal center. This is particularly interesting taking into account that a coordination via N<sup>1</sup> or N<sup>2</sup> (Chart 1) has so far been observed for the majority of triazaphosphole-based complexes.<sup>4,19</sup> The calculated 0.6 kcal mol<sup>-1</sup> energy difference between the two complexation modes of 1 indicates that subtle steric effects determine the complexation site in triazaphospholes.

A comparison of the IR spectra of **6**, **7** and **9** further shows, that the wavenumbers of the CO stretching frequencies are shifted to higher values in **6** compared to the ones found for **7** and **9** (Table 1). This is in line with the expected lower netdonor properties of **3** compared to **1** and **8**.

Thus, the CF<sub>3</sub>-substituted diazaphosphole **3** is a stronger  $\pi$ -accepting ligand than triazaphospholes **1** and **8**, if coordination to the metal center proceeds *via* the phosphorus donor.





Scheme 2 Synthesis of W(0)-complexes 6 and 7.

[M]

Table 1 Experimental wavenumbers [in  $\rm cm^{-1}]$  for the CO stretching modes. These data were also supported by DFT calculations (see Table S3 in the ESI)



Fig. 3 Molecular structures of **6** (left) and **9** (right) in the crystal. Displacement ellipsoids are shown at the 50% probability level. Selected bond lengths (Å) and angles (°): **6**: P(1)–N(2): 1.677(2), P(1)–C(8): 1.707(2), C(8)–C(7): 1.413(3), C(7)–N(1), N(1)–N(2): 1.361(3), P(1)–W(1): 2.3890(6), N(2)–C(1): 1.441(3). N(1)–N(2)–C(1)–C(2): 64.1(3). **7**: Only one independent molecule is shown. P(1)–N(2): 1.675(2), P(1)–C(1): 1.712(2), C(1)–N(3): 1.357(3), N(3)–N(2): 1.303(2), N(2)–N(1): 1.358(2), N(1)–C(2): 1.446(2), C(1)–C(14): 1.521(3), P(1)–W(1): 2.4512(5). N(1)–P(1)–C(1): 88.67(9), N(2)–N(1)–C(2)–C(3): 86.3(2).

Finally, single crystals of **6** and **9**, suitable for X-ray diffraction, could be obtained by slow evaporation of the solvent of a saturated solution of **6** and **9** in *n*-pentane. Fig. 3 shows the molecular structures of **6** and **9** in the crystal, along with selected bond lengths and angles. The W(0) complexes **6** and **9** show a slightly distorted octahedral coordination geometry and unequivocally confirm that the heterocycles coordinates *via* the phosphorus atom to the W(CO)<sub>5</sub> fragment. Compared to the solid state structure of the free ligand **3** (Fig. 1), the P(1)–C(8) and P(1)–N(2) bonds in **6** are slightly shortened upon coordination of the ligand to the metal center (1.707(2) Å and 1.677(2) Å in **6** *vs*. 1.712(2) Å and 1.693(2) Å in **3**). For steric reasons, the aryl rings in **6** and **9** are rotated out of the heterocyclic plane (see also Fig. 1).

We could demonstrate for the first time that a 3H-1,2,3,4triazaphosphole derivative undergoes a selective [4 + 2] cycloaddition with hexafluoro-2-butyne with subsequent elimination of pivaloyl nitrile to afford a bis-CF<sub>3</sub>-substituted diazaphosphole in high yield. According to the isolobal relationship between a trivalent phosphorus atom and a C–H fragment, this heterocycle represents a phosphorus congener of a bis-CF<sub>3</sub>substituted pyrazole, which finds applications as a bioactive nitrogen heterocycle. The novel diazaphosphole forms an (L)W(CO)<sub>5</sub>-complex, in which the ligand coordinates *via* the phosphorus atom to the metal center. In combination with DFT-calculations, the experimental results show that the bis-CF<sub>3</sub>-substituted diazaphosphole is a stronger  $\pi$ -acceptor than the corresponding triazaphosphole, which was used as a starting material. Our results demonstrate that bis-CF<sub>3</sub>-substituted diazaphospholes are accessible in a facile manner. Their use as novel  $\pi$ -accepting ligands in coordination chemistry and homogeneous catalysis as well as the investigation of their potential bioactive properties is currently explored.

The authors are grateful for financial support provided by Freie Universität Berlin. Z. K. is grateful for the general support of Hungarian Academy of Science under the Premium Postdoctoral Research Program 2019.

## Conflicts of interest

There are no conflicts to declare.

### Notes and references

- 1 L. Nyulászi, T. Veszprémi, J. Réffy, B. Burkhardt and M. Regitz, J. Am. Chem. Soc., 1992, 114, 9080.
- 2 (a) Y. Y. C. Yeung Lam, Ko, R. Carrié, A. Muench and G. Becker, J. Chem. Soc., Chem. Commun., 1984, 1634; (b) W. Rösch and M. Regitz, Angew. Chem., Int. Ed. Engl., 1984, 23, 900.
- 3 L. Nyulászi, Chem. Rev., 2001, 101, 1229.
- 4 (a) S. L. Choong, A. Nafady, A. Stasch, A. M. Bond and C. Jones, *Dalton Trans.*, 2013, 42, 7775; (b) S. L. Choong, C. Jones and A. Stasch, *Dalton Trans.*, 2010, 39, 5774.
- 5 M. Papke, L. Dettling, J. A. W. Sklorz, D. Szieberth, L. Nyulászi and C. Müller, *Angew. Chem., Int. Ed.*, 2017, **56**, 16484.
- 6 E. Yue, L. Dettling, J. A. W. Sklorz, S. Kaiser, M. Weber and C. Müller, *Chem. Commun.*, 2022, **58**, 310.
- 7 (a) N. Avavari, P. Le Floch and F. Mathey, J. Am. Chem. Soc., 1996, 118, 11978; (b) A. Schmidpeter and H. Klehr, Z. Naturforsch., B: Anorg. Chem., Org. Chem., 1983, 38, 1484.
- 8 (a) G. Baccolini, R. Dalpozzo and E. Errani, *Tetrahedron*, 1987,
  43, 2755; (b) J. H. Weinmaier, G. Brunnhuber and A. Schmidpeter,
  *Chem. Ber.*, 1980, 113, 2278; (c) F. Armbruster, U. Klingebiel and
  M. Noltemeyer, *Z. Naturforsch.*, 2006, 61b, 225.
- 9 (a) A. Mack, U. Bergsträßer, G. J. Reiß and M. Regitz, Eur. J. Org. Chem., 1999, 587; (b) S. Asmus, L. Nyulászi and M. Regitz, J. Chem. Soc., Perkin Trans. 2, 2001, 1968.
- 10 H. Beer, J. Bresien, D. Michalik, A.-K. Rölke, A. Schulz, A. Villinger and R. Wustrack, J. Org. Chem., 2020, 85, 14435.
- 11 Á. Díaz-Ortiz, A. de Cózar, P. Prieto, A. de la Hoz and A. Moreno, *Tetrahedron Lett.*, 2006, 47, 8761.
- (a) S. Eguchi. *Bioactive Heterocycles I*, Springer, Heidelberg, 2006;
  (b) Y. L. Yagupolskii, N. V. Pavlenko, I. I. Gerus, S. Peng and M. Nappa, *ChemistrySelect*, 2019, 4, 4604.
- 13 J. A. W. Sklorz, S. Hoof, N. Rades, N. De Rycke, L. Könczöl, D. Szieberth, M. Weber, J. Wiecko, L. Nyulászi, M. Hissler and C. Müller, *Chem. – Eur. J.*, 2015, 21, 11096.
- 14 S. Giese, D. Buzsáki, L. Nyulászi and C. Müller, Chem. Commun., 2019, 55, 13812.
- 15 C. Batich, E. Heilbronner, V. Hornung, A. J. Ashe III, D. T. Clark, U. T. Gobley, D. Kilcast and I. Scanlan, *J. Am. Chem. Soc.*, 1973, 95, 929.
- 16 (a) J. G. Kraaijkamp, D. M. Grove, G. van Koten and A. Schmidpeter, *Inorg. Chem.*, 1988, 27, 2612; (b) J. G. Kraaijkamp, G. van Koten, K. Vrieze, D. M. Grove, E. A. Klop, A. L. Spek and A. Schmidpeter, *J. Organomet. Chem.*, 1983, 256, 375.
- 17 P. Kozáček, L. Dostál, A. Růžička, I. Císařová, Z. Černošek and M. Erben, New J. Chem., 2019, 43, 13388.
- 18 See also:(a) A. B. Grimm, S. Evariste, A. L. Rheingold, C. E. Moore and J. D. Protasiewicz, *J. Organomet. Chem.*, 2017, 851, 9; (b) J. Heinicke, N. Gupta, S. Singh, A. Surana, O. Kühl, R. K. Bansal, K. Karaghiosoff and M. Vogt, *Z. Anorg. Allg. Chem.*, 2002, 628, 2869.
- 19 J. A. W. Sklorz, S. Hoof, M. G. Sommer, F. Weißer, M. Weber, J. Wiecko, B. Sarkar and C. Müller, Organometallics, 2014, 33, 511.