



Cite this: *Chem. Commun.*, 2022, **58**, 2580

DOI: 10.1039/d2cc90023h

rsc.li/chemcomm

Correction: Synthesis of glycerolipids containing simple linear acyl chains or aromatic rings and evaluation of their Mincle signaling activity

Takanori Matsumaru,^{*ab} Risa Ikeno,^a Yusuke Shuchi,^a Toshiki Iwamatsu,^a Takashi Tadokoro,^a Sho Yamasaki,^c Yukari Fujimoto,^b Atsushi Furukawa^a and Katsumi Maenaka^{*a}

Correction for 'Synthesis of glycerolipids containing simple linear acyl chains or aromatic rings and evaluation of their Mincle signaling activity' by Takanori Matsumaru *et al.*, *Chem. Commun.*, 2019, **55**, 711–714, DOI: 10.1039/C8CC07322H.

The authors regret that the structures of brartemicin and compounds **6a** and **b** presented in Fig. 2 of the article were incorrect. The correct structures are depicted below. In addition, explanations of the R' groups have been added below each compound.




Fig. 2 (A) Synthesis of **2a**–**2e**, (B) the structure of brartemicin, and (C) synthesis of **5a** and **b** and **8a** and **b**.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo, Hokkaido 060-0812, Japan. E-mail: maenaka@pharm.hokudai.ac.jp

^b Faculty of Science and Technology, Keio University, 3-14-1 Hiyoshi, Kohoku-ku, Yokohama, Kanagawa 223-8522, Japan. E-mail: matsumaru@keio.chem.ac.jp

^c Department of Molecular Immunology Research Institute for Microbial Diseases, and Laboratory for Molecular Immunology, Osaka University, 3-1 Yamadaoka, Suita, Osaka 565-0871, Japan

