



Cite this: *Chem. Commun.*, 2022, 58, 9437

DOI: 10.1039/d2cc90272a


rsc.li/chemcomm

Correction: Visible-light-mediated photoredox minisci C–H alkylation with alkyl boronic acids using molecular oxygen as an oxidant

Jiayang Dong,^a Fuyang Yue,^a Hongjian Song,^a Yuxiu Liu^a and Qingmin Wang^{*ab}

Correction for 'Visible-light-mediated photoredox minisci C–H alkylation with alkyl boronic acids using molecular oxygen as an oxidant' by Jiayang Dong *et al.*, *Chem. Commun.*, 2020, **56**, 12652–12655, <https://doi.org/10.1039/D0CC05946C>.

The authors regret that the regioselectivity of products **19** and **20** in Table 2 was incorrect. The correct structures, regioisomers of the originally proposed structures, are shown below.



The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a State Key Laboratory of Elemento-Organic Chemistry, Research Institute of Elemento-Organic Chemistry, College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China. E-mail: wangqm@nankai.edu.cn

^b Collaborative Innovation Center of Chemical Science and Engineering (Tianjin), Tianjin 300071, People's Republic of China

