Dalton Transactions

PAPER

Check for updates

Cite this: *Dalton Trans.*, 2022, **51**, 2050

Received 13th December 2021, Accepted 11th January 2022 DOI: 10.1039/d1dt04192d

rsc.li/dalton

Introduction

Bond activation reactions mediated by low-valent main group element complexes gained increasing interest within the past two decades due to the potential of establishing main group element-based catalytic transformations.¹ In this light, the reactivity of monovalent group 13 β -diketiminates LM (M = Al, Ga, In; L = HC[C(Me)N(Dip)]₂, Dip = 2,6-i-Pr₂C₆H₃) in the formal oxidation state +1 has been extensively investigated.² Typical reaction patterns include the formation of Lewis acidbase adducts,³ oxidative addition reactions of σ - and π -bonds,⁴ and two-electron reduction reactions, which can also be utilized consecutively by optimizing the reaction conditions.⁵ A key step in transition metal-catalyzed C-C cross coupling reactions is the initial oxidative addition of C-X (X = F, Cl, Br, I) bonds to the metal center.⁶ Remarkably, group 13 diyls LM have been found to be capable of C-X oxidative addition. Indanediyl LIn was shown to react with alkyl bromides and iodides to yield the corresponding bond insertion products L(X)InR (R = Me, i-Pr, t-Bu; X = Br, I), whereas it was found

^aFaculty of Chemistry, University of Duisburg-Essen, Universitätsstr. 5-7,

S07 S03 C30, D-45117 Essen, Germany. E-mail: stephan.schulz@uni-due.de;

Fax: (+49) 201-1833830https://www.uni-due.de/ak_schulz/index_en.php

of chloromethanes and chlorosilanes by gallanediyl LGa†

Geminal C–Cl and Si–Cl bond activation

Christoph Helling,^a Chelladurai Ganesamoorthy,^a Christoph Wölper^a and Stephan Schulz ¹/₂*^{a,b}

The activation of relatively inert E–X σ -bonds by low-valent main group metal complexes is receiving increasing interest. We here confirm the promising potential of gallanediyl LGa (L = HC[C(Me)N(Dip)]₂, Dip = 2,6-i-Pr₂C₆H₃) to activate E–Cl (E = C, Si) σ -bonds of group 14 element compounds. Equimolar reactions of LGa with chloromethanes and chlorosilanes EH_xCl_{4-x} (E = C, x = 0-2; E = Si, x = 0, 1) occurred with E–Cl bond insertion and formation of gallylmethanes and -silanes L(Cl)GaEH_xCl_{3-x} (E = C, x = 2 (**1**), 1 (**2**), 0 (**3**); E = Si, x = 1 (**4**). In contrast, consecutive insertion into a geminal E–Cl bond was observed with two equivalents of LGa, yielding digallyl complexes [L(Cl)Ga]₂EH_xCl_{2-x} (E = C, x = 2 (**5**); E = Si, x = 1 (**6**), 0 (**7**). Compounds **1–7** were characterized by heteronuclear NMR (¹H, ¹³C, ²⁹Si (**4**, **6**)), IR spectroscopy and elemental analysis, and their solid-state structures were determined by single-crystal X-ray diffraction (sc-XRD).

unreactive towards alkyl chlorides and aryl iodides due to the stronger C-Cl and C(sp²)-I bonds, respectively.⁷ In contrast, stronger reducing gallanediyl LGa was found to activate the C-Cl σ -bond in t-BuCl and even the C-F σ -bonds in C₆F₆ and C_6F_5H under forcing conditions, yielding L(Cl)Gat-Bu (I)⁸ and $L(F)GaAr^{F}$ ($Ar^{F} = C_{6}F_{5}$ II, $C_{6}F_{4}H$ III),⁹ respectively. The lightest and most reactive alanediyl LAl readily activated strong C(sp³)-F and C(sp²)-F bonds in alkyl, vinyl, and aryl fluorides via oxidative addition, α -fluoride elimination, and β -fluoride elimination mechanisms with formation of alanes L(F)AlR.¹⁰ Furthermore, LGa reacted with SnCl₂ in reduction and bond insertion cascades to metalloid Sn clusters, [L(Cl)Ga]₂Sn₇ and [L(Cl)Ga]₄Sn₁₇,¹¹ while intermetallic compounds [LGa]₂Ge₂ and [LGa]₂Ge₄ were obtained from reactions with (Cy₃P)GeCl₂/ KC₈ and (NHC)GeCl₂, respectively.¹² Oxidative additions of Sn-H and Pb-Cl bonds were observed with Ph₃SnH and Me₃PbCl, yielding L(H)GaSnPh3^{4a} and L(Cl)GaPbMe3,¹³ respectively, and two-fold bond insertion reactions produced [L(Cl)Ga]₂SnCl₂⁸ and plumbylene [L(F₃C(O)₂SO)Ga]₂Pb(THF).¹³

In case of Si, reactions of LGa with SiX₄ gave L(X)GaSiX₃ (X = Cl IV, Br V),^{8,14} and compound V reacted with another equivalent of LGa to $[L(Br)Ga]_2SiBr_2$ (VI) *via* consecutive Si–Br bond insertion. Most remarkably, the reaction of VI with LGa under CO atmosphere produced the stable Si–carbonyl complex $[L(Br)Ga]_2Si(CO)$ (VII), whereas in the absence of CO transient silylene $[L(Br)Ga]_2Si$ was found to activate a C–C bond of the ligand backbone to form compound VIII (Fig. 1).¹⁴ In contrast, no further reactivity with LGa was observed for IV.⁸



View Article Online

^bCenter for NanoIntegration Duisburg-Essen (CENIDE), University of Duisburg-Essen, 47057 Duisburg, Germany

[†]Electronic supplementary information (ESI) available: Spectroscopic (NMR and IR) and crystallographic data. CCDC 2127285–2127291. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1dt04192d



Fig. 1 Selected compounds derived from LGa relevant to this study.

We have reported in recent years on σ -bond activation reactions of group 13 diyls LM (M = Al, Ga, In) with main group element complexes of group 13,¹⁵ 15,¹⁶ and 16,¹⁷ respectively, resulting in the formation of a large variety of complexes with unusual bonding properties and electronic structures including double bonded species, cluster-type complexes and main group metal centered radicals. We now became interested to study the general reactivity of LGa towards chloromethanes and chlorosilanes with geminal E–Cl bonds (E = C, Si), and report herein on the synthesis and characterization of mono-and bis-insertion products.

Results and discussion

Synthesis and properties

We initially studied reactions of LGa with dichloromethane (DCM, CH_2Cl_2), chloroform (CHCl₃), and carbon tetrachloride (CCl₄). Dissolution of LGa in both DCM and chloroform at ambient temperature immediately resulted in the formation of colorless solutions containing the C–Cl bond activation products L(Cl)GaCH₂Cl (1) and L(Cl)GaCHCl₂ (2), which were isolated as colorless crystalline solids in good yields after removal of the solvent and crystallization from *n*-pentane (1) and *n*-hexane (2), respectively (Scheme 1).

In situ ¹H NMR spectroscopic studies revealed the virtually quantitative formation of **1** and **2**, and no other reaction products were detected in the reaction solutions. This observation contrasts those reported by Fischer *et al.*, who observed subsequent decomposition reactions and formation of undefined products in reactions of LGa with DCM and chloroform.⁸ In contrast, dissolution of LGa in carbon tetrachloride at ambient temperature gave a new species together with considerable amounts of LGaCl₂¹⁸ as the major compounds in solution as determined by *in situ* ¹H NMR spectroscopy. The formation of LGaCl₂ indicates the presence of a reduction pathway, which



Scheme 1 Synthesis of compounds 1-4.

was successfully suppressed by addition of carbon tetrachloride to a toluene solution of LGa at low temperature. Pure L(Cl)GaCCl₃ (3) was obtained as a colorless crystalline material in moderate yield after crystallization from DCM (Scheme 1). Under inert gas atmosphere, compounds 1 and 2 are stable in the solid-state and in solution even at elevated temperatures (80 °C), whereas 3 is stable in the solid-state but slowly decomposes in solution with formation of LGaCl₂ and minor amounts of so far unidentified products, potentially arising from reactions with transient dichlorocarbene. The selective formation of compounds 1-3 is remarkable since only a few (low-valent) main group species were found to activate DCM and chloroform via simple C-Cl bond insertion, i.e. neutral stannylenes,¹⁹ germyliumylidene,²⁰ phosphenium cations,²¹ and silylenes,²² while such reactivity was not yet observed towards carbon tetrachloride. With respect to group 13 elements, $[In(L)]OSO_2CF_3$ (L = 18-c-6, dibenzo-18-c-6) are the only low-oxidation state compounds reported to undergo distinct C-Cl bond insertion with DCM and chloroform.²³

Since the formation of compound **IV** by Si–Cl oxidative addition of SiCl₄ to LGa was previously reported,⁸ we became interested to study the selectivity of Si–H ν s. Si–Cl bond insertion by reaction of LGa with HSiCl₃. Addition of HSiCl₃ to a solution of LGa in benzene afforded a colorless solution, from which the Si–Cl bond activation product L(Cl)GaSiHCl₂ (4) was isolated as a colorless crystalline solid in good yield (Scheme 1). *In situ* ¹H NMR spectroscopy proved the selective formation of 4 with no observable signals corresponding to a potential Si–H bond activation product. Compound 4 is stable in the solid-state and in solution at ambient temperature in the absence of air and moisture even after prolonged storage. Similar Si–Cl bond activations of HSiCl₃ were observed by a vinyl silylsilylene²⁴ and a diamidosilylene,^{22b} albeit in the latter case with low selectivity.

Intrigued by the consecutive Si-Br bond activation of compound **V** by reaction with LGa, the reactions of compounds **1–4** and **IV** with one equivalent of LGa were investigated. Compound **1** reacted with LGa at 100 °C within 7 hours *via* C– Cl bond insertion and formation of a rare Ga methylene complex, $[L(Cl)Ga]_2CH_2$ (5), which was isolated in moderate yield after recrystallization from *n*-hexane (Scheme 2). In contrast, compounds **2** and **3** reacted with LGa at 60 °C and



Scheme 2 Synthesis of compounds 5–7.

ambient temperature, but the reactions only resulted in formation of LGaCl₂ as the main product and several minor unidentified species as shown by *in situ* ¹H NMR spectroscopy. The large steric pressure in the hypothetically formed products [L(Cl)Ga]₂C(X)Cl (X = H, Cl) in conjunction with the presence of a reactive C–Cl bond most likely induces the elimination of LGaCl₂ concomitant with reduction of the carbon center, thus accounting for the several side products observed.

Compound 5, which is exceptionally stable in both solidstate and solution under inert gas atmosphere, belongs to the very short list of structurally characterized group 13 element methylene complexes, which includes the parent Ga methylene $[Ga_8(\mu-CH_2)_{12}]$,²⁵ Al methylene $[R_2Al]_2CH_2$ (R = Cl, CH(SiMe_3)_2, 2,4,6-i-Pr_3C_6H_2, N(Me_2CCH_2)_2CH_2, N(SiMe_3)_2),²⁶ and In methylene complexes $[(L)X_2In]_2CH_2$ (X = Cl, Br; L = $Me_2NC_2H_4NMe_2$),²⁷ respectively. In addition, several methylene complexes were obtained from two-fold C–Cl bond activation of DCM by silylenes,²⁸ stannylenes,²⁹ and a digermene,³⁰ respectively.

Gallylsilanes 4 and II were found to react with LGa at 80 °C with Si-Cl bond insertion to afford digallylsilanes [L(Cl) Ga]₂SiHCl (6) and [L(Cl)Ga]₂SiCl₂ (7) in good yields after recrystallization from benzene. In the reaction of LGa with 4, selective Si-Cl bond activation occurred and no indications for a Si-H activated product were observed. Fischer and co-workers⁸ only obtained compound IV even in the presence of excess LGa as conversion of IV to 7 requires thermal treatment to initiate the second bond insertion as was observed for V.14 Most conveniently, compounds 6 and 7 are synthesized by direct reactions of two equivalents of LGa with HSiCl₃ and SiCl₄, respectively. Both compounds, 6 and 7, are stable in the solid-state and in solution at ambient temperature in the absence of air and moisture. Apart from compound V,¹⁴ two-fold bond insertion into geminal Si-halogen bonds has only been observed by a cyclic digermene via initial Si-X addition to the double bond.³¹ Attempts to dehydrochlorinate 6 by addition of strong bases as well as further reduction of 7 using LGa or other reducing agents to obtain the corresponding silvlene failed, showing the decisive influence of the Si-X bond strength.

Spectroscopic characterization

The ¹H and ¹³C NMR spectra of compounds 1–5 and 7 feature the expected signals for the β -diketiminate ligand in reduced $C_{\rm s}$ symmetry due to the presence of two additional different substituents at each Ga center, *e.g.*, two singlets (γ -CH, CCH₃), two septets $(CH(CH_3)_2)$, and four doublets $(CH(CH_3)_2)$ were observed in the ¹H NMR spectra. In contrast, compound 6 exhibits signals corresponding to a C_1 -symmetric β -diketiminate ligand, *e.g.*, three singlets (γ -CH, CCH₃), four septets $(CH(CH_3)_2)$, and eight doublets $(CH(CH_3)_2)$ in the ¹H NMR spectrum, resulting from the heteroleptic substitution pattern at the central Si atom. The CH₂Cl (2.62 ppm) and $CHCl_2$ (5.31 ppm) resonances of 1 and 2, respectively, are observed at higher chemical shift compared to those of CH_2Cl_2 (4.27 ppm) and $CHCl_3$ (6.15 ppm) in C_6D_6 due to the substitution of an electronegative Cl atom by an electropositive Ga

atom. Virtually identical ¹H NMR chemical shifts were observed for L'SiCH_{3-x}Cl_x complexes containing a related N,N'chelating ligand L' (x = 1, 2.68 ppm; x = 2, 5.34 ppm),^{22b} whereas the shift to higher frequency is less pronounced in cationic Cl-In- $CH_{3-x}Cl_x$ crown ether complexes (x = 1, 3.47, 4.37 ppm; x = 2, 5.87, 5.65 ppm).²³ Similar higher chemical shifts compared to CH₂Cl₂ (53.5 ppm) and CHCl₃ (77.8 ppm) are observed for the CH₂Cl (25.4 ppm) and CHCl₂ (59.6 ppm) ¹³C NMR resonances of **1** and **2**, respectively, which could only be reliably located by the cross-peaks in the corresponding 2D-¹H-¹³C-HSOC spectra due to their broadness. For the same reason, the CCl_3 resonance of 3 was not observed. The CH_2 (-1.02 ppm) and CH_2 (-12.4 ppm) resonances of Ga methylene complex 5 appear at even higher frequency as a result of the high electron density at the central CH₂ moiety imposed by the two electropositive gallyl substituents, and are comparable to those of THF and pyridine adducts of the parent Ga methylene $[Ga_8(\mu-CH_2)_{12}] (\delta_{^{1}H} 0.63-0.49 \text{ ppm})^{25}$ and $[R_2AI]_2CH_2 (\delta_{^1H} R = CH(SiMe_3)_2 -0.50; 2,4,6-i-Pr_3C_6H_2 -0.66;$ $N(SiMe_3)_2$ –0.67 ppm),²⁶ while those of the double silylene insertion product {[H₂CC(SiMe₃)₂]₂(Cl)Si}₂CH₂ formed by reaction of two equivalents of silvlene [H2CC(SiMe3)2]2Si with dichloromethane are found at lower frequency (δ_{1H} 1.43 ppm; $\delta_{^{13}\text{C}}$ 20.2 ppm).^{28b} In contrast to CH₂Cl₂, **1**, and **5**, the observed ¹H NMR chemical shifts of the SiH protons in HSiCl₃ (5.39 ppm), 4 (5.69 ppm), and 6 (4.30 ppm) show no distinct trend. However, the ${}^{1}J_{HSi}$ coupling constants measured from ²⁹Si satellites gradually decrease from HSiCl₃ (${}^{1}J_{HSi}$ = 371.0 Hz) over 4 (${}^{1}J_{HSi}$ = 244.0 Hz) to 6 (${}^{1}J_{HSi}$ = 178.0 Hz) most likely reflecting the change in hybridization: bonding to Ga will redirect the Si s character to that electropositive atom, which increases the p character in the Si-H bond and results in a smaller ¹J_{HSi} coupling constant.³² Moreover, these values are in accordance to those of Cl-Si-SiHCl₂ complexes (5.34 ppm,^{22b} 4.64 ppm (ref. 24)) formed in Si-Cl bond insertion of HSiCl₃ with silvlenes. Unfortunately, no meaningful ²⁹Si NMR spectra of compound 7 was obtained due to its low solubility in organic solvents, whereas the ²⁹Si spectra of 4 and 6 show resonces at 9.0 ppm (${}^{1}J_{SiH}$ = 244 Hz, 4) and -30.5 ppm (¹ J_{Si-H} = 178 Hz, 6), respectively. The γ -*CH* resonances of compounds 1-7 remain largely unaffected by the different substituents at the Ga centers ($\delta_{^1\mathrm{H}}$ 4.79–4.88; $\delta_{^{13}\mathrm{C}}$ 97.7–98.6). The IR absorption bands corresponding to the Si-H stretching mode $(\nu_{\text{Si-H}})$ are found at 2181 cm⁻¹ and 2108 cm⁻¹ for compounds 4 and 6, respectively.

Single-crystal structure analysis

sc-XRD analyses confirmed the molecular structures of 1–7 in the solid state as one- and two-fold C–Cl and Si–Cl bond-activated complexes (Fig. 2, 3, S22 and S23†). Suitable crystals of 1–7 were obtained from saturated solutions in *n*-pentane (1), *n*-hexane (2, 5), benzene (4, 6, 7), and DCM/*n*-hexane (3). Compounds 2 and 4 as well as 3 and IV^8 crystallize isomorphous in the orthorhombic space group *Pnma* and the monoclinic space group $P2_1/m$, respectively, while compound 1 crystallizes in the monoclinic space group $P2_1/n$ with two indepen**Dalton Transactions**

View Article Online



Fig. 2 Molecular structures of 1, 2, and 3 in the solid-state. Only one of the two independent molecules of 1 is shown. Hydrogen atoms (except CH_2Cl and CH_2Cl and $CHCl_2$) and the minor component of the disordered CH_2Cl molety were omitted for clarity. Displacement ellipsoids are drawn at 50% probability level, whereas hydrogen atoms are displayed as spheres of arbitrary radius. Symmetry generated parts in 2 and 3 are depicted with pale inner lines.



Fig. 3 Molecular structures of 5 and 6 in the solid-state. Hydrogen atoms (except CH₂ and SiHCl), the minor component of the disordered SiHCl moiety, and co-crystallized solvent molecules were omitted for clarity. Displacement ellipsoids are drawn at 50% probability level, whereas hydrogen atoms are displayed as spheres of arbitrary radius. Symmetry generated parts in 5 are depicted with pale inner lines.

dent yet similar molecules in the asymmetric unit, in which the CH₂Cl and Ga(Cl)CH₂Cl moieties are disordered over two positions, respectively. The SiHCl moiety in compound **6**, which crystallized in the monoclinic space group $P2_1/c$, is dis-

Table 1 Selected bond lengths [Å] and angles [°] of compounds 1–7

ordered over two positions. Digallyl complexes 5 and 7 crystallize in the monoclinic space group C2/c and the triclinic space group $P\bar{1}$, respectively, and both incorporate co-crystallized solvent molecules (n-hexane 5, benzene 7). Selected bond lengths and angles are summarized in Table 1. The LGa moieties in the molecular structures of 1-7 feature the typical metrics observed in complexes of the type LGa(X)Y, *i.e.*, the Ga atoms are located significantly out of the C₃N₂ planes of the ligand backbones. The Ga atoms adopt distorted tetrahedral coordination geometries with Cl-Ga-E bond angles ranging from $105.15(4)^{\circ}$ to $117.90(3)^{\circ}$, which is slightly larger than the values observed in digallyl complexes 5-7. The Ga-C and Ga-Cl bond lengths of 1-3 are virtually identical to those of L(Cl) GaMe (Ga-C 1.956(2) Å, Ga-Cl 2.223(1) Å),³³ I (Ga-C 2.029(8) Å, Ga-Cl 2.251(2) Å),8 and L(Cl)GaCp (Ga-C 2.0006(17) Å, Ga-Cl 2.1887(5) Å),³⁴ and the C–Cl bond lengths agree with those of related Si-CH₂Cl (1.758 Å) and Si-C(Me)Cl₂ (1.773 Å, 1.775 Å) complexes.^{22b} The Ga-Cl bond lengths in 4 are similar to those of 1-3, and the Ga-Si and Si-Cl bonds lengths correspond to those of compounds IV (Ga-Si 2.3860(13) Å, Si-Cl 1.999(2) Å, 2.0135(15) Å)⁸ and V (Ga-Si 2.3992(14) Å).¹⁴ The

	$1^{a,b}$	2	3	4	5	6 ^b	7
Ga–Cl	2.230(14)	2.1933(5)	2.1928(5)	2.2016(5)	2.2264(4)	2.2131(4) 2.2173(5)	2.2171(3) 2.2249(4)
Ga-E	2.035(3)	2.0172(19)	2.0150(17)	2.3809(6)	1.9413(8)	2.4540(14) 2.3632(14)	2.4240(4) 2.4168(4)
Ga–N	1.938(3) 1.899(3)	1.9283(9)	1.9233(9)	1.9281(9)	$\begin{array}{c} 1.9628(10) \\ 1.9481(10) \end{array}$	1.9548(12) 1.9556(12) 1.9546(12) 1.9519(12)	$1.9565(9) \\ 1.9537(9) \\ 1.9480(9) \\ 1.9494(10)$
E-Cl	1.753(2)	1.7731(12)	1.7617(19) 1.7827(11)	2.0528(5)	—	2.0948(17)	2.0807(5) 2.0775(5)
Cl-Ga-E	114.2(7)	110.58(6)	106.84(6)	109.56(2)	117.90(3)	116.20(4) 105.15(4)	115.11(2) 113.69(2)
N–Ga–E	113.0(8) 110.8(7)	115.04(4)	118.10(4)	117.00(3)	109.96(5) 124.78(3)	$117.81(5) \\118.18(5) \\112.90(5) \\131.26(5)$	$122.68(3) \\114.65(3) \\116.68(3) \\121.50(3)$
Ga–E–Ga	—	—	—	—	127.16(9)	123.01(6)	129.09(2)

^{*a*} Average values of the two independent molecules. ^{*b*} Values given for the major component of disorder only.

[L(Cl)Ga] ligands in digallyl complexes 5-7 adopt twisted syn,syn conformations with respect to the Cl atoms, and the compounds feature Ga-E-Ga bond angles ranging from 123.01(6)° to 129.09(2)° comparable to those reported for other complexes of the type $[L(X)Ga]_2E$ (E = PBr,³⁵ InEt,¹⁵ SnMe₂,⁸ SiH₂,¹⁴ SiBr₂;¹⁴ Ga-E-Ga 120.01°-129.83°). Interestingly, the Ga-C bonds in 5 are considerably shorter than those in 1-3, while the Ga-Cl and Ga-N bonds are slightly elongated, which can be explained by a negative hyperconjugative interaction from the C-H bond orbital to the antibonding Ga-Cl/N orbitals $(\sigma_{\rm C-H} \rightarrow \sigma^*_{\rm Ga-Cl/N})$ as were observed in related complexes containing electron-rich pnictogen centers instead of the CH₂ group.^{16i,36} The Ga–C bond lengths in 5 are comparable to those of $[Ga_8(\mu-CH_2)_{12}]$ (1.960(2) Å, 1.961(1) Å, 1.972(2) Å).²⁵ The Ga–Si bond lengths in 7 agree with those of compound V (2.4247(5) Å, 2.4344(5) Å),¹⁴ while those of **6** significantly differ from each other (by 0.09 Å) with one longer and one shorter Ga-Si bond. The shorter Ga-Si bond in 6 is close to that of [L(Br)Ga]₂SiH₂ (2.3788(7) Å),¹⁴ which suggests uneven steric repulsion between the H and Cl, and the [L(Cl)Ga] ligands, respectively, leading to the observed distortions of bond lengths. The Ga-Cl and Si-Cl bonds in 6 and 7 are similar to those of compound 4 and IV (Ga-Cl 2.1980(11) Å, Si-Cl 1.999(2) Å, 2.0135(15) Å), respectively.⁸

Conclusions

Gallanediyl LGa reacts with chloromethane solvents and chlorosilanes selectively with E-Cl bond activation, yielding oxidative addition products $L(Cl)GaCH_xCl_{3-x}$ (x = 0 (3), 1 (2), 2 (1)) and L(Cl)GaSiH_xCl_{3-x} (x = 0 (**IV**), 1 (4)). Consecutive insertion of LGa into a geminal E-Cl bond furnished digallyl complexes $[L(Cl)Ga]_2CH_2$ (5), $[L(Cl)Ga]_2SiHCl$ (6), and $[L(Cl)Ga]_2SiCl_2$ (7), with 5 representing a rare Ga methylene complex. Hence, the scope of C-X (X = F, Cl, Br, I) bond activation reactions mediated by Ga(I) complexes was extended to compounds containing geminal C-X bonds. Moreover, the influence of the Si-X bond strength was disclosed as [L(Cl) Ga]₂SiCl₂ failed to give a silvlene carbonyl complex analogous to V. Further studies on the C-X bond activation of polyhaloalkanes by Ga(1) as well as the utilization of carbenoids 1-3 as precursors for the release of halogenated and functionalized carbenes, and the use of 5 as methylene transfer reagent in synthetic transformations are under current investigation in our laboratory.

Experimental

General procedures, materials, and instrumentation

All manipulations were carried out under an atmosphere of purified argon using standard Schlenk and glovebox techniques. Toluene, *n*-hexane, and *n*-pentane were dried with a MBraun Solvent Purification System (SPS), and benzene was distilled from Na/K alloy. Dichloromethane and chloroform were distilled from CaH₂ and CaCl₂, respectively, while carbon tetrachloride was dried with molecular sieves (4 Å) and degassed. Deuterated benzene was dried over activated molecular sieves (4 Å) and degassed prior to use. LGa³⁷ was prepared according to literature procedures. SiCl₄ and HSiCl₃ were obtained from commercial sources and used as received. NMR spectra (δ in ppm) were recorded using a Bruker Avance DPX 300 (¹H 300.1 MHz, ¹³C{¹H} 75.5 MHz) or a Bruker Avance Neo 400 (¹H 400.1 MHz, ${}^{13}C{}^{1}H{}$ 100.6 MHz) spectrometer and were referenced to internal C₆D₅H (¹H δ = 7.16, ¹³C δ = 128.06). IR spectra were recorded in a glovebox with an ALPHA-T FT-IR spectrometer equipped with a single-reflection ATR sampling module. Microanalyses were performed at the Elemental Analysis Laboratory of the University of Duisburg-Essen. Melting points were measured in wax-sealed glass capillaries under argon atmosphere using a Thermo Scientific 9300 apparatus and are uncorrected.

Synthesis of L(Cl)GaCH₂Cl (1)

LGa (50 mg, 0.1026 mmol) was dissolved in CH₂Cl₂ (2 mL) yielding a colorless solution. Volatiles were removed in vacuo and the residue was dissolved in n-pentane (0.8 mL). Storage of the solution at ambient temperature overnight afforded colorless analytically pure crystals of 1. Yield: 38 mg (0.0664 mmol, 65%). Mp: 177 °C (dec.). Anal. calcd for C₃₀H₄₃Cl₂GaN₂: C, 62.96; H, 7.57; N, 4.99. Found: C, 63.1; H, 7.58; N, 5.17. ¹H NMR (400.1 MHz, C_6D_6): δ 7.13 (m, 4 H, $C_6H_3(i-Pr)_2$), 7.03 (m, 2 H, $C_6H_3(i-Pr)_2$), 4.83 (s, 1 H, γ -CH), 3.74 (sept, ${}^3J_{HH} = 6.7$ Hz, 2 H, $CH(CH_3)_2$), 3.18 (sept, ${}^{3}J_{HH} = 6.8$ Hz, 2 H, $CH(CH_3)_2$), 2.62 (s, 2 H, CH_2Cl), 1.53 (s, 6 H, CCH_3), 1.50 (d, ${}^{3}J_{HH}$ = 6.8 Hz, 6 H, CH(CH₃)₂), 1.25 (d, ${}^{3}J_{HH}$ = 6.8 Hz, 6 H, CH(CH₃)₂), 1.19 (d, ${}^{3}J_{\text{HH}} = 6.7 \text{ Hz}, 6 \text{ H}, \text{CH}(\text{CH}_{3})_{2}), 1.03 \text{ (d, }{}^{3}J_{\text{HH}} = 6.7 \text{ Hz}, 6 \text{ H}, \text{CH}$ $(CH_3)_2$). ¹³C{¹H} NMR (100.6 MHz, C₆D₆): δ 170.4 (CCH₃), 145.8 $(C_6H_3(i-Pr)_2)$, 143.1 $(C_6H_3(i-Pr)_2)$, 140.0 $(C_6H_3(i-Pr)_2)$, 127.9 $(C_6H_3(i-Pr)_2)$, 125.5 $(C_6H_3(i-Pr)_2)$, 124.0 $(C_6H_3(i-Pr)_2)$, 97.8 $(\gamma-CH)$, 29.0 $(CH(CH_3)_2)$, 28.3 $(CH(CH_3)_2)$, 26.6 $(CH(CH_3)_2)$, 25.4 (CH₂Cl), 25.0 (CH(CH₃)₂), 24.5 (CH(CH₃)₂), 24.0 (CH(CH₃)₂), 23.4 (CCH₃). IR (neat): v 2964, 2962, 2867, 1526, 1438, 1383, 1316, 1260, 1178, 1021, 935, 872, 800, 758, 553, 450 cm⁻¹.

Synthesis of L(Cl)GaCHCl₂ (2)

LGa (50 mg, 0.1026 mmol) was dissolved in CHCl₃ (2 mL) yielding a colorless solution. Volatiles were removed *in vacuo* and the residue was dissolved in hot *n*-hexane (1 mL). Storage of the solution at ambient temperature overnight afforded colorless analytically pure crystals of 2. Yield: 48 mg (0.0791 mmol, 77%). Mp: 241 °C (dec.). Anal. calcd for $C_{30}H_{42}Cl_3GaN_2$: C, 59.39; H, 6.98; N, 4.62. Found: C, 59.3; H, 6.89; N, 4.79. ¹H NMR (400.1 MHz, C₆D₆): δ 7.13 (m, 4 H, C₆H₃(i-Pr)₂), 7.01 (m, 2 H, C₆H₃(i-Pr)₂), 5.31 (s, 1 H, CHCl₂), 4.79 (s, 1 H, γ -CH), 3.71 (sept, ³J_{HH} = 6.7 Hz, 2 H, CH(CH₃)₂), 1.55 (d, ³J_{HH} = 6.8 Hz, 2 H, CH(CH₃)₂), 1.55 (d, ³J_{HH} = 6.8 Hz, 6 H, CH(CH₃)₂), 1.51 (s, 6 H, CCH₃), 1.26 (d, ³J_{HH} = 6.8 Hz, 6 H, CH(CH₃)₂), 1.19 (d, ³J_{HH} = 6.7 Hz, 6 H, CH(CH₃)₂), 0.99 (d, ³J_{HH} = 6.7 Hz, 6 H, CH(CH₃)₂), 142.6 (C₆H₃(i-Pr)₂),

Dalton Transactions

140.1 $(C_6H_3(i-Pr)_2)$, 125.6 $(C_6H_3(i-Pr)_2)$, 123.8 $(C_6H_3(i-Pr)_2)$, 97.7 $(\gamma$ -CH), 59.6 $(CHCl_2)$, 29.4 $(CH(CH_3)_2)$, 28.2 $(CH(CH_3)_2)$, 26.4 $(CH(CH_3)_2)$, 25.0 $(CH(CH_3)_2)$, 24.6 $(CH(CH_3)_2)$, 23.7 $(CH(CH_3)_2)$, 23.4 (CCH_3) . IR (neat): ν 3060, 2965, 2926, 2867, 1528, 1439, 1378, 1314, 1254, 1178, 1021, 934, 873, 799, 760, 684, 519, 453 cm⁻¹.

Synthesis of L(Cl)GaCCl₃ (3)

An excess of CCl₄ (0.1 mL) was added to a solution of LGa (200 mg, 0.4104 mmol) in toluene (5 mL) at -40 °C. The solution was warmed to ambient temperature and volatiles were removed in vacuo. The residue was dissolved in warm dichloromethane and stored at -30 °C overnight to afford colorless analytically pure crystals of 3. Yield: 95 mg (0.1481 mmol, 36%). Mp: 158 °C (dec.). Anal. calcd for C₃₀H₄₁Cl₄GaN₂: C, 56.20; H, 6.45; N, 4.37. Found: C, 56.35; H, 6.31; N, 4.46. ¹H NMR (400.1 MHz, C_6D_6): 7.12 (m, 4 H, $C_6H_3(i-Pr)_2$), 7.04 (m, 2 H, C₆ H_3 (i-Pr)₂), 4.88 (s, 1 H, γ -CH), 3.80 (sept, ${}^{3}J_{HH}$ = 6.7 Hz, 2 H, $CH(CH_3)_2$), 3.40 (sept, ${}^{3}J_{HH}$ = 6.8 Hz, 2 H, $CH(CH_3)_2$), 1.56 (s, 6 H, CCH₃), 1.52 (d, ${}^{3}J_{HH} = 6.8$ Hz, 6 H, CH(CH₃)₂), 1.30 (d, ${}^{3}J_{\text{HH}} = 6.8 \text{ Hz}, 6 \text{ H}, \text{CH}(\text{CH}_{3})_{2}), 1.18 \text{ (d, }{}^{3}J_{\text{HH}} = 6.7 \text{ Hz}, 6 \text{ H}, \text{CH}$ $(CH_3)_2$, 1.00 (d, ${}^{3}J_{HH}$ = 6.7 Hz, 6 H, $CH(CH_3)_2$). ${}^{13}C{}^{1}H$ NMR (100.6 MHz, C₆D₆): δ 172.0 (CCH₃), 145.7 (C₆H₃(i-Pr)₂), 143.2 $(C_6H_3(i-Pr)_2)$, 140.5 $(C_6H_3(i-Pr)_2)$, 125.5 $(C_6H_3(i-Pr)_2)$, 124.0 $(C_6H_3(i-Pr)_2)$, 98.5 (γ -CH), 29.8 (CH(CH_3)_2), 28.3 (CH(CH_3)_2), 25.6 (CH(CH₃)₂), 23.8 (CH(CH₃)₂), 25.0 (CH(CH₃)₂), 24.0 (CCH₃) (CCl₃ not observed). IR (neat): v 3050, 2954, 2917, 2857, 1521, 1430, 1378, 1310, 1255, 1174, 1016, 933, 872, 793, 754, 710, 687, 635, 530, 441 cm⁻¹.

Synthesis of L(Cl)GaSiHCl₂ (4)

HSiCl₃ (27 mg, 0.1990 mmol, 20 µL) was added to a solution of LGa (97 mg, 0.1990 mmol) in benzene (1 mL) and the solution was stirred for 1 h at ambient temperature. Storage of the solution at 8 °C overnight afforded colorless analytically pure crystals of 4. Yield: 80 mg (0.1280 mmol, 65%). Mp: 190 °C (dec.). Anal. calcd for C₂₉H₄₂Cl₃GaN₂Si: C, 55.93; H, 6.80; N, 4.50. Found: C, 56.0; H, 6.75; N, 4.53. ¹H NMR (300.1 MHz, C_6D_6): δ 7.18–7.03 (m, 6 H, $C_6H_3(i-Pr)_2$), 5.69 (s, satellite: ${}^1J_{HSi} = 244.0$ Hz, 1 H, Si*H*), 4.85 (s, 1 H, γ-C*H*), 3.73 (sept, ${}^{3}J_{HH}$ = 6.6 Hz, 2 H, $CH(CH_3)_2$), 3.16 (sept, ${}^{3}J_{HH}$ = 6.9 Hz, 2 H, $CH(CH_3)_2$), 1.54 (s, 6 H, CCH₃), 1.51 (d, ${}^{3}J_{HH}$ = 6.6 Hz, 6 H, CH(CH₃)₂), 1.33 (d, ${}^{3}J_{\text{HH}}$ = 6.9 Hz, 6 H, CH(CH₃)₂), 1.18 (d, ${}^{3}J_{\text{HH}}$ = 6.9 Hz, 6 H, CH $(CH_3)_2$, 1.00 (d, ${}^{3}J_{HH}$ = 6.9 Hz, 6 H, $CH(CH_3)_2$). ${}^{13}C{}^{1}H$ NMR (C₆D₆, 75.5 MHz): δ 170.2 (CCH₃), 146.1 (C₆H₃(i-Pr)₂), 142.3 $(C_6H_3(i-Pr)_2)$, 140.2 $(C_6H_3(i-Pr)_2)$, 128.2 $(C_6H_3(i-Pr)_2)$, 125.8 $(C_6H_3(i-Pr)_2)$, 124.1 $(C_6H_3(i-Pr)_2)$, 98.0 $(\gamma$ -CH), 29.8 $(CH(CH_3)_2)$, 28.2 (CH(CH₃)₂), 26.8 (CH(CH₃)₂), 24.9 (CH(CH₃)₂), 24.5 (CH(CH₃)₂), 23.8 (CH(CH₃)₂), 23.2 (CCH₃). ²⁹Si NMR (119 MHz, C_6D_6 , DEPT90): 9.0 (¹ J_{HSi} = 244.0 Hz). IR (neat): ν 2966, 2923, 2864, 2181, 1525, 1462, 1435, 1380, 1314, 1252, 1177, 1099, 1021, 932, 873, 794, 756, 638, 592, 541, 513, 435 cm⁻¹.

Synthesis of [L(Cl)Ga]₂CH₂ (5)

A solution of LGa (90 mg, 0.1573 mmol) and 1 (77 mg, 0.1573 mmol) in benzene (1 mL) was heated to 100 $^{\circ}$ C for 7 h.

Volatiles were removed in vacuo and the residue was dissolved in hot *n*-hexane (5 mL). Storage of the solution at -30 °C overnight afforded colorless analytically pure crystals of 5. Yield: 80 mg (0.0755 mmol, 48%). Mp: 358 °C. Anal. calcd for C₅₉H₈₄Cl₂Ga₂N₄·C₆H₁₄: C, 68.13; H, 8.62; N, 4.89. Found: C, 68.1; H, 8.37; N, 4.86. ¹H NMR (400.1 MHz, C₆D₆): δ 7.19 (dd, $J_{\rm HH}$ = 7.6, 1.6 Hz, 4 H, C₆ H_3 (i-Pr)₂), 7.13 (t, ${}^{3}J_{\rm HH}$ = 7.6 Hz, 4 H, $C_6H_3(i-Pr)_2)$, 6.98 (dd, J_{HH} = 7.6, 1.6 Hz, 4 H, $C_6H_3(i-Pr)_2)$, 4.88 (s, 2 H, γ -CH), 3.65 (sept, ${}^{3}J_{HH}$ = 6.7 Hz, 4 H, CH(CH₃)₂), 2.93 (sept, ${}^{3}J_{HH} = 6.7$ Hz, 4 H, CH(CH₃)₂), 1.45 (s, 12 H, CCH₃), 1.25 $(d, {}^{3}J_{HH} = 6.7 \text{ Hz}, 12 \text{ H}, CH(CH_{3})_{2}), 1.20 (d, {}^{3}J_{HH} = 6.7 \text{ Hz}, 12 \text{ H},$ $CH(CH_3)_2$, 0.98 (d, ${}^{3}J_{HH}$ = 6.7 Hz, 12 H, $CH(CH_3)_2$), 0.94 (d, ${}^{3}J_{\text{HH}} = 6.7 \text{ Hz}, 12 \text{ H}, \text{CH}(\text{CH}_{3})_{2}), -1.02 \text{ (s, 2 H, CH}_{2}). {}^{13}\text{C}{}^{1}\text{H}$ NMR (C_6D_6 , 100.6 MHz): δ 168.9 (CCH_3), 145.9 ($C_6H_3(i-Pr)_2$), 142.1 $(C_6H_3(i-Pr)_2)$, 142.0 $(C_6H_3(i-Pr)_2)$, 127.0 $(C_6H_3(i-Pr)_2)$, 125.5 $(C_6H_3(i-Pr)_2)$, 123.9 $(C_6H_3(i-Pr)_2)$, 97.9 $(\gamma$ -CH), 28.9 $(CH(CH_3)_2)$, 28.4 $(CH(CH_3)_2)$, 27.4 $(CH(CH_3)_2)$, 25.1 $(CH(CH_3)_2)$, 24.7 (CH(CH₃)₂), 24.6 (CH(CH₃)₂), 24.2 (CCH₃), -12.4 (CH₂). IR (neat): v 2953, 2916, 2857, 1520, 1430, 1377, 1312, 1254, 1174, 1095, 1018, 993, 933, 862, 793, 757, 684, 550, 497, 449 cm⁻¹.

Synthesis of [L(Cl)Ga]₂SiHCl (6)

HSiCl₃ (14 mg, 0.0990 mmol, 10 µL) was added to a solution of LGa (97 mg, 0.1990 mmol) in benzene (1 mL) and the solution was heated to 80 °C for 2 days. Storage of the solution at 8 °C overnight afforded colorless analytically pure crystals of 6. Yield: 67 mg (0.0600 mmol, 61%). Mp: 244 °C (dec.). Anal. calcd for C₅₈H₈₃Cl₃Ga₂N₄Si: C, 62.75; H, 7.54; N, 5.05. Found: C, 62.78; H, 7.57; N, 5.15. ¹H NMR (300.1 MHz, C_6D_6): δ 7.19–7.02 (m, 12 H, $C_6H_3(i-Pr)_2$), 4.88 (s, 2 H, γ -CH), 4.30 (s, satellite: ${}^{1}J_{HSi}$ = 178.0 Hz, 1 H, Si*H*), 3.78 (sept, ${}^{3}J_{HH}$ = 6.9 Hz, 2 H, $CH(CH_3)_2$, 3.54 (sept, ${}^{3}J_{HH} = 6.6$ Hz, 2 H, $CH(CH_3)_2$), 3.18 (m, 4 H, CH(CH₃)₂), 1.50 (s, 6 H, CCH₃), 1.49 (s, 6 H, CCH₃), 1.30 (d, ${}^{3}J_{HH} = 6.6$ Hz, 6 H, CH(CH₃)₂), 1.29 (d, ${}^{3}J_{HH} = 6.9$ Hz, 6 H, $CH(CH_3)_2$, 1.23 (d, ${}^{3}J_{HH}$ = 6.9 Hz, 6 H, $CH(CH_3)_2$), 1.21 (d, ${}^{3}J_{HH}$ = 6.9 Hz, 6 H, CH(CH₃)₂), 1.19 (d, ${}^{3}J_{HH}$ = 6.9 Hz, 6 H, CH $(CH_3)_2$, 1.15 (d, ${}^{3}J_{HH}$ = 6.6 Hz, 6 H, $CH(CH_3)_2$), 0.99 (d, ${}^{3}J_{HH}$ = 6.6 Hz, 6 H, CH(CH₃)₂), 0.95 (d, ${}^{3}J_{HH}$ = 6.6 Hz, 6 H, CH(CH₃)₂). $^{13}C{^{1}H}$ NMR (C₆D₆, 75.5 MHz): δ 169.6 (CCH₃), 169.3 (CCH₃), 146.2 $(C_6H_3(i-Pr)_2)$, 146.0 $(C_6H_3(i-Pr)_2)$, 143.2 $(C_6H_3(i-Pr)_2)$, 142.4 $(C_6H_3(i-Pr)_2)$, 142.3 $(C_6H_3(i-Pr)_2)$, 141.2 $(C_6H_3(i-Pr)_2)$, 127.7 $(C_6H_3(i-Pr)_2)$, 127.5 $(C_6H_3(i-Pr)_2)$, 125.3 $(C_6H_3(i-Pr)_2)$, 125.0 $(C_6H_3(i-Pr)_2)$, 123.7 $(C_6H_3(i-Pr)_2)$, 123.5 $(C_6H_3(i-Pr)_2)$, 98.6 (γ-CH), 29.8 (CH(CH₃)₂), 29.5 (CH(CH₃)₂), 28.3 (CH(CH₃)₂), 28.1 $(CH(CH_3)_2)$, 27.9 $(CH(CH_3)_2)$, 27.8 $(CH(CH_3)_2)$, 25.1 (CH(CH₃)₂), 24.9 (CH(CH₃)₂), 24.4 (CH(CH₃)₂), 24.3 (CH(CH₃)₂), 24.1 (CH(CH₃)₂), 24.0 (CH(CH₃)₂), 23.6 (CCH₃), 23.3 (CCH₃). ²⁹Si NMR (119 MHz, C₆D₆, DEPT90): –30.5 (¹J_{HSi} = 178.0 Hz). IR (neat): v 2958, 2922, 2866, 2108, 1521, 1434, 1382, 1314, 1255, 1176, 1100, 1017, 937, 866, 799, 759, 735, 683, 636, 501, 441, 410 cm^{-1} .

Synthesis of [L(Cl)Ga]₂SiCl₂ (7)

SiCl₄ (15 mg, 0.0870 mmol, 10 μ L) was added to a solution of LGa (85 mg, 0.1740 mmol) in benzene (1 mL) and the solution was heated to 80 °C for 3 days. Storage of the solution at 8 °C

overnight afforded colorless analytically pure crystals of 7. Yield: 78 mg (0.0680 mmol, 78%). Mp: 208 °C (dec.). Anal. calcd for C₅₈H₈₂Cl₄Ga₂N₄Si: C, 60.86; H, 7.22; N, 4.89. Found: C, 60.83; H, 7.28; N, 4.81. ¹H NMR (300.1 MHz, C_6D_6): δ 7.20-7.03 (m, 12 H, C₆H₃(i-Pr)₂), 4.85 (s, 2 H, γ-CH), 3.67 (sept, ${}^{3}J_{\text{HH}}$ = 6.9 Hz, 4 H, CH(CH₃)₂), 3.23 (sept, ${}^{3}J_{\text{HH}}$ = 6.9 Hz, 4 H, $CH(CH_3)_2$, 1.46 (s, 12 H, CCH₃), 1.33 (d, ${}^{3}J_{HH}$ = 6.6 Hz, 12 H, $CH(CH_3)_2$, 1.32 (d, ${}^{3}J_{HH}$ = 6.6 Hz, 12 H, $CH(CH_3)_2$), 1.23 (d, ${}^{3}J_{\text{HH}}$ = 6.9 Hz, 12 H, CH(CH₃)₂), 0.94 (d, ${}^{3}J_{\text{HH}}$ = 6.6 Hz, 12 H, CH(CH₃)₂). ¹³C 1 H} NMR (C₆D₆, 75.5 MHz): δ 170.0 (CCH₃), 145.9 $(C_6H_3(i-Pr)_2)$, 143.0 $(C_6H_3(i-Pr)_2)$, 142.4 $(C_6H_3(i-Pr)_2)$, 127.7 $(C_6H_3(i-Pr)_2)$, 125.2 $(C_6H_3(i-Pr)_2)$, 123.8 $(C_6H_3(i-Pr)_2)$, 98.6 $(\gamma$ -CH), 29.6 (CH(CH₃)₂), 28.2 (CH(CH₃)₂), 27.9 (CH(CH₃)₂), 25.0 (CH(CH₃)₂), 24.6 (CH(CH₃)₂), 24.1 (CH(CH₃)₂), 24.0 (CCH₃). A ²⁹Si NMR spectrum of 7 could not be obtained due to its poor solubility. IR (neat): v 2957, 2924, 2868, 1521, 1433, 1380, 1312, 1256, 1179, 1098, 1022, 937, 865, 796, 760, 711, 638, 496, 441 cm^{-1} .

Single-crystal X-ray diffraction

The crystals of 1-7 were mounted on nylon loops in inert oil. Crystallographic data of 1 was collected on a Bruker AXS D8 Venture diffractometer with Photon II detector (CuKa radiation, $\lambda = 1.54178$ Å, micro-focus source) at 102(2) K (Table S1[†]). Absorption corrections were performed semiempirically from equivalent reflections on the basis of multiscans (Bruker AXS APEX2). Crystallographic data of 2-7 were collected on a Bruker D8 Kappa diffractometer with APEX2 detector (MoK_{α} radiation, $\lambda = 0.71073$ Å) at 100(2) K and are summarized in Tables S1 and S2.† Absorption corrections were performed semi-empirically from equivalent reflections on the basis of multiscans (Bruker AXS APEX3). The structures were solved by direct methods (SHELXS-97)38 and refined anisotropically by full-matrix least-squares on F^2 (SHELXL-2014).³⁹ Hydrogen atoms were refined using a riding model or rigid methyl groups. In the crystal of 1, the central Ga(Cl)CH₂Cl moiety in residue 1 and the CH₂Cl group in residue 2 are disordered. Two alternate positions each were used for the modelling. In residue 1 a third orientation was identified yet strong parameter correlations made a refinement impossible thus the rather large anisotropic displacement parameters had to be accepted. All corresponding bond lengths of the disordered parts were restrained to be equal (SADI), additionally, the bond angles were restrained (SADI) in residue 1. RIGU restraints were applied to the anisotropic displacement parameters of the disordered atoms of residue 1, and C30_1 and C30'_1 were refined with common displacement parameters (EADP). The crystal of 2 was a non-merohedral twin and the model was refined against de-twinned HKLF4 data. In 4, one Dip group is disordered over two positions. Its bond lengths were restrained with SADI. In 5, an n-hexane molecule is disordered over two positions. All bond lengths and angles were restrained to be equal (SADI) and RIGU restraints were applied to the anisotropic displacement parameters. In 6, the central Si(H)Cl unit is disordered over two positions. The SiH hydrogen atoms were refined freely and the Si-H bond lengths were restrained to be equal (SADI). In 7, one of the four benzene

molecules is disordered over two positions. The ADPs of the others suggest minor disorder. Any attempt to separate different orientations failed. All 1,2 and 1,3 distances of the solvent molecules were restraint to be equal (SADI) and the atoms were restrained to be on a common plane (FLAT). RIGU and ISOR restraints were applied to the atoms of the disorder model.

CCDC-2127285 (1), 2127286 (2), 2127287 (3), 2127288 (4), 2127289 (5), 2127290 (6), 2127291 (7) contain the supplementary crystallographic data for this paper.†

Author contributions

C. H. and C. G. planned and executed all the experimental work and analysed the corresponding data. C. W. conducted the sc-XRD measurements and processed corresponding data. C. H. wrote the manuscript. S. S. supervised the project and revised the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Financial support by the University of Duisburg-Essen, the German Research Foundation DFG (grant no. SCHU 1069/26-1), and Evonik Industries (C. H., doctoral fellowship) is gratefully acknowledged. We thank J. Schoening for performing ²⁹Si NMR measurements with compounds **4** and **6**.

Notes and references

- (a) T. Chu and G. I. Nikonov, *Chem. Rev.*, 2018, **118**, 3608;
 (b) P. P. Power, *Nature*, 2010, **463**, 171;
 (c) C. Weetman and S. Inoue, *ChemCatChem*, 2018, **10**, 4213.
- 2 M. Zhong, S. Sinhababu and H. W. Roesky, *Dalton Trans.*, 2020, **49**, 1351.
- 3 (*a*) N. J. Hardman and P. P. Power, *Chem. Commun.*, 2001, 1866; (*b*) C. Ganesamoorthy, M. Matthias, D. Bläser, C. Wölper and S. Schulz, *Dalton Trans.*, 2016, 45, 11437.
- 4 (a) A. Seifert, D. Scheid, G. Linti and T. Zessin, *Chem. Eur.* J., 2009, 15, 12114; (b) T. Chu, I. Korobkov and G. I. Nikonov, J. Am. Chem. Soc., 2014, 136, 9195; (c) C. Bakewell, A. J. P. White and M. R. Crimmin, *Chem.* Sci., 2019, 10, 2452; (d) H. Zhu, J. Chai, H. Fan, H. W. Roesky, C. He, V. Jancik, H.-G. Schmidt, M. Noltemeyer, W. A. Merrill and P. P. Power, *Angew. Chem.*, *Int. Ed.*, 2005, 44, 5090.
- 5 (a) G. Prabusankar, C. Gemel, P. Parameswaran, C. Flener, G. Frenking and R. A. Fischer, Angew. Chem., Int. Ed., 2009, 48, 5526; (b) L. Tuscher, C. Ganesamoorthy, D. Bläser, C. Wölper and S. Schulz, Angew. Chem., Int. Ed., 2015, 54, 10657; (c) J. Krüger, C. Ganesamoorthy, L. John, C. Wölper

and S. Schulz, *Chem. – Eur. J.*, 2018, 24, 9157; (*d*) J. Krüger, C. Wölper, L. John, L. Song, P. R. Schreiner and S. Schulz, *Eur. J. Inorg. Chem.*, 2019, 1669; (*e*) J. Krüger, C. Wölper and S. Schulz, *Inorg. Chem.*, 2020, **59**, 11142.

- 6 K. M. Korch and D. A. Watson, Chem. Rev., 2019, 119, 8192.
- 7 M. S. Hill, P. B. Hitchcock and R. Pongtavornpinyo, *Inorg. Chem.*, 2007, 46, 3783.
- 8 A. Kempter, C. Gemel and R. A. Fischer, *Inorg. Chem.*, 2008, 47, 7279.
- 9 O. Kysliak, H. Görls and R. Kretschmer, *J. Am. Chem. Soc.*, 2021, **143**, 142.
- (a) M. R. Crimmin, M. J. Butler and A. J. P. White, *Chem. Commun.*, 2015, 51, 15994; (b) T. Chu, Y. Boyko, I. Korobkov and G. I. Nikonov, *Organometallics*, 2015, 34, 5363; (c) C. Bakewell, A. J. P. White and M. R. Crimmin, *Angew. Chem., Int. Ed.*, 2018, 57, 6638.
- 11 G. Prabusankar, A. Kempter, C. Gemel, M.-K. Schröter and R. A. Fischer, *Angew. Chem., Int. Ed.*, 2008, 47, 7234.
- 12 A. Doddi, C. Gemel, M. Winter, R. A. Fischer, C. Goedecke, H. S. Rzepa and G. Frenking, *Angew. Chem., Int. Ed.*, 2013, 52, 450.
- 13 G. Prabusankar, C. Gemel, M. Winter, R. W. Seidel and R. A. Fischer, *Chem. – Eur. J.*, 2010, **16**, 6041.
- 14 C. Ganesamoorthy, J. Schoening, C. Wölper, L. Song, P. R. Schreiner and S. Schulz, *Nat. Chem.*, 2020, **12**, 608.
- 15 C. Ganesamoorthy, D. Bläser, C. Wölper and S. Schulz, *Organometallics*, 2015, **34**, 2991.
- 16 (a) L. Tuscher, C. Ganesamoorthy, D. Bläser, C. Wölper and S. Schulz, Angew. Chem., Int. Ed., 2015, 54, 10657; (b) L. Tuscher, C. Helling, C. Ganesamoorthy, J. Krüger, C. Wölper, W. Frank, A. S. Nizovtsev and S. Schulz, Chem. -Eur. J., 2017, 23, 12297; (c) C. Ganesamoorthy, C. Helling, C. Wölper, W. Frank, E. Bill, G. E. Cutsail III and S. Schulz, Nat. Commun., 2018, 9, 87; (d) C. Helling, C. Wölper and S. Schulz, J. Am. Chem. Soc., 2018, 140, 5053; (e) J. Krüger, C. Ganesamoorthy, L. John, C. Wölper and S. Schulz, Chem. - Eur. J., 2018, 24, 9157; (f) C. Helling, C. Wölper, Y. Schulte, G. Cutsail III and S. Schulz, Inorg. Chem., 2019, 58, 10323; (g) J. Krüger, C. Wölper, L. John, L. Song, P. R. Schreiner and S. Schulz, Eur. J. Inorg. Chem., 2019, 1669; (h) J. Schoening, L. John, C. Wölper and S. Schulz, Dalton Trans., 2019, 48, 17729; (i) C. Helling, G. E. Cutsail III, H. Weinert, C. Wölper and S. Schulz, Angew. Chem., Int. Ed., 2020, 59, 7561; (j) J. Krüger, C. Wölper and S. Schulz, Inorg. Chem., 2020, 59, 11142; (k) C. Helling, C. Wölper and S. Schulz, Dalton Trans., 2020, 49, 11835; (l) C. Helling, C. Wölper and S. Schulz, Eur. J. Inorg. Chem., 2020, 4225; (m) B. Li, C. Wölper, G. Haberhauer and S. Schulz, Angew. Chem., Int. Ed., 2021, 60, 1986; (n) B. Li, C. Wölper, G. Haberhauer and S. Schulz, Angew. Chem., Int. Ed., 2021, 60, 1986; (o) M. K. Sharma, C. Wölper, G. Haberhauer and S. Schulz, Angew. Chem., Int. Ed., 2021, 60, 6784; (p) H. M. Weinert, C. Wölper, J. Haak, G. E. Cutsail III and S. Schulz, Chem. Sci., 2021, 12, 14024.
- 17 S. Schulz, C. Ganesamoorty, G. Bendt, D. Bläser and C. Wölper, *Dalton Trans.*, 2015, **44**, 5153.

- 18 M. Stender, B. E. Eichler, N. J. Hardman, P. P. Power, J. Prust, M. Noltemeyer and H. W. Roesky, *Inorg. Chem.*, 2001, 40, 2794.
- (a) D. A. Dickie, P. T. K. Lee, O. A. Labeodan, G. Schatte, N. Weinberg, A. R. Lewis, G. M. Bernard, R. E. Wasylishen and J. A. C. Clyburne, *Dalton Trans.*, 2007, 2862;
 (b) C. Drost, P. B. Hitchcock and M. F. Lappert, *Organometallics*, 2002, 21, 2095.
- 20 (a) A. Rit, R. Tirfoin and S. Aldridge, Angew. Chem., Int. Ed., 2016, 55, 378; (b) M. M. D. Roy, S. Fujimori, M. J. Ferguson, R. McDonald, N. Tokitoh and E. Rivard, Chem. Eur. J., 2018, 24, 14392.
- 21 (a) N. Burford, P. Losier, P. K. Bakshi and T. S. Cameron, Dalton Trans., 1993, 201; (b) N. Burford, P. Losier, C. Macdonald, V. Kyrimis, P. K. Bakshi and T. S. Cameron, Inorg. Chem., 1994, 33, 1434.
- 22 (a) D. C. H. Do, A. V. Protchenko, M. A. Fuentes, J. Hicks, P. Vasko and S. Aldridge, *Chem. Commun.*, 2020, 56, 4684;
 (b) Y. Xiong, S. Yao and M. Driess, *Organometallics*, 2009, 28, 1927.
- 23 (a) C. G. Andrews and C. L. B. Macdonald, Angew. Chem., Int. Ed., 2005, 44, 7453; (b) B. F. T. Cooper, C. G. Andrews and C. L. B. Macdonald, J. Organomet. Chem., 2007, 692, 2843; (c) C. Peppe, D. G. Tuck, F. M. de Andrade, J. A. Nóbrega, M. A. Brown and R. A. Burrow, J. Organomet. Chem., 2005, 690, 925.
- 24 M. M. D. Roy, M. J. Ferguson, R. McDonald, Y. Zhou and E. Rivard, *Chem. Sci.*, 2019, **10**, 6476.
- 25 M. Bonath, C. Maichle-Mössmer, P. Sirsch and R. Anwander, *Angew. Chem., Int. Ed.*, 2019, **58**, 8206.
- 26 (a) H. Lehmkuhl and R. Schäfer, *Tetrahedron Lett.*, 1966, 7, 2315; (b) M. Layh and W. Uhl, *Polyhedron*, 1990, 9, 277; (c) W. Uhl, C. Rösener, C. Stefaniak, T. Choy, B. Jasper-Peter, J. Kösters, M. Layh and A. Hepp, *Z. Naturforsch., B: J. Chem. Sci.*, 2012, 67, 1081; (d) K. Knabel, H. Nöth and T. Seifert, *Z. Naturforsch., B: J. Chem. Sci.*, 2002, 57, 830.
- 27 M. A. Khan, C. Peppe and D. G. Tuck, *Organometallics*, 1986, 5, 525.
- 28 (a) B. Gehrhus and M. F. Lappert, J. Organomet. Chem., 2001, 617–618, 209; (b) S. Ishida, T. Iwamoto, C. Kabuto and M. Kira, Chem. Lett., 2001, 30, 1102; (c) S. S. Sen, J. Hey, D. Kratzert, H. W. Roesky and D. Stalke, Organometallics, 2012, 31, 435.
- 29 M. J. S. Gynane, M. F. Lappert, S. J. Miles, A. J. Carty and N. J. Taylor, *Dalton Trans.*, 1977, 2009.
- 30 V. Y. Lee, H. Yasuda, M. Ichinohe and A. Sekiguchi, *Angew. Chem., Int. Ed.*, 2005, **44**, 6378.
- 31 T. Sugahara, T. Sasamori and N. Tokitoh, *Angew. Chem., Int. Ed.*, 2017, **56**, 9920.
- 32 (a) W. Scherer, P. Meixner, K. Batke, J. E. Barquera-Lozada, K. Ruhland, A. Fischer, G. Eickerling and K. Eichele, Angew. Chem., Int. Ed., 2016, 55, 11673; (b) P. Meixner, K. Batke, A. Fischer, D. Schmitz, G. Eickerling, M. Kalter, K. Ruhland, K. Eichele, J. E. Barquera-Lozada, N. P. M. Casati, F. Montisci, P. Macchi and W. Scherer, J. Phys. Chem. A, 2017, 121, 7219.

- 33 S. Singh, H.-J. Ahn, A. Stasch, V. Jancik, H. W. Roesky, A. Pal, M. Biadene, R. Herbst-Irmer, M. Noltemeyer and H.-G. Schmidt, *Inorg. Chem.*, 2006, 45, 1853.
- 34 C. Helling, C. Wölper, G. E. Cutsail III, G. Haberhauer and S. Schulz, *Chem. – Eur. J.*, 2020, 26, 13390.
- 35 L. Tuscher, C. Helling, C. Wölper, W. Frank, A. S. Nizovtsev and S. Schulz, *Chem. – Eur. J.*, 2018, **24**, 3241.
- 36 (a) C. Helling, C. Wölper and S. Schulz, *Dalton Trans.*, 2020,
 49, 11835; (b) J. Krüger, C. Wölper and S. Schulz, *Angew. Chem., Int. Ed.*, 2021, 60, 3572.
- 37 N. J. Hardman, B. E. Eichler and P. P. Power, *Chem. Commun.*, 2000, 1991.
- 38 G. M. Sheldrick, Acta Crystallogr., Sect. A: Found. Crystallogr., 1990, 46, 467.
- 39 (a) G. M. Sheldrick, SHELXL-2014, Program for the Refinement of Crystal Structures, University of Göttingen, Göttingen, Germany, 2014. See also; G. M. Sheldrick, Acta Crystallogr., Sect. A: Found. Crystallogr., 2008, 64, 112(b) shelXle, A Qt GUI for SHELXL: C. B. Hübschle, G. M. Sheldrick and B. Dittrich, J. Appl. Crystallogr., 2011, 44, 1281.