Food & Function

CORRECTION

Check for updates

Cite this: Food Funct., 2022, 13, 2372

Correction: Gut microbial metabolite urolithin B attenuates intestinal immunity function *in vivo* in aging mice and *in vitro* in HT29 cells by regulating oxidative stress and inflammatory signalling

Peng Chen,^a Fuchao Chen,^b Jiexin Lei^c and Benhong Zhou*^{a,d}

DOI: 10.1039/d2fo90008d

Correction for 'Gut microbial metabolite urolithin B attenuates intestinal immunity function *in vivo* in aging mice and *in vitro* in HT29 cells by regulating oxidative stress and inflammatory signalling' by Peng Chen *et al., Food Funct.,* 2021, **12**, 11938–11955, DOI: 10.1039/D1FO02440J.

The authors regret that incorrect versions of Fig. 5 and 8 were included in the original article. The panel for occludin 100 mg kg⁻¹ Uro B was incorrect in Fig. 5A. The panel for HE 300 mg kg⁻¹ Uro B was incorrect in Fig. 8A, and the panel for occludin 450 mg kg⁻¹ Uro B was incorrect in Fig. 8B. The correct figures are presented below.

^aKey Laboratory of Combinatorial Biosynthesis and Drug Discovery, Ministry of Education, Wuhan University School of Pharmaceutical Sciences, Wuhan University, Wuhan, Hubei, P. R. China. E-mail: benhongzh@whu.edu.cn; Tel: +15335898431

^bSinopharm Dongfeng General Hospital, Hubei University of Medicine, China

^cDepartment of Endocrinology, Renmin Hospital of Wuhan University, Wuhan, Hubei, 430060, P. R. China

^dDepartment of Pharmacy, Renmin Hospital of Wuhan University, Wuhan, Hubei, 430060, P. R. China



View Article Online



Fig. 5 (A) Representative HE and Masson staining of the small intestine in d-gal-induced aging mice. (B) Immunohistochemistry of ZO-1, occludin, and MUC2 in the small intestine of each group. (C) Protein expression of ZO-1, occludin, and MUC2 was detected using western blotting. All results are described as means \pm SD (n = 12). #P < 0.05 and ##P < 0.01 vs. the control group; *P < 0.05 and **P < 0.01 vs. the d-gal-induced aging group.



Fig. 8 (A) Representative HE and Masson staining of the small intestine in normal mice. (B) Immunohistochemistry of ZO-1, occludin, and MUC2 in the small intestine of each group. All results are described as means \pm SD (n = 12). n.s. represents no statistical significance.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.