



Cite this: *Food Funct.*, 2022, **13**, 3077

Correction: Chestnut polysaccharides restore impaired spermatogenesis by adjusting gut microbiota and the intestinal structure

Zhong-Yi Sun,^a Shuai Yu,^b Yu Tian,^c Bao-Quan Han,^b Yong Zhao,^d Ya-Qi Li,^e Yan Wang,^b Yu-Jiang Sun^{c,f} and Wei Shen^{*c}

DOI: 10.1039/d2fo90009b

rsc.li/food-function

Correction for 'Chestnut polysaccharides restore impaired spermatogenesis by adjusting gut microbiota and the intestinal structure' by Zhong-Yi Sun *et al.*, *Food Funct.*, 2022, **13**, 425–436, DOI: 10.1039/D1FO03145G.

The authors regret that the panel for busulfan MVH in Fig. 1 was shown incorrectly in the original article. The correct version of Fig. 1 is shown below.




Fig. 1 Chestnut polysaccharides (CPs) increased germ cell quantity. Histopathology photos of HE staining and MVH and DAZL staining of mouse testes.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aUrology Department, Shenzhen University General Hospital, Shenzhen 518055, China

^bUrology Department, Peking University Shenzhen Hospital, Shenzhen 518036, China

^cCollege of Life Sciences, Key Laboratory of Animal Reproduction and Biotechnology in Universities of Shandong, Qingdao Agricultural University, Qingdao 266109, China. E-mail: wshen@qau.edu.cn, shenwei427@163.com

^dState Key Laboratory of Animal Nutrition, Institute of Animal Sciences, Chinese Academy of Agricultural Sciences, Beijing, 100000, China

^eUrology Department, Zaozhuang Hospital of Zaozhuang Mining Group, Zaozhuang 277100, China

^fDongying Vocational Institute, Dongying 257091, China

