



Cite this: *Green Chem.*, 2022, **24**, 6319


DOI: 10.1039/d2gc90068h
 rsc.li/greenchem

Correction: Visible-light-mediated minisci C–H alkylation of heteroarenes with 4-alkyl-1,4-dihydropyridines using O₂ as an oxidant

Jianyang Dong,^a Fuyang Yue,^a Wentao Xu,^a Hongjian Song,^a Yuxiu Liu^a and Qingmin Wang^{*a,b}

Correction for 'Visible-light-mediated minisci C–H alkylation of heteroarenes with 4-alkyl-1,4-dihydropyridines using O₂ as an oxidant' by Jianyang Dong *et al.*, *Green Chem.*, 2020, **22**, 5599–5604, <https://doi.org/10.1039/D0GC02111C>.

The authors found that a recently corrected *ACS Catalysis* article reported that the addition of an alkyl radical to 6-chloroimidazo[1,2-*b*]pyridazine should be at the C8 position in Minisci reactions through crystal structure analysis.¹ The authors found that they have a similar NMR spectrum compared with this corrected *ACS Catalysis* article for a 6-chloroimidazo[1,2-*b*]pyridazine compound. The authors regret that the regioselectivity of products **19**, **20** and **21** in Fig. 1 were incorrect. The correct structures/regio-isomers of the originally proposed structures are shown below.



The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

References

- 1 R. A. Garza-Sanchez, A. Tlahuext-Aca, G. Tavakoli and F. Glorius, *ACS Catal.*, 2022, **12**, 6640–6640.

^aState Key Laboratory of Elemento-Organic Chemistry, Research Institute of Elemento-Organic Chemistry, College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China. E-mail: wangqm@nankai.edu.cn

^bCollaborative Innovation Center of Chemical Science and Engineering (Tianjin), Tianjin 300071, People's Republic of China

