

CORRECTION

View Article Online
View Journal | View Issue

Cite this: *Lab Chip*, 2022, 22, 652

Correction: Quantifying nanotherapeutic penetration using a hydrogel-based microsystem as a new 3D *in vitro* platform

Saba Goodarzi,^a Audrey Prunet,^a Fabien Rossetti,^a Guillaume Bort,^a
Olivier Tillement,^a Erika Porcel,^b Sandrine Lacombe,^b Ting-Di Wu,^{cd}
Jean-Luc Guerquin-Kern,^{cd} Hélène Delanoë-Ayari,^a
François Lux^{ae} and Charlotte Rivière^{*ae}

DOI: 10.1039/d1lc90127c

rsc.li/loc

Correction for 'Quantifying nanotherapeutic penetration using a hydrogel-based microsystem as a new 3D *in vitro* platform' by Saba Goodarzi *et al.*, *Lab Chip*, 2021, **21**, 2495–2510, DOI: 10.1039/D1LC00192B.

In Fig. 2E the legend of the graph should be inversed. The corrected legend is: white bars#2D, black bars#3D. The corrected figure is shown below.



^a University of Lyon, Université Claude Bernard Lyon 1, CNRS, Institut Lumière Matière, F-69622, Villeurbanne, France. E-mail: charlotte.riviere@univ-lyon1.fr

^b Université Paris-Saclay, CNRS, Institut des Sciences Moléculaires d'Orsay, 91405, Orsay, France

^c *Institut Curie, Université PSL, Paris, France*

^d Université Paris-Saclay, CNRS, Inserm, Centre d'Imagerie Multimodale, 91401, Orsay, France

^e Institut Universitaire de France (IUF), France




Fig. 2 Quantification of penetration and cellular uptake of AGuIX®-Cy5.5 nanoparticles in HCT-116 tumour spheroids and monolayer cell culture. (A) Representative confocal fluorescence images of HCT-116 spheroids incubated with 0.8, 1.5 and 2 mM concentration of AGuIX®-Cy5.5 for 24 h for four different depths (0, 30, 60 and 90 μm). (B) Mean intensity along with standard deviation (light colors) of AGuIX®-Cy5.5 as a function of the distance from the spheroid periphery (see the orthogonal view in the inset, green = nuclei, red = AGuIX®-Cy5.5) for 0.8 mM (yellow, $N = 73$), 1.5 mM (green, $N = 68$) and 2 mM (red, $N = 121$); three independent experiments. (C) Representative confocal fluorescence images of monolayer HCT-116 cells exposed to AGuIX®-Cy5.5 nanoparticles with 0.8, 1.5 and 2 mM concentration. (D) Quantification of the mean intensity of AGuIX®-Cy5.5 nanoparticles in maximal projection of confocal fluorescence images of monolayer cells after 24 h of incubation with different AGuIX®-Cy5.5 concentrations: 0.8 mM (yellow, $N = 40$), 1.5 mM (green, $N = 40$) and 2 mM (red, $N = 40$); three independent experiments. Error bars represent the standard deviations. (E) Mean and standard deviation of the concentration of Gd ($\text{ppb } \mu\text{m}^{-3}$) uptaken by the cells after incubation with 0.8, 1.5 and 2 mM concentration of AGuIX® for 24 h in HCT-116 cell spheroids and monolayer cell culture measured with ICP-MS ($N = 6$, two independent experiments).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

