



Cite this: *New J. Chem.*, 2022, 46, 17080

Correction: Mechanoresponsive, proteolytically stable and biocompatible supergelators from ultra short enantiomeric peptides with sustained drug release propensity

Radha Rani Mehra,^a Anindya Basu,^b Ryann M. Christman,^c Jeena Harjit,^d Anil K. Mishra,^e Amit K. Tiwari^c and Anita DuttKonar^{*ab}

DOI: 10.1039/d2nj90109a

rsc.li/njc

Correction for 'Mechanoresponsive, proteolytically stable and biocompatible supergelators from ultra short enantiomeric peptides with sustained drug release propensity' by Radha Rani Mehra *et al.*, *New J. Chem.*, 2020, **44**, 6346–6354, <https://doi.org/10.1039/D0NJ00102C>.

The authors regret that Fig. 9 in the original article was incorrect. The correct version of Fig. 9 is included below.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Department of Chemistry, Rajiv Gandhi Technological University, Bhopal, India. E-mail: anitaduttkonar@rgtu.net

^b School of Pharmaceutical Sciences, Rajiv Gandhi Technological University, Bhopal, India

^c University of Toledo, Ohio, USA

^d Truba College Bhopal, India

^e Ujjain Engineering College Ujjain, India






Fig. 9 (A and B) The cell death curve of hydrogelators I and II at various concentrations in different cell lines PANC-1, S-1, HCT-116, and MDAMB-231. The statistical significance was calculated with Student's *t*-test, where (*) is shown at $p < 0.05$. The morphological image is a reflection of MTT assay. It indicates the effect of hydrogelators I and II when treated with four different cell lines PANC-1, HCT-116, MDA-MB 231 and S1 of diversified nature, for 72 h (100 μM concentration of the hydrogelators is shown).

