



Cite this: *Org. Biomol. Chem.*, 2022, **20**, 8756

## Visible-light-mediated aerobic Ritter-type C–H amination of diarylmethanes using DDQ/*tert*-butyl nitrite†

Tianci Li,<sup>a</sup> Jiangyu Yang,<sup>a,b</sup> Xin Yin,<sup>a,b</sup> Jinhua Shi,<sup>a</sup> Qun Cao,<sup>id</sup>\*<sup>c</sup> Miaomiao Hu,<sup>a</sup> Xiaowen Xu,<sup>a</sup> Meichao Li<sup>\*a</sup> and Zhenlu Shen<sup>id</sup>\*<sup>a</sup>

Received 20th September 2022,  
Accepted 14th October 2022

DOI: 10.1039/d2ob01713j

rsc.li/obc

A metal-free photocatalytic Ritter-type C–H amination of unactivated sp<sup>3</sup> carbons using molecular oxygen as a terminal oxidant has been developed. By employing a co-catalytic system of 3-dichloro-4,5-dicyano-1,4-benzoquinone (DDQ) and *tert*-butyl nitrite (TBN), this novel strategy provides a low cost, sustainable and scalable way to synthesise a broad range of secondary amides in moderate to excellent yields under mild conditions.

### Introduction

Catalytic transformation of omnipresent C–H bonds to C–N bonds offers an efficient synthetic strategy for the construction of valuable nitrogen-containing compounds.<sup>1–3</sup> As one of the most popular strategies to construct C–N bonds, the Ritter reaction has been established as a powerful tool for the synthesis of amides,<sup>4–6</sup> which widely exist in modern pharmaceutical molecules.<sup>7,8</sup> The classic Ritter reaction involves the generation of a carbocation from a functionalized substrate (*e.g.*, alcohols or alkenes), which then reacts with a nitrile leading to the formation of an amide product (Fig. 1A).<sup>4–6</sup> Recently, much effort has been directed towards the development of catalytic Ritter-type C–H amination, which allows amides to be selectively synthesised from non-functionalized alkanes, with the carbocation intermediate generated from C–H bond oxidation using a variety of oxidants such as ceric ammonium nitrate (CAN),<sup>9</sup> Selectfluor,<sup>10</sup> hypervalent iodine,<sup>11</sup> iodic acid,<sup>12</sup> 3-dichloro-4,5-dicyano-1,4-benzoquinone (DDQ)<sup>13</sup> and sodium persulfate<sup>14</sup> (Fig. 1B). However, these methods still suffer from high loadings of transition metals, high temperature and low sustainability due to unwanted by-products derived from the use of stoichiometric oxidants. Considering the strive for greener chemical production, molecular oxygen (O<sub>2</sub>) is the most ideal oxidant as it is inexpensive and environmentally friendly.<sup>15,16</sup> In the past 10 years, important advances

have been made in the catalytic aerobic oxidation of C(sp<sup>3</sup>)–H bonds;<sup>17–21</sup> however, to the best of our knowledge, there is no aerobic Ritter type C–H amination method reported to date.

Therefore, developing an environmentally friendly and low-cost metal-free catalytic system for C–H amination is highly desirable. Herein, we report the first photocatalytic aerobic Ritter-type C–H amination for the synthesis of various amides under mild and metal-free conditions using a DDQ/*tert*-butyl nitrite (TBN) catalytic system (Fig. 1C).

### Results and discussion

We began our studies by using the DDQ/TBN/O<sub>2</sub> catalytic system, which has previously been successfully used for the oxidation of C(sp<sup>3</sup>)–H bonds,<sup>22–25</sup> oxidative C–O coupling<sup>26</sup> and C–N coupling reactions.<sup>27–29</sup> Under the optimal conditions, diphenylmethane (**1a**, 0.5 mmol), DDQ (20 mol%), TBN (20 mol%), H<sub>2</sub>O (4 mmol, 8 equiv.) and trifluoroacetic acid (TFA, 5 equiv.) in 2 mL of benzonitrile (**2a**) under an oxygen atmosphere (1 bar) led to the desired product **3aa** in 76% yield under the irradiation of blue LED light for 12 h at room temperature (Table 1, entry 1). When the loading of DDQ was decreased from 20 mol% to 10 mol%, the yield of **3aa** decreased slightly to 74% (Table 1, entry 2). Further decreasing the loading of DDQ or the co-catalyst TBN led to unsatisfactory yields of **3aa** (41–61%, for details see ESI Table S1†). It was found that the solvent played a key role in this catalytic reaction. Attempts to replace benzonitrile with other organic solvents (*e.g.*, dichloroethane, toluene, THF, dioxane) led to poor yields (Table 1, entries 3–6), and no amide product was observed when the reaction was carried out in dimethylformamide (DMF, Table 1, entry 7) and dimethoxyethane


<sup>a</sup>College of Chemical Engineering, Zhejiang University of Technology, Hangzhou, 310014, China. E-mail: limc@zjut.edu.cn, zhenlushen@zjut.edu.cn

<sup>b</sup>Zhejiang Jitai New Materials Co., Ltd, Shaoxing 312369, China

<sup>c</sup>School of Chemistry and Chemical Engineering, Queen's University Belfast, David Keir Building, Belfast, BT9 5AG, UK. E-mail: qcao01@qub.ac.uk

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2ob01713j>





**Fig. 1** (A) Ritter reaction with a carbocation as an intermediate. (B) Representative examples of catalytic Ritter-type amination methods with stoichiometric oxidants. (C) This work: metal-free aerobic photo-catalytic Ritter-type C–H amination.

(DME, Table 1, entry 8). Both DDQ and TBN were essential for this catalytic system. No desired product was formed in the absence of DDQ (Table 1, entry 9), whereas a lower yield of **3aa** (32%) was obtained when TBN was omitted or replaced by other NO donors (Table 1, entry 10 and Table S2 in the ESI†). Control experiment also confirmed that the absence of TFA or replacing TFA with other acids (*e.g.*, acetic acid, aq. HCl, H<sub>2</sub>SO<sub>4</sub>) resulted in lower yields of **3aa** (11–21%, Table 1, entries 11–14), which clearly shows the importance of TFA as an additive. Attempts to carry out the reaction under a N<sub>2</sub> atmosphere or without blue LED light led to a yield of 16% and 8% of **3aa**, respectively (Table 1, entries 15 & 16; for further light source and temperature screening, see Table S2 in the ESI†).

With the optimized reaction conditions in hand, the applicability of these conditions to a range of diarylmethanes was explored (Fig. 2). We found that with 10–20 mol% DDQ and 20 mol% TBN, we could obtain the desired amides in good yields. Having methyl/*tert*-butyl substituents at the *ortho*-, *para*-, and *meta*-positions of diphenylmethane led to the corresponding amide in 64–85% yields (**3ba–3ga**). Unfortunately, diphenylmethane bearing a methoxy group did not participate in the reaction (**3ha**), as the substrate was preferentially oxidized to its corresponding ketone under the opti-

**Table 1** Optimization of reaction conditions


Entry	Variation from the standard conditions	Yields <sup>a</sup> [%]
1	None	76
2	10 mol% DDQ	74
3 <sup>b</sup>	DCE as the solvent + 15 equiv. of <b>2a</b>	61
4 <sup>b</sup>	Toluene as the solvent + 15 equiv. of <b>2a</b>	42
5 <sup>b</sup>	THF as the solvent + 15 equiv. of <b>2a</b>	13
6 <sup>b</sup>	Dioxane as the solvent + 15 equiv. of <b>2a</b>	<1
7 <sup>b</sup>	DMF as the solvent + 15 equiv. of <b>2a</b>	N.D. <sup>c</sup>
8 <sup>b</sup>	DME as the solvent + 15 equiv. of <b>2a</b>	N.D. <sup>c</sup>
9	No DDQ	N.D. <sup>c</sup>
10	No TBN	32
11	No TFA	21
12	CH <sub>3</sub> COOH instead of TFA	52
13 <sup>d</sup>	HCl instead of TFA	28
14 <sup>e</sup>	H <sub>2</sub> SO <sub>4</sub> instead of TFA	11
15	Under N <sub>2</sub> instead of O <sub>2</sub>	16
16	Dark environment	8

Standard conditions: the reaction was performed with **1a** (0.5 mmol), **2a** (2 mL, as a reagent/solvent), DDQ (20 mol%, 0.10 mmol), TBN (20 mol%, 0.10 mmol), TFA (5 equiv., 2.5 mmol), and H<sub>2</sub>O (4 mmol, 8 equiv.) with blue LED light and an oxygen balloon at room temperature for 12 h. <sup>a</sup> Yields were determined by GC using biphenyl as an internal standard. <sup>b</sup> 10 mol% DDQ with a specified solvent (2 mL) was used. <sup>c</sup> N.D. implies that no product could be detected by GC-MS and GC. <sup>d</sup> 35 wt% aq. HCl was used. <sup>e</sup> 98 wt% H<sub>2</sub>SO<sub>4</sub> was used.

mized conditions. It was found that fluoro-, chloro-, and bromo-substituted diphenylmethanes were successfully transformed to their desired amides in fair to good yields (58–76%, **3ia–3oa**). Notably, strong electron withdrawing substituents (*e.g.*, CF<sub>3</sub>– and NO<sub>2</sub>–) were well tolerated in the catalytic system and afforded amides in moderate yields (**3pa**, **3qa**). Although benzylheteroarene was not suitable in our reaction system (**3ra**), benzyl-naphthalenes showed good activities, affording the desired products (**3sa**, **3ta**) in 67% and 64% yields, respectively.


In previous related Ritter-type C–H amination reports,<sup>4–9,30</sup> the scope of nitriles was less studied and often limited to acetonitrile and benzonitrile. With a broad range of diarylmethanes examined, we tested the system on a variety of nitriles. Excellent functional group tolerance was exhibited by a wide range of nitriles as shown in Fig. 3. We were delighted that benzonitriles bearing an electron donating group (CH<sub>3</sub>–) and halogens (F–, Cl–, Br–) could participate in the reaction to furnish their corresponding amides in good to very good yields (**3ab–3ad**, **3ae–3ag**). Furan and thiophene derivatives (*e.g.*, **3ah** and **3ai**) could also be prepared, albeit in moderate yields, highlighting the applicability of this method to synthesize substituted heterocycles. It was found that  $\alpha,\beta$ -unsaturated nitriles could also react successfully, as exemplified by the preparation of **3aj** and **3ak**. Additionally, this system was applied to aliphatic nitriles; they were transformed into their corresponding *N*-benzhydryl amides (**3al**, **3am**, **3an**) in good yields.






**Fig. 2** Scope of diarylmethanes for the aerobic photocatalytic Ritter-type C–H amination reaction using the DDQ/TBN system. Reaction conditions: **1** (0.5 mmol), **2a** (2 mL, as a reagent/solvent), DDQ (10–20 mol% as specified), TBN (20 mol%, 0.10 mmol), TFA (5 equiv., 2.5 mmol) and H<sub>2</sub>O (8 equiv., 4 mmol) with blue LED light and an oxygen balloon at room temperature for 12 h. Unless otherwise noted, isolated yields are shown. <sup>a</sup> 10 mol% DDQ was used. <sup>b</sup> 20 mol% DDQ was used. <sup>c</sup> (4-Methoxyphenyl)(phenyl)methanone was isolated in 89% yield. <sup>d</sup> 87% **1r** was recovered.

To further explore the utility of our Ritter-type amination method, we expanded the reaction to a gram scale under the optimized conditions with 10 mol% DDQ and 20 mol% TBN as the co-catalyst (Fig. 4). This work thus far highlights the improved performance of the system compared with the previous system, given the low cost of the catalyst, sustainable mild reaction conditions, and the absence of any metal salts and stoichiometric oxidants.




**Fig. 3** Scope of nitriles for the aerobic photocatalytic Ritter-type C–H amination reaction using the DDQ/TBN system. Standard conditions: **1** (0.5 mmol), **2** (2 mL, as a reagent/solvent), DDQ (20 mol%, 0.10 mmol), TBN (20 mol%, 0.10 mmol), TFA (5 equiv., 2.5 mmol) and H<sub>2</sub>O (8 equiv., 4 mmol) with blue LED light and an oxygen balloon at room temperature for 12 h. Unless otherwise noted, isolated yields are shown. <sup>a</sup> Reaction was carried out with 15 equiv. of the corresponding nitrile in 2 mL of DCE. <sup>b</sup> 42% **1a** was recovered. <sup>c</sup> 40% **1a** was recovered.



**Fig. 4** Gram scale reaction.

To gain insights into the reaction mechanism, several control experiments were carried out (Fig. 5). When 2 equiv. of 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) were added as a radical scavenger, the photocatalytic oxidation reaction was completely inhibited and an adduct (**4**) of the N-radical from **1a** with TEMPO was observed by HPLC-MS analysis (Fig. 5a and Fig. S1 in the ESI<sup>†</sup>). These results suggested that the reaction involved a radical pathway with a carbon centered radical being involved as a key intermediate, which agrees with the mechanism proposed for previously reported Ritter type C–H amination methods.<sup>9–14</sup>






**Fig. 5** Control experiments. <sup>a</sup> Yields were determined by GC using biphenyl as an internal standard. <sup>b</sup> Reaction was carried out using 3 Å molecular sieves pre-dried **1a**, **2a** and TBN without the addition of H<sub>2</sub>O. <sup>c</sup> Water content was determined by Karl-Fischer titration.

It was reported that the DDQ/TBN system could be employed for visible-light photocatalytic aerobic benzylic C(sp<sup>3</sup>)-H oxygenations with alcohols being formed as either intermediates or final products,<sup>31</sup> which could then serve as substrates for the Ritter reaction.<sup>4–6</sup> In addition, it was found that a small amount of **2a** could be hydrolysed to benzylamide (**7**) under the standard reaction conditions (Fig. S2 in the ESI†). To investigate the roles of alcohols and amides, **1a** was first replaced with diphenylmethanol (**5**) as the starting material, which led to only 14% yield of **3aa** with benzophenone (**6**) formed as the major product in 57% yield (Fig. 5b). In addition, the catalytic reaction with benzamide (**7**) instead of **2a** as the starting material led to a sluggish reaction with 21% yield of **3aa** obtained (Fig. 5c). These observations indicate that alcohols or amides generated *in situ* are not involved as the major intermediates during the catalytic cycle. Consistent with previous reports,<sup>14</sup> it was found that the amount of water also played an important role in the catalytic cycle. When the amount of water in the catalytic system was decreased from 8 equiv. to 0.32 equiv., the yield of **3aa** decreased to 55% (Fig. 5d).

According to the above experimental results and literature reports,<sup>14,28,31,32</sup> a mechanism exemplified by the reaction of diphenylmethane (**1a**) with benzonitrile (**2a**) is proposed as shown in Fig. 6. Initial photolysis and acidolysis of TBN liberate NO, which can be readily oxidised to NO<sub>2</sub> by O<sub>2</sub>.<sup>31,33,34</sup> Under visible light irradiation, DDQ is excited to its triplet excited state (<sup>3</sup>DDQ\*).<sup>35</sup> The hydrogen atom transfer (HAT) reaction between **1a** and <sup>3</sup>DDQ\* generates DDQH• and an alkyl radical (**I**),<sup>36,37</sup> which undergoes subsequent single electron transfer (SET) and leads to the formation of diphenylmethyl cation (**II**) and DDQH<sup>-</sup>.<sup>31,37</sup> The carbocation (**II**) then reacts with



**Fig. 6** The proposed mechanism of the aerobic photocatalytic Ritter-type C–H amination using the DDQ/TBN system.

**2a** to give the nitrilium ion intermediate (**III**), which undergoes hydrolysis to the corresponding amide (**3a**). Meanwhile, DDQH<sup>-</sup> is protonated to the hydroquinone DDQH<sub>2</sub>, which can be re-oxidized to DDQ by NO<sub>2</sub>.<sup>38,39</sup>

## Conclusions

In summary, by using the DDQ/TBN catalytic system, we have developed the first photocatalytic metal-free Ritter-type C–H amination method with molecular oxygen as a green oxidant. With high functional group tolerance and improved nitrile substrate scope, this protocol provides a simple, sustainable, and scalable method for the construction of secondary C–N bonds under mild conditions with moderate to excellent selectivities. We expect this new protocol to complement existing amination methods and to expand the toolbox for C–H functionalization of unreactive C(sp<sup>3</sup>)-H bonds.

## Author contributions

ZS and ML conceived the project. ZS, ML and QC designed the methodology and supervised the project. TL, JY, XY, JS, MH and XX performed synthetic and catalytic experiments. TL, QC and ZS analysed the data and wrote the manuscript.

## Conflicts of interest

CN Patent CN113943199A by Z. Shen *et al.* contains intellectual property described in this article. The other authors (*e.g.*, J. Yang, X. Yin, J. Shi, Q. Cao, M. Hu and X. Xu) have no conflict of interest to declare.



## Acknowledgements

This work was financially supported by the National Natural Science Foundation of China (22178321, 21773211 and 21776260) and the Key Research and Development Project of Zhejiang Province (no. 2021C01079).

## Notes and references

- 1 Y. Park, Y. Kim and S. Chang, *Chem. Rev.*, 2017, **117**, 9247–9301.
- 2 J. L. Roizen, M. E. Harvey and J. Du Bois, *Acc. Chem. Res.*, 2012, **45**, 911–922.
- 3 M.-L. Louillat and F. W. Patureau, *Chem. Soc. Rev.*, 2014, **43**, 901–910.
- 4 G. Mohammadi Ziarani, F. Soltani Hasankiadeh and F. Mohajer, *ChemistrySelect*, 2020, **5**, 14349–14379.
- 5 D. Jiang, T. He, L. Ma and Z. Wang, *RSC Adv.*, 2014, **4**, 64936–64946.
- 6 A. Guérinot, S. Reymond and J. Cossy, *Eur. J. Org. Chem.*, 2012, 19–28.
- 7 X. Wang, *Nat. Catal.*, 2019, **2**, 98–102.
- 8 C. L. Allen and J. M. J. Williams, *Chem. Soc. Rev.*, 2011, **40**, 3405–3415.
- 9 S. Sakaguchi, T. Hirabayashi and Y. Ishii, *Chem. Commun.*, 2002, 516–517.
- 10 Q. Michaudel, D. Thevenet and P. S. Baran, *J. Am. Chem. Soc.*, 2012, **134**, 2547–2550.
- 11 G.-X. Li, C. A. Morales-Rivera, F. Gao, Y. Wang, G. He, P. Liu and G. Chen, *Chem. Sci.*, 2017, **8**, 7180–7185.
- 12 K. Kiyokawa, K. Takemoto and S. Minakata, *Chem. Commun.*, 2016, **52**, 13082–13085.
- 13 Y. Zhang, J. Dong, L. Liu, L. Liu, Y. Zhou and S.-F. Yin, *Org. Biomol. Chem.*, 2017, **15**, 2897–2901.
- 14 H. Lai, J. Xu, J. Lin, B. Su and D. Zha, *Org. Chem. Front.*, 2022, **9**, 1541–1549.
- 15 Z. Shi, C. Zhang, C. Tang and N. Jiao, *Chem. Soc. Rev.*, 2012, **41**, 3381–3430.
- 16 T. Punniyamurthy, S. Velusamy and J. Iqbal, *Chem. Rev.*, 2005, **105**, 2329–2364.
- 17 H. Sterckx, B. Morel and B. U. W. Maes, *Angew. Chem., Int. Ed.*, 2019, **58**, 7946–7970.
- 18 H. Y. Kim and K. Oh, *Org. Biomol. Chem.*, 2021, **19**, 3569–3583.
- 19 J. Xie, C. Pan, A. Abdukader and C. Zhu, *Chem. Soc. Rev.*, 2014, **43**, 5245–5256.
- 20 Y. Bian, Y. Gu, X. Zhang, H. Chen and Z. Li, *Chin. Chem. Lett.*, 2021, **32**, 2837–2840.
- 21 Y. Ma, F. Gao, W. Xiao, N. Li, S. Li, B. Yu and X. Chen, *Chin. Chem. Lett.*, 2022, **33**, 4395–4399.
- 22 Z. Shen, J. Dai, J. Xiong, X. He, W. Mo, B. Hu, N. Sun and X. Hu, *Adv. Synth. Catal.*, 2011, **353**, 3031–3038.
- 23 J. Ma, Z. Hu, M. Li, W. Zhao, X. Hu, W. Mo, B. Hu, N. Sun and Z. Shen, *Tetrahedron*, 2015, **71**, 6733–6739.
- 24 D. Pan, Y. Wang, M. Li, X. Hu, N. Sun, L. Jin, B. Hu and Z. Shen, *Synlett*, 2019, 218–224.
- 25 Z. Shen, T. Li, J. Ma and M. Li, CN113943199A, 2021.
- 26 D. Pan, Z. Pan, Z. Hu, M. Li, X. Hu, L. Jin, N. Sun, B. Hu and Z. Shen, *Eur. J. Org. Chem.*, 2019, 5650–5655.
- 27 C. Song, X. Dong, H. Yi, C.-W. Chiang and A. Lei, *ACS Catal.*, 2018, **8**, 2195–2199.
- 28 C. Song, H. Yi, B. Dou, Y. Li, A. K. Singh and A. Lei, *Chem. Commun.*, 2017, **53**, 3689–3692.
- 29 S. Das, P. Natarajan and B. König, *Chem. – Eur. J.*, 2017, **23**, 18161–18165.
- 30 T. Shen and T. H. Lambert, *J. Am. Chem. Soc.*, 2021, **143**, 8597–8602.
- 31 F. Rusch, J.-C. Schober and M. Brasholz, *ChemCatChem*, 2016, **8**, 2881–2884.
- 32 Y. Zhang, W. Chen, L. Wang and P. Li, *Org. Chem. Front.*, 2018, **5**, 3562–3566.
- 33 J. Piera and J.-E. Bäckvall, *Angew. Chem., Int. Ed.*, 2008, **47**, 3506–3523.
- 34 T. Ting, R. Stanger and T. Wall, *Int. J. Greenhouse Gas Control*, 2013, **18**, 15–22.
- 35 K. Ohkubo, A. Fujimoto and S. Fukuzumi, *J. Am. Chem. Soc.*, 2013, **135**, 5368–5371.
- 36 X. Guo, H. Zipse and H. Mayr, *J. Am. Chem. Soc.*, 2014, **136**, 13863–13873.
- 37 V. S. Batista, R. H. Crabtree, S. J. Konezny, O. R. Luca and J. M. Praetorius, *New J. Chem.*, 2012, **36**, 1141–1144.
- 38 E. Bosch, R. Rathore and J. K. Kochi, *J. Org. Chem.*, 1994, **59**, 2529–2536.
- 39 R. Rathore, E. Bosch and J. K. Kochi, *Tetrahedron Lett.*, 1994, **35**, 1335–1338.

