Organic & Biomolecular Chemistry



View Article Online

PAPER

Check for updates

Cite this: Org. Biomol. Chem., 2023, **21**, 402

Hydrogen bond templated synthesis of catenanes and rotaxanes from a single isophthalic acid derivative[†]

Hydrogen bond templated [2]catenanes and [2]rotaxanes have been synthesized using azide precursors derived from a single isophthalic acid derivative precursor. The interlocked molecules were prepared

using either stoichiometric or near stoichiometric amounts of macrocycle and CuAAC "click" precursors,

with yields of up to 70% for the mechanical bond formation step. Successful preparation of the inter-

locked structures was confirmed by NMR spectroscopy and mass spectrometry, with detail of co-confor-

Sean R. Barlow, ២ Geoffrey R. Akien ២ and Nicholas H. Evans 🕩 *

mational behaviour being elucidated by a range of ¹H NMR spectroscopic experiments.

Received 4th November 2022, Accepted 5th December 2022

DOI: 10.1039/d2ob02019j

rsc.li/obc

Introduction

Mechanically interlocked molecules such as catenanes¹ (consisting of interlocked macrocyclic rings) and rotaxanes² (macrocyclic rings(s) trapped on stoppered axles) have unusual properties arising from the mechanical bond such as the potential for the controlled, large amplitude motion of the interlocked components and the 3D cavities and spaces arising from their exotic architectures.³

An array of template methodologies have been used to prepare interlocked molecules, including but not limited to, metal cations,⁴ π – π stacking (or π -donor/ π -acceptor interactions),⁵ hydrogen bonding⁶ and anions.⁷ However, the bespoke nature of many syntheses of catenanes and rotaxanes is often a drawback when trying to exploit their properties and potential applications. Development of modular methodologies based on easy to access building blocks is desirable.

While our group has reported upon the rapid hydrogen bond templated synthesis of [2]rotaxanes in good yield,^{8,9} our previous serendipitous [2]catenane synthesis is low yielding (12%) with regard to the crucial covalent capture step that forms the interlocked compound.^{10,11} We hypothesized that with design modification to allow for more efficient hydrogen bonding templation, that synthesis of [2]catenanes in higher yields should be possible.

Indeed, here in we report the successful synthesis of three novel [2]catenanes derived from a simple isophthalic acid precursor in good yield. Furthermore, starting from the same isophthalic acid derivative allowed for the synthesis of two novel [2]rotaxanes in reasonable yield. All interlocked molecules were characterized by NMR and IR spectroscopy and mass spectrometry.

Results and discussion

Synthesis and characterization of catenanes

The catenane syntheses all build upon azido-functionalized isophthalic acid derivative **1** (Scheme 1), which may be rapidly prepared from commercially available mono-methyl ester of isophthalic acid.^{12,13}

Three azide–alkyne coupling partners were prepared. Initially DCC/*N*-hydroxysuccinimide-mediated amide coupling of alkyne carboxylic acids $2-4^{14-16}$ with commercially available 4-(Boc-amino)benzylamine afforded alkynes 5–7. These were subsequently deprotected using TFA, with removal of the Boc group in quantitative yield confirmed by ¹H NMR spectroscopy.

The alkyne–azide precursor targets **11–13** were synthesized *via* the coupling of the amine trifluoroacetate salts **8–10** with isophthalic acid derivative **1** using DCC and *N*-hydroxysuccinimide. Following workup and purification by silica gel column chromatography, **11–13** were isolated in reasonable yields (51–64%). The successful preparation of novel compounds **11–13** was confirmed by NMR spectroscopy and mass spectrometry (see ESI, pp. S9 and S10; S14 and S15; S19 and S20†).

Mechanical bond formation was accomplished *via* the threading and cyclization of alkyne-azides **11–13** through previously reported macrocycle **14**.¹⁰ In each case, to a solution of **14** in dry CH₂Cl₂, **1.0** equivalents of **11–13** was added and allowed to stir for 15 minutes to facilitate pseudorotaxane for-

Department of Chemistry, Lancaster University, Lancaster, LA1 4YB, UK.

E-mail: n.h.evans@lancaster.ac.uk

[†]Electronic supplementary information (ESI) available: Further experimental procedures; copies of spectral characterization data. See DOI: https://doi.org/ 10.1039/d2ob02019j

Paper



Scheme 1 Synthesis of [2]catenanes 15–17.

mation (Scheme 1). Then, catalytic $Cu(CH_3CN)_4BF_4$ and TBTA (tris((1-benzyl-4-triazolyl)methyl)amine), and 1.2 equivalents of *N*,*N*-diisopropylamine were added. The reactions were stirred overnight at room temperature under an inert atmosphere, and then submitted to aqueous workup and purification by silica gel column chromatography. Novel heterocir-

cuit [2]catenanes 15-17^{17,18} were isolated in good yields (59-70%). The [2]catenanes were characterized by ¹H & ¹³C NMR and IR spectroscopy, with molecular ions being detected by high resolution mass spectrometry (see ESI, pp. S21-S26†).

Catenane formation is evident upon comparison of the ¹H NMR spectra of precursor **13**, macrocycle **14** and [2]catenane

Paper

17 (Fig. 1). The upfield shift and splitting of aromatic protons 14/15 and h/i in catenane 17 compared to 13 and 14 is consistent with the intercalation of aromatic rings within the interlocked structure. The downfield shift of internal isophthalamide proton d is indicative of interactions with a hydrogen bond acceptor on the cyclized thread. Another notable characteristic of the ¹H NMR spectra of catenanes 15–17 is the splitting of certain protons (*e.g. f* and *k*) due to the two faces of the rotationally symmetric macrocycle becoming inequivalent due to the directionality of the newly formed rotationally asymmetric macrocyclic ring.

Further evidence of the interlocked nature of the macrocyclic rings is provided by a molecular ion peak being identifiable by positive ion electrospray mass spectrometry (see ESI, pp. S22, S24 & S26†). In addition, for each catenane there is the appearance of multiple through-space correlations in the ¹H–¹H ROESY NMR spectra between resonances arising from protons in the two interlocked macrocycles (*e.g.* Fig. 2 and see ESI, pp. S39–



Fig. 1 1 H NMR spectra of (a) alkyne–azide 13, (b) [2]catenane 17 and (c) macrocycle 14 (1 : 1 CDCl₃/CD₃OD, 400 MHz, 298 K).



Fig. 2 Section of ${}^{1}H-{}^{1}H$ ROESY NMR spectrum of [2]catenane 17 with intercomponent through-space correlations highlighted (1:1 CDCl₃/ CD₃OD, 400 MHz, 298 K).

S41[†]). Inspection of the entire spectrum in each case reveals sufficient intercomponent correlations to indicate the macrocyclic rings of the catenane are switching between multiple co-conformations in 1:1 CDCl₃/CD₃OD (see ESI, p. S42[†]).

To probe potential dynamic interactions (*e.g.* the pirouetting of the interlocked macrocyclic rings) the most soluble [2] catenane, glycol catenane 17, was studied by variable temperature (VT) ¹H NMR spectroscopy in $C_2D_2Cl_4$ (Fig. 3). At (and below) room temperature, several C-H resonances are broad, but these sharpen upon heating. Three of the amide N-H resonances not only sharpen but move upfield, while one (δ) has almost no change in chemical shift, indicating that this amide N-H is participating in intramolecular hydrogen bonding.

Synthesis and characterization of rotaxanes

The azido-functionalized isophthalic acid derivative **1** can also be derivatized with bulky stopper groups to allow for the formation of rotaxanes (Scheme 2). Novel half-axle components





18 and **19** were prepared in good yield by reaction of 3,5-bis(trifluoromethyl)benzylamine and 3,5-bis(trifluoromethyl)-aniline with isophthalic acid derivative **1** activated using DCC and *N*-hydroxysuccinimide.

To prepare [2]rotaxanes 21 and 23, 1.1 equivalents of 18 or 19 and alkyne 20⁸ were added to a solution of macrocycle 14 in dry CH₂Cl₂, followed by catalytic Cu(CH₃CN)₄BF₄ and TBTA, and 1.2 equivalents of *N*,*N*-diisopropylamine. Following aqueous workup and purification of the crude material by preparative TLC, [2]rotaxanes 21 and 23 were isolated in 29% and 40% yields respectively.¹⁹ While the isolated yields are reasonable, it should be noted the actual yields of rotaxane formation are almost certainly higher. Isolation of pure rotaxane was hindered in both cases by incomplete separation of product bands during chromatographic purification.

Successful formation of the [2]rotaxanes was confirmed by analysis of NMR spectra and detection of molecular ion peaks in high resolution mass spectrometry (see ESI, pp. S31–S34†). The ¹H NMR spectrum of rotaxane **21**, along with that of noninterlocked macrocycle **14** and axle **22** for comparison, is shown in Fig. 4. Once again, the upfield shift and splitting of



Fig. 4 ¹H NMR spectra of (a) axle 22, (b) [2]rotaxane 21 and (c) macrocycle 14 (CDCl₃, 400 MHz, 298 K).

aromatic protons *h* and *i* in the interlocked molecule compared to non-interlocked macrocycle **14** is consistent with presence of the second component (here an axle) between the aromatic rings of the macrocycle. The downfield shift of proton *d* and the amide protons α of the macrocycle in the rotaxane are indicative of interactions with a hydrogen bond acceptor on the axle. In the ¹H-¹H ROESY NMR spectra of both rotaxanes there are through-space correlations between resonances arising from protons in the two interlocked components (see ESI, pp. S43 & S44†). For rotaxane **21**, the multiple observed intercomponent correlations strongly support the primary location of the macrocyclic ring component as being in the vicinity of the axle isophthalamide in CDCl₃ (see ESI, p. S45†). Meanwhile for rotaxane **23** there is evidence the macrocyclic ring preferentially resides over the isophthalamide amide N–



Open Access Article. Published on 07 December 2022. Downloaded on 6/12/2025 10:07:05 AM.

H γ due to the observable correlations being between stopper proton 22 and ring protons *i* and *l* (see ESI, p. S45[†]).

Conclusions

Using precursors prepared from isophthalic acid derivative 1, we have demonstrated the hydrogen bond templated synthesis of both [2]catenanes and [2]rotaxanes. Despite using only stoichiometric (or near stoichiometric) quantities of precursors, yields of up to 70% for the crucial covalent capture step are possible. This work demonstrates synthetic methodologies that have the potential to incorporate a range of functionality into either the ring of a catenane or the stoppers of a rotaxane in a modular approach in good yields. Investigations into deploying these methodologies, including preparing mechanically chiral analogues,²⁰ are ongoing in our laboratories.

Experimental

General information

All reagents and solvents were used as obtained from commercial suppliers, unless otherwise stated. Dry solvents, Et_3N and DIPEA were purchased dry and stored under an inert atmosphere. $Cu(CH_3CN)_4BF_4$ was stored in a desiccator over P_4O_{10} . Petrol refers to the fractions of petroleum that boil between 40 °C and 60 °C. Deionized water was used in all cases. All aqueous solutions are saturated unless otherwise stated.

Azido-functionalized isophthalic acid derivative 1,¹² alkyne benzoic acids 2^{14} and 3,¹⁵ alkyne ethylene glycol acid 4,¹⁶ macrocycle 14,^{9,10} alkyne 20^8 were all synthesized based on previously reported procedures.

Silica gel with a 60 Å particle size was used as the stationary phase for column chromatography. Analytical TLC was used to monitor the progress of column chromatography, with TLC plates examined under short wavelength (254 nm) UV light, or staining with potassium permanganate and phosphomolybdic acid solutions as appropriate. Preparatory TLC was carried out on silica gel possessing a fluorescent indicator to allow for examination with short wavelength UV light.

IR spectra were recorded on an Agilent Technologies Cary 630 FTIR spectrometer. NMR spectra were recorded on a Bruker AVANCE III 400 or a Bruker Fourier 300 spectrometer at 298 K (unless otherwise stated). Mass spectra were recorded on a Shimadzu LCMS IT ToF instrument. Melting points were recorded on a Gallenkamp capillary melting point apparatus and are uncorrected.

Experimental procedures

Alkyne 5. To a solution of 2 (200 mg, 1.14 mmol) dissolved in dry CH₃CN (10 mL) was added DCC (281 mg, 1.36 mmol) and N-hydroxysuccinimide (157 mg, 1.36 mmol). The solution was stirred at room temperature under nitrogen for 16 hours. The solution was then filtered under gravity and the solvent removed in vacuo to afford an offwhite solid. The crude material was redissolved in dry CH₂Cl₂ (20 mL) and 4-(Boc-amino)benzylamine (268 mg, 1.14 mmol) was added to the solution followed by dry Et₃N (0.19 mL, 1.36 mmol). The solution was stirred at room temperature under nitrogen for 16 hours. The mixture was then washed with aq. 1 M HCl (2×20 mL), aq. NaHCO₃ (2 \times 20 mL) and water (1 \times 20 mL). The organic layer was dried (MgSO₄) and the solvent was removed in vacuo to afford an off-white solid. Purification by flash column chromatography (EtOAc: petrol 2:1) gave the title product (262 mg, 59%) as a colourless solid.



 $R_{\rm f}$ 0.60 (EtOAc : petrol 2 : 1).

m.p. 84-85 °C.

*ν*_{max}/cm⁻¹ (neat): 3315 (N−H), 3274 (N−H), 2928 (C−H), 1677 (C=O), 1632 (C=O).

δ_H (400 MHz, CDCl₃): 7.77 (2H, d, J = 9.0 Hz, H⁵), 7.34–7.20 (4H, m, H¹¹ & H¹²), 7.01 (2H, d, J = 9.0 Hz, H⁶), 6.28 (1H, bs, NH^δ), 4.83 (1H, bs, NH^γ), 4.75 (2H, d, J = 2.4 Hz, H³), 4.63 (2H, d, J = 5.7 Hz, H⁹), 4.31 (2H, d, J = 6.1 Hz, H¹⁴), 2.54 (1H, t, J = 2.4 Hz, H¹), 1.47 (9H, s, H¹⁷).

 $δ_{\rm C}$ (100 MHz, CDCl₃): 166.6 (C⁸) 166.5 (C¹⁵), 160.0 (C⁴), 138.4 (C⁷), 137.4 (C¹⁰), 128.7 (C⁶), 128.2 (C¹²), 127.8 (C¹¹), 127.4 (C¹³), 114.6 (C⁵), 79.5 (C²), 79.1 (C¹⁶), 76.0 (C¹), 55.6 (C³), 44.3 (C¹⁴), 43.7 (C⁹), 28.3 (C¹⁷).

m/z (ESI): 417.1785 ([M + Na]⁺ C₂₃H₂₆N₂NaO₄ requires 417.1769).

Deprotected amine–alkyne salt 8. To a solution of Boc-alkyne 5 (255 mg, 0.65 mmol) in CH_2Cl_2 (5 mL) cooled to 0 °C was added trifluoroacetic acid (1 mL, 3.7 mmol). The solution was warmed to room temperature and stirred for 2 hours, then the volatiles were removed *in vacuo*. The product was isolated as the TFA salt. Boc deprotection was confirmed by ¹H NMR analysis and the material was taken forward immediately without further purification.



δ_H (300 MHz, D₆-DMSO): δ 8.96 (1H, t, J = 6.0 Hz, NH^δ), 8.28–7.96 (3H, m, NH₃⁺), 7.90–7.82 (2H, m, H⁵), 7.42–7.30 (4H, m, H¹² & H¹¹), 7.08–7.02 (2H, m, H⁶), 4.87 (2H, d, J = 2.4 Hz, H³), 4.46 (2H, d, J = 6.0 Hz, H⁹), 4.00 (2H, q, J = 5.6 Hz, H¹⁴), 3.60 (1H, t, J = 2.4 Hz, H¹).

Alkyne–azide 11. To a solution of 1 (62 mg, 0.25 mmol) dissolved in dry CH₃CN (5 mL) was added DCC (61 mg, 0.32 mmol) and *N*-hydroxysuccinimide (31 mg, 0.29 mmol). The reaction was then stirred at room temperature under nitrogen for 16 hours. The resulting suspension was filtered under gravity and the solvent removed *in vacuo* to afford a white solid. The crude material was then redissolved in dry CH₂Cl₂ (20 mL) and **8** (102 mg, 0.25 mmol) and dry Et₃N (0.10 mL, 0.72 mmol) were added. The solution was then stirred at room temperature under nitrogen for 16 hours. The mixture was then washed with aq. 1 M HCl (2 × 20 mL), aq. NaHCO₃ (2 × 20 mL) and water (1 × 20 mL). The combined organic layers were dried (MgSO₄) and the solvent was removed *in vacuo* to afford a colourless solid. Purification by flash column chromatography (EtOAc : petrol 3 : 1–4 : 1) gave the *title product* (77 mg, 59%) as a colourless solid.



*R*_f 0.35 (EtOAc : petrol 4 : 1). m.p. 120–122 °C.

 ν_{max}/cm^{-1} (neat): 3281 (C=C-H), 2929 (C-H), 2102 (N=N=N), 1621 (C=O), 1541 (C-N).

$$\begin{split} & \pmb{\delta_{\rm H}} \ (\text{400 MHz}, \ 9:1 \ \text{CDCl}_3/\text{CD}_3\text{OD}): \ 8.17 \ (1\text{H}, \ \text{bs}, \ \text{H}^{10}), \ 7.94 \\ & (1\text{H}, \ \text{d}, J = 9.0 \ \text{Hz}, \ \text{H}^{6/8}) \ 7.90 \ (1\text{H}, \ \text{d}, J = 9.0 \ \text{Hz}, \ \text{H}^{6/8}), \ 7.74 \ (2\text{H}, \\ & \text{d}, J = 8.9 \ \text{Hz}, \ \text{H}^{21}), \ 7.44 \ (1\text{H}, \ \text{app. t}, \ \text{H}^7), \ 7.20\text{-}7.10 \ (4\text{H}, \ \text{m}, \ \text{H}^{14} \\ & \text{8} \ \text{H}^{15}), \ 6.94 \ (2\text{H}, \ \text{d}, J = 8.9 \ \text{Hz}, \ \text{H}^{20}), \ 4.69 \ (2\text{H}, \ \text{d}, J = 2.4 \ \text{Hz}, \\ & \text{H}^{23}), \ 4.48 \ (2\text{H}, \ \text{d}, J = 6.0 \ \text{Hz}, \ \text{H}^{17}), \ 4.43 \ (2\text{H}, \ \text{d}, J = 5.7 \ \text{Hz}, \ \text{H}^{12}), \\ & 3.39\text{-}3.35 \ (2\text{H}, \ \text{m}, \ \text{H}^3), \ 3.30 \ (2\text{H}, \ \text{t}, J = 9.0 \ \text{Hz}, \ \text{H}^1), \ 2.55 \ (1\text{H}, \ \text{t}, J = 2.4 \ \text{Hz}, \ \text{H}^{25}), \ 1.78 \ (2\text{H}, \ \text{quintet}, J = 6.5 \ \text{Hz}, \ \text{H}^2). \end{split}$$

 $δ_c$ (100 MHz, 9:1 CDCl₃/CD₃OD): 167.6 (C¹¹), 167.4 (C¹⁸), 167.2 (C⁴), 160.1 (C²²), 137.6 (C¹³), 137.1 (C^{5/9/19}) 137.0 (C¹⁶), 134.3 (C^{5/9/19}), 134.2 (C^{5/9/19}), 130.4 (C^{6/8}), 130.4 (C^{6/8}) 129.1 (C²¹), 129.0 (C⁷), 127.9 (C¹⁵), 127.0 (C¹⁴), 125.2 (C¹⁰), 114.6 (C²⁰), 77.8 (C²⁴), 76.0 (C²⁵), 55.7 (C²³), 49.1 (C¹), 43.7 (C¹⁷), 43.4 (C¹²), 37.4 (C³), 28.5 (C²).

m/z (ESI): 525.2245 ([M + H]⁺ C₂₉H₂₉N₆O₄ requires 525.2231).

Alkyne 6. To a solution of 3 (224 mg, 1.27 mmol) dissolved in dry CH₃CN (10 mL) was added DCC (314 mg, 1.35 mmol) and N-hydroxysuccinimide (175 mg, 1.27 mmol). The solution was stirred at room temperature under nitrogen for 16 hours. The solution was then filtered under gravity and the solvent removed in vacuo to afford an off-white solid. The crude material was redissolved in dry CH2Cl2 (20 mL) and 4-(Bocamino)benzylamine (300 mg, 1.27 mmol) was added to the solution followed by dry Et₃N (0.22 mL, 1.67 mmol). The solution was stirred at room temperature under nitrogen for 16 hours. The mixture was then washed with aq. 1 M HCl (2 \times 20 mL), aq. NaHCO₃ (2 \times 20 mL) and water (1 \times 20 mL). The combined organic layers were dried (MgSO₄) and the solvent was removed in vacuo to afford an off-white solid. Purification by flash column chromatography (EtOAc: Hexane 2:3-1:1) gave the title product (306 mg, 61%) as a colourless solid.

 $\begin{array}{c} & & & & \\ & & & & \\ & & & & \\ & & & & \\$

 $R_{\rm f}$ 0.87 (EtOAc : petrol 1 : 2).

т.р. 118–120 °С.

 $\nu_{\text{max}}/\text{cm}^{-1}$ (neat): 3310 (C=C-H), 3270 (N-H), 2930 (C-H), 1672 (C=O), 1635 (C=O).

 $δ_{\rm H}$ (400 MHz, CDCl₃): 7.48–7.45 (1H, m, H⁸), 7.38–7.29 (6H, m, H¹⁴, H¹³, H⁵ & H⁷), 7.16–7.11 (1H, m, H⁶), 6.38 (1H, bs, NH^δ), 4.88 (1H, bs, NH^γ), 4.75 (2H, d, J = 2.4 Hz, H³), 4.64 (2H, d, J = 5.8 Hz, H¹¹), 4.32 (2H, d, J = 5.8 Hz, H¹⁶), 2.55 (1H, t, J = 2.4 Hz, H¹), 1.48 (9H, s, H¹⁹).

 $δ_{\rm C}$ (100 MHz, CDCl₃): 166.9 (C¹⁰) 166.8 (C¹⁷), 157.8 (C⁴), 138.5 (C¹²), 137.1 (C¹⁵), 135.9 (C¹³), 129.7 (C⁶), 128.2 (C¹⁴), 127.8 (C¹⁹), 119.6 (C⁵), 118.5 (C⁷), 113.5 (C⁸), 79.1 (C¹⁸), 78.0 (C²), 75.9 (C¹), 55.9 (C³), 44.3 (C¹¹), 43.8 (C¹⁶), 28.4 (C¹⁹). m/z (ESI): 417.1785 ([M + Na]⁺ C₂₃H₂₆N₂NaO₄ requires 417.1771).

Deprotected amine-alkyne salt 9. To a solution of Boc-alkyne **6** (299 mg, 0.76 mmol) in CH_2Cl_2 (5 mL) cooled to 0 °C was added trifluoroacetic acid (1 mL, 3.7 mmol). The solution was warmed to room temperature and stirred for 2 hours, then volatiles were removed *in vacuo*. The product was isolated as the TFA salt. Boc deprotection was confirmed by ¹H NMR analysis and the material was taken forward immediately without further purification.



δ_H (400 MHz, D₆-DMSO): 9.06 (1H, t, J = 6.0 Hz, NH^δ), 8.28-7.96 (3H, m, NH₃⁺), 7.54-7.47 (2H, m, H⁵ & H⁷), 7.44-7.34 (5H, m, H⁸, H¹⁴ & H¹³), 7.19-7.14 (2H, m, H⁶), 4.86 (2H, d, J = 2.3 Hz, H³), 4.48 (2H, d, J = 6.0 Hz, H¹¹), 4.02 (2H, q, J = 5.8 Hz, H¹⁶), 3.60 (1H, t, J = 2.4 Hz, H¹).

Alkyne-azide 12. To a solution of 1 (182 mg, 0.73 mmol) dissolved in dry CH₃CN (10 mL) was added DCC (185 mg, 0.90 mmol) and N-hydroxysuccinimide (103 mg, 0.90 mmol). The reaction was stirred at room temperature under nitrogen for 16 hours. The resulting suspension was filtered under gravity and the solvent removed in vacuo to afford a white solid. The crude material was then redissolved in dry CH₂Cl₂ (20 mL) and 9 (220 mg, 0.75 mmol) and dry Et₃N (0.26 mL, 2.00 mmol) were added. The solution was stirred at room temperature under nitrogen for 16 hours. The mixture was then washed with aq. 1 M HCl (2×20 mL), aq. NaHCO₃ ($2 \times$ 20 mL) and water (1 \times 20 mL). The combined organic layers were dried (MgSO₄) and the solvent was removed in vacuo to afford a colourless solid. Purification by flash column chromatography (EtOAc: petrol 5:1) gave the *title product* (253 mg, 64%) as a colourless solid.



 $R_{f} = 0.23$ (EtOAc : petrol 3 : 1).

т.р. 122–124 °С.

*ν*_{max}/cm⁻¹ (neat): 3291 (N−H), 2926 (C−H), 2098 (N=N=N), 1634 (C=O), 1533 (C−N).

δ_H (400 MHz, CDCl₃): 8.36 (1H, s, H¹⁰), 8.08 (1H, d, J = 7.7 Hz, H^{6/8}), 7.98 (1H, d, J = 7.7 Hz, H^{6/8}), 7.53–7.48 (1H, m, H⁷), 7.49–7.47 (1H, m, H²⁰), 7.46–7.43 (1H, m, H^{24/22}), 7.35 (1H, t, J = 7.7 Hz, H²³), 7.21–7.13 (4H, m, H^{22/24} & NH^{β, γ, δ}), 7.10 (2H,

d, J = 8.9 Hz, H¹⁵), 7.01 (2H, d, J = 8.9 Hz, H¹⁴), 4.71 (2H, d, J = 2.4 Hz, H²⁵), 4.51 (2H, d, J = 5.8 Hz, H¹⁷), 4.36 (2H, d, J = 5.8 Hz, H¹²), 3.32 (2H, q, J = 6.6 Hz, H³), 3.25 (2H, t, J = 6.6 Hz, H¹), 2.53 (1H, t, J = 2.4 Hz, H²⁷), 1.71 (2H, quintet, J = 6.6 Hz, H²).

 $δ_{\rm C}$ (100 MHz, CDCl₃): 167.6 (C¹⁸), 166.8 (C¹¹), 166.3 (C⁴), 157.7 (C²¹), 137.4 (C¹⁶), 137.0 (C¹³), 135.2 (C^{5/9/19}), 134.1 (C^{5/9/19}), 134.0 (C^{5/9/19}), 130.7 (C^{6/8}), 130.6 (C^{6/8}), 129.7 (C²³), 129.0 (C⁷), 128.3 (C¹⁵), 127.3 (C¹⁴), 124.9 (C¹⁰), 119.8 (C²⁴), 118.6 (C²²), 113.7 (C²⁰), 78.0 (C²⁶), 75.9 (C²⁷), 55.9 (C²⁵), 49.1 (C¹), 43.9 (C¹⁷), 43.8 (C¹²), 37.6 (C³), 28.5 (C²).

m/z (ESI): 525.2245 ([M + H]⁺ C₂₉H₂₉N₆O₄ requires 525.2261).

Alkyne 7. To a solution of 4 (1.91 g, 12.1 mmol) dissolved in dry THF (30 mL) was added DCC (4.98 g, 24.2 mmol) and N-hydroxysuccinimide (2.78 g, 24.2 mmol). The reaction was stirred at room temperature under nitrogen for 16 hours. The resulting suspension was filtered under gravity and the solvent removed in vacuo to afford a white solid. The crude material was then redissolved in dry CH₂Cl₂ (30 mL) and 4-(Boc-amino) benzylamine (1.38 g, 5.85 mmol) and dry Et₃N (0.96 mL, 7.00 mmol) were added. The solution was stirred at room temperature under nitrogen for 16 hours. The mixture was then washed with aq. 1 M HCl (2 \times 20 mL), aq. NaHCO₃ (2 \times 20 mL) and water (1 \times 20 mL). The combined organic layers were dried (MgSO₄) and the solvent was removed in vacuo to afford a colourless solid. Purification by flash column chromatography (EtOAc: Hexane 1:1-2:1) gave the title product (1.67, 75%) as a clear oil.

and the crude material was taken forward immediately without further purification.



δ_H (400 MHz, D₆-DMSO): 8.35 (1H, t, J = 6.0 Hz, NH^δ), 8.28–7.96 (3H, m, NH₃⁺), 7.42–7.37 (2H, m, H¹¹), 7.34–7.29 (2H, m, H¹⁰), 4.32 (2H, d, J = 6.2 Hz, H⁸), 4.15 (2H, d, J = 2.3Hz, H³), 4.01 (2H, q, J = 5.8 Hz, H¹³), 3.95 (2H, s, H⁶), 3.65–3.59 (4H, m, H⁴ & H⁵), 3.44 (1H, t, J = 2.4 Hz, H¹).

Alkyne-azide 13. To a solution of 1 (250 mg, 1.00 mmol) dissolved in dry CH₃CN (10 mL) was added DCC (247 mg, 1.20 mmol) and N-hydroxysuccinimide (138 mg, 1.20 mmol). The reaction was stirred at room temperature under nitrogen for 16 hours. The resulting suspension was filtered under gravity and the solvent removed in vacuo to afford a colourless solid. The crude material was then redissolved in dry CH₂Cl₂ (20 mL) and 10 (390 mg, 1.00 mmol) and dry Et₃N (0.60 mL, 4.4 mmol) were added. The solution was stirred at room temperature under nitrogen for 16 hours. The solution was then washed with aq. 1 M HCl $(2 \times 20 \text{ mL})$, aq. NaHCO₃ $(2 \times 20 \text{ mL})$ and water (1×20 mL). The combined organic layers were dried (MgSO₄) and the solvent was removed in vacuo to afford a colourless solid. Purification by flash column chromatography (EtOAc: Hexane 6: 1-8:1) gave the *title product* (259 mg, 51%) as a colourless solid.



*R*_f 0.15 (EtOAc : hexane 1 : 1).

*ν*_{max}/cm⁻¹ (neat): 3330 (C≡C-H), 2976 (C-H), 1664 (C=O), 1515 (C-N), 1095 (C-O-C).

 $δ_{\rm H}$ (400 MHz, CDCl₃): 7.30–7.23 (5H, m, H¹⁰, H¹¹ & NH^δ), 4.88 (1H, bs, NH^γ), 4.47 (2H, d, *J* = 5.9 Hz, H⁸), 4.31 (2H, d, *J* = 5.7 Hz, H¹³), 4.06–4.03 (4H, m, H⁶ & H³), 3.72–3.65 (4H, H⁴ & H⁵), 2.41 (1H, t, *J* = 2.4 Hz, H¹), 1.46 (9H, s, H¹⁶).

 $δ_{\rm C}$ (100 MHz, CDCl₃): 169.6 (C⁷), 155.8 (C¹⁴), 138.2 (C⁹), 137.1 (C¹²), 128.2 (C¹⁰), 127.7 (C¹¹), 79.5 (C²), 79.1 (C¹⁵), 74.9 (C¹), 70.8 (C⁵), 70.4 (C⁴), 68.6 (C⁶), 58.2 (C³), 44.3 (C¹³), 42.6 (C⁸), 28.3 (C¹⁶).

m/z (ESI): 399.1890 ([M + Na]⁺ C₂₀H₂₈N₂NaO₅ requires 399.1890).

Deprotected amine–alkyne salt 10. To a solution of Bocalkyne 7 (386 mg, 1.03 mmol) in CH_2Cl_2 (5 mL) cooled to 0 °C was added trifluoroacetic acid (1 mL, 3.7 mmol). The solution was warmed to room temperature and stirred for 2 hours, then the volatiles removed *in vacuo*. The product was isolated as the TFA salt. Boc deprotection was confirmed by ¹H NMR analysis



 R_{f} 0.13 (EtOAc : Hexane 4 : 1).

m.p. 78–80 °C.

 $\nu_{\text{max}}/\text{cm}^{-1}$ (neat): 3291 (C=C-H), 3060 (N-H), 2916 (C-H), 2102 (N=N=N), 1649 (C=O), 1530 (C-N), 1097 (C-O-C).

δ_H (400 MHz, CDCl₃): 8.22 (1H, s, H¹⁰), 8.00–7.92 (2H, m, H⁶ & H⁸), 7.53 (1H, app. t, H⁷), 7.34 (1H, bs, NH^δ), 7.31–7.22 (4H, m, H¹⁴ & H¹⁵), 6.75–6.66 (2H, m, NH^{β,γ}), 4.61 (2H, d, J = 5.6 Hz, H¹²), 4.44 (2H, d, J = 5.9 Hz, H¹⁷), 4.08 (2H, d, J = 2.4 Hz, H²²), 4.03 (2H, s, H¹⁹), 3.73–3.66 (4H, m, H²¹ & H²⁰), 3.52 (2H, q, J = 6.6 Hz, H³), 3.42 (2H, t, J = 6.5 Hz, H¹), 2.42 (1H, t, J = 2.4 Hz, H²⁴), 1.88 (2H, quintet, J = 6.6 Hz, H²).

 $δ_{\rm C}$ (100 MHz, CDCl₃): 169.9 (C¹⁸), 166.6 (C⁴), 166.3 (C¹¹), 137.6 (C¹⁶), 137.0 (C¹³), 134.6 (C^{5/9}), 134.5 (C^{5/9}), 130.1 (C^{6/8}), 130.1 (C^{6/8}), 129.0 (C⁷), 128.2 (C¹⁴), 128.1 (C¹⁵), 125.2 (C¹⁰), 79.1 (C²³), 75.0 (C²⁴), 70.5 (C¹⁹), 70.0 (C²⁰), 68.7 (C²¹), 58.3 (C²²), 49.4 (C¹), 43.9 (C¹²), 42.5 (C¹⁷), 37.8 (C³), 28.7 (C²).

m/z (ESI): 529.2170 ([M + Na]⁺ C₂₆H₃₀N₆O₅ requires 529.2175).

Catenane 15. Macrocycle 14 (60 mg, 0.13 mmol) and alkyneazide 11 (68 mg, 0.13 mmol) were dissolved in dry CH_2Cl_2 (20 mL) under an argon atmosphere. To the solution, [Cu (CH₃CN)₄BF₄] (9.4 mg, 0.03 mmol), TBTA (16 mg, 0.03 mmol) and dry DIPEA (0.022 mL, 0.13 mmol) were added. The reaction was stirred at room temperature for 16 hours. The reaction mixture was diluted with further CH₂Cl₂ (10 mL) and then the solution was washed with 0.02 M EDTA in aq. 1 M NH₃ (2 \times 15 mL) and brine (1 \times 15 mL). The organic layer was dried $(MgSO_4)$ and solvent removed *in vacuo* to afford a vellow solid. Purification by flash column chromatography (CH₂Cl₂:CH₃OH 98:2) gave the *title product* (76 mg, 59%) as a colourless solid.



 $R_{\rm f}$ 0.29 (CH₂Cl₂:CH₃OH 98:2).

m.p. 165–167 °C.

 $\nu_{\text{max}}/\text{cm}^{-1}$ (neat): 3345 (N-H), 2868 (C-H), 1630 (C=O), 1578 (C-N), 1533 (C-N), 1416 (Ar-C).

 $\delta_{\rm H}$ (400 MHz, 50:50 CDCl₃/CD₃OD): 8.70 (1H, s, H^d), 8.21–8.16 (2H, m, H^b), 8.11 (1H, d, J = 7.7 Hz, H^{6/8}), 8.02 (1H, d, J = 7.8 Hz, $H^{6/8}$), 7.82–7.72 (3H, m, H^{20} & H^{25}), 7.68–7.59 (2H, m, H^7 & H^a), 7.52 (1H, s, H^{10}), 7.40 (2H, d, J = 7.9 Hz, H^{14}), 7.07–6.97 (4H, m, H^{15} & H^{21}), 6.94 (4H, d, J = 7.9 Hz, H^{h}), 6.56 (4H, d, J = 7.9 Hz, Hⁱ), 5.42 (2H, s, H²³), 4.58 (2H, dd, J = 14.2, 5.1 Hz, H^f), 4.47-4.36 (4H, m, H¹² & H^{f'}), 4.25 (2H, bs, H¹), 3.97-3.79 (6H, m, H^k, H^{k'} & H¹⁷), 2.98 (2H, bs, H³), 2.93-2.75 (4H, m, H^l), 2.64-2.48 (4H, m, H^m), 1.84 (2H, bs, H^2).

 $\delta_{\rm C}$ (100 MHz, 50:50 CDCl₃/CD₃OD): 167.4 (C^e), 167.0 (C⁴), 166.4 (C¹⁸), 166.3 (C¹¹), 159.8 (C²²), 143.4 (C²⁴), 139.3 (C¹³), 139.2 (C¹⁶), 135.2 (C^j), 133.9 (C^c), 133.9 (C^{5/9}), 133.8 (C^{5/9}), 133.5 (C^{6/8}), 131.8 (C^{6/8}), 131.6 (C^b), 131.5 (C^g), 129.6 (Cⁱ), 129.3 (C¹⁵), 129.3 (C¹⁴), 128.9 (C^a), 128.9 (C⁷), 128.7 (C²⁰), 128.1 (C^h), 126.3 (C^{19}), 124.6 (C^{d}) 124.6 (C^{10}), 123.4 (C^{25}) 115.3 (C^{21}), 73.4 (C^{k}) , 69.9 (C^{m}) , 68.4 (C^{l}) , 60.5 (C^{23}) , 47.9 (C^{1}) , 44.3 (C^{f}) , 43.4 $(C^{17}), 43.2 (C^{12}), 37.4 (C^3), 29.5 (C^2).$

m/z (ESI): 999.4400 ([M + H]⁺ C₅₇H₅₉N₈O₉ requires 999.4426).

Catenane 16. Macrocycle 14 (60 mg, 0.13 mmol) and alkyneazide 12 (68 mg, 0.13 mmol) were dissolved in dry CH₂Cl₂ (20 mL) and placed under an argon atmosphere. To the solution, [Cu(CH₃CN)₄BF₄] (9.4 mg, 0.03 mmol), TBTA (16 mg, 0.03 mmol) and dry DIPEA (0.022 mL, 0.13 mmol) were added. The reaction was stirred at room temperature for 16 hours. The reaction mixture was diluted with further CH_2Cl_2 (10 mL)

and then the solution was washed with 0.02 M EDTA in aq. 1 M NH₃ (2 \times 15 mL) and brine (1 \times 15 mL). The combined organic layers were dried (MgSO₄) and solvent removed in vacuo to afford a yellow solid. Purification by flash column chromatography (CH₂Cl₂:CH₃OH 98:2) gave the *title product* (79 mg, 60%) as a colourless solid.



*R*_f 0.29 (CH₂Cl₂:CH₃OH 98:2).

m.p. 168-170 °C.

 $\nu_{\text{max}}/\text{cm}^{-1}$ (neat): 3341(N-H), 2868 (C-H), 1630 (C=O), 1578 (C-N), 1533 (C-N), 1416 (Ar-C).

 $\delta_{\rm H}$ (400 MHz, 50:50 CDCl₃/CD₃OD): 8.71 (1H, s, H^d), 8.19 $(2H, dd, J = 7.8 Hz, H^{b}), 8.12 (1H, d, J = 7.7 Hz, H^{6/8}), 8.05 (1H, d, J = 7.7 Hz, H^{10}), 8.05 (1H, d, J = 7.8 Hz, H^{10}), 8.05 (1H$ d, J = 7.8 Hz, $H^{6/8}$), 7.80 (1H, s, H^{27}), 7.69–7.63 (2H, m, H^7 & H^{a}), 7.62 (1H, s, H^{10}), 7.51 (1H, d, J = 7.5 Hz, H^{24}), 7.43–7.35 $(3H, m, H^{23} \& H^{14}), 7.32 (1H, s, H^{20}), 7.22 (1H, dd, J = 7.2, 2.4)$ Hz, H^{22}), 7.04 (2H, d, J = 7.7 Hz, H^{15}), 6.94 (4H, d, J = 7.9 Hz, H^h), 6.56 (4H, d, J = 7.9 Hz, Hⁱ), 5.34 (2H, s, H²⁵), 4.59 (2H, d, J = 14.2 Hz, H^f), 4.47–4.35 (4H, m, H¹² & H^f), 4.26 (2H, bs, H¹), 3.98-3.90 (4H, m, H^k & H^{k'}), 3.88 (2H, bs, H¹⁷), 3.07-2.92 (8H, m, H¹, H^{1'}, H^m & H³), 2.78–2.71 (2H, m, H^{m'}), 1.96 (2H, bs, H²).

 $\delta_{\rm C}$ (100 MHz, 50:50 CDCl₃/CD₃OD): 167.8 (C^e), 166.5 (C¹⁸), 166.4 (C¹¹), 165.8 (C⁴), 157.24 (C²¹), 143.4 (C²⁶), 144.3 (C^{5/9}), 138.4 (C^{5/9}), 137.1 (C^j), 136.7 (C^{16/19/c}), 136.5 (C¹³), 135.0 (C^g), 134.6 ($C^{16/19/c}$), 131.6 ($C^{6/8}$), 131.6 ($C^{6/8}$), 131.6 (C^{b}), 130.0 (C^{23}), 129.8 (Cⁱ) 129.5 (C¹⁵) 129.3 (C¹⁴) 129.1 (C^a), 129.0 (C⁷), 128.4 (C^{h}) , 124.6 (C^{10}) , 124.6 (C^{d}) , 123.1 (C^{27}) , 119.3 (C^{24}) , 117.3 $(C^{22}), 114.7 (C^{20}), 72.9 (C^{k}), 69.9 (C^{l}), 68.5 (C^{m}), 60.9 (C^{25}), 52.8$ $(C^{1}), 43.6 (C^{f}), 42.8 (C^{12}), 42.7 (C^{17}), 36.3 (C^{1}), 29.2 (C^{2}).$

m/z (ESI): 999.4400 ([M + H]⁺ C₅₇H₅₉N₈O₉ requires 999.4427).

Catenane 17. Macrocycle 14 (60 mg, 0.13 mmol) and alkyneazide 13 (63 mg, 0.13 mmol) were dissolved in dry CH₂Cl₂ (20 mL) and placed under an argon atmosphere. To the solution, [Cu(CH₃CN)₄BF₄] (9.4 mg, 0.03 mmol), TBTA (16 mg, 0.03 mmol) and dry DIPEA (0.022 mL, 0.13 mmol) were added. The reaction was stirred at room temperature for 16 hours. The reaction mixture was diluted with further CH₂Cl₂ (10 mL) and then the solution was washed with 0.02 M EDTA in aq. 1 M NH₃ (2 \times 15 mL) and brine (1 \times 15 mL). The combined organic layers were dried (MgSO₄) and solvent removed in

View Article Online

vacuo to afford a yellow solid. Purification by flash column chromatography ($CH_2Cl_2:CH_3OH 96:4-94:6$) gave the *title* product (89 mg, 70%) as a colourless solid.



 $R_{f} 0.32 (CH_{2}Cl_{2}: CH_{3}OH 96: 4).$

m.p. 118–121 °C.

 ν_{max} /cm⁻¹ (neat): 3215 (C-H), 2860 (C-H), 1638 (C=O), 1522 (C-N), 1017 (C-O-C).

 $\delta_{\rm H}$ (400 MHz, 50:50 CDCl₃/CD₃OD): 8.70 (1H, s, H^d), 8.18 $(2H, dd, J = 7.8, 1.8 Hz, H^{b}), 8.13 (1H, d, J = 7.7 Hz, H^{6/8}), 8.05$ (1H, d, J = 7.8 Hz, H^{6/8}), 7.80 (1H, s, H²⁴), 7.70–7.63 (3H, m, H⁷ & H^{a}), 7.62 (1H, s, H^{10}), 7.26 (2H, d, J = 8.1 Hz, H^{14}), 7.03 (2H, d, J = 8.1 Hz, H¹⁵), 6.96 (4H, d, J = 7.9 Hz, H^h), 6.63 (4H, d, J = 7.9 Hz, Hⁱ), 4.63-4.53 (4H, m, H²² & H^f), 4.42-4.36 (4H, m, H^f & H¹²), 4.32 (2H, bs, H¹), 4.11-4.02 (4H, m, H^k), 3.98 (2H, m, H¹⁹), 3.90 (2H, bs, H¹⁷), 3.71 (4H, bs, H²⁰ & H²¹), 3.15-3.00 (8H, dt, m, H¹, H³, H^m & H¹), 2.90–2.85 (2H, m, H^{m'}), 2.08 (2H, bs, H^2).

 $\delta_{\rm C}$ (100 MHz, 50:50 CDCl₃/CD₃OD): 170.0 (C¹⁸), 167.4 (C^e), 167.0 (C⁴), 166.4 (C¹¹), 144.3 (C²³), 138.7 (C¹⁶),138.4 (C^{5/9/c}), 138.3 (C^{5/9/c}), 137.1 (C^{5/9/c}), 136.9 (C¹³), 135.4 (C^j), 131.7 (C^b) 131.6 (C^{6/8}), 131.4 (C^{6/8}), 131.2 (C^g), 129.5 (Cⁱ) 129.3 (C¹⁵), 129.1 (C^a), 128.7 (C⁷), 128.5 (C¹⁴), 128.2 (C^h), 124.8 (C¹⁰), 124.6 (C^d), 123.6 (C^{24}), 73.5 (C^{k}), 70.4 (C^{19}), 70.3 (C^{m}), 70.2 (C^{21}), 69.7 (C²⁰) 68.7 (C¹), 64.2 (C²²), 47.5 (C¹), 44.2 (C^f), 43.3 (C¹⁷), 42.1 $(C^{12}), 36.6 (C^3), 29.5 (C^2).$

m/z (ESI): 1003.4325 ([M + H]⁺ C₅₄H₆₁N₈O₁₀ requires 1003.4372).

Azide 18. To a solution of 1 (100 mg, 0.40 mmol) in dry CH₃CN (15 mL) under an argon atmosphere was added DCC (99 mg, 0.48 mmol) and N-hydroxysuccinimide (55 mg, 0.48 mmol). The reaction mixture was stirred at room temperature for 16 hours. The resulting suspension was filtered under gravity and the solvent removed in vacuo. The resulting crude material was re-dissolved in dry CH₂Cl₂ (20 mL). 3,5-Bis(trifluoromethyl)benzylamine (117 mg, 0.48 mmol) and Et₃N (0.11 mL, 0.80 mmol) were then added, and the reaction mixture was stirred at room temperature overnight under an argon atmosphere. The reaction mixture was then washed with aq. 1 M HCl (2 \times 15 mL) and aq. NaHCO₃ (2 \times 15 mL). The organic layer was dried (MgSO₄) and solvent removed *in vacuo*. The crude material was purified by silica gel column chromatography (EtOAc/hexane 1:1) to afford the title product as a waxy clear oil (129 mg, 68%).



 R_{f} 0.15 (EtOAc : Hexane 3 : 7).

 $\nu_{\rm max}/{\rm cm}^{-1}$ (neat): 3330 (CH₂), 2933 (ArCH), 2096 (N=NN), 1638 (C=O), 1531 (C-N).

 $\delta_{\rm H}$ (400 MHz, CDCl₃): 8.23 (1H, s, H¹⁰), 8.01-7.96 (1H, m, H⁸), 7.93-7.88 (1H, m, H⁶), 7.80 (3H, s, H¹⁴ & H¹⁶), 7.54 (1H, app. t, H^7), 6.95 (1H, bs, NH^{γ}), 6.54 (1H, bs, NH^{β}), 4.77 (2H, d, J = 6.1 Hz, H¹²), 3.56 (2H, t, J = 6.6 Hz, H³), 3.45 (2H, t, J = 6.5Hz, H^{1}), 1.91 (2H, quintet, I = 6.6 Hz, H^{2}).

 $\delta_{\rm C}$ (100 MHz, CDCl₃): 166.6 (C¹¹), 166.5 (C⁴), 140.7 (C¹³), 134.8 (C⁹), 134.0 (C⁵), 131.9 (q, ${}^{2}J$ = 34 Hz, C¹⁵), 130.2 (C⁸), 130.0 (C⁶), 129.2 (C⁷), 128.0 (C¹⁴), 125.5 (C¹⁰), 124.6 (C¹⁶), 121.9 $(q, {}^{1}J = 271 \text{ Hz}, C^{17}), 49.5 (C^{1}), 43.4 (C^{12}), 38.0 (C^{3}), 28.7 (C^{2}).$

 $\delta_{\rm F}$ (377 MHz, CDCl₃): -62.8

m/z (ESI): 474.1359 ([M + H]⁺ C₂₀H₁₈N₅O₂F₆ requires 474.1346).

Azide 19. To a solution of 1 (95 mg, 0.38 mmol) in dry CH₃CN (15 mL) under an argon atmosphere was added DCC (95 mg, 0.46 mmol) and N-hydroxysuccinimide (52 mg, 0.46 mmol). The reaction mixture was stirred at room temperature for 16 hours maintaining the argon atmosphere. The resulting suspension was filtered under gravity and excess CH3CN removed in vacuo. The resulting crude material was redissolved in CH₂Cl₂ (20 mL). 3.5-Bis(trifluoromethyl)aniline (110 mg, 0.42 mmol) and Et₃N (0.11 mL, 0.80 mmol) were then added, and the reaction mixture was stirred at room temperature overnight. The reaction mixture was then washed with 1 M HCl (2 \times 15 mL) and NaHCO₃ (2 \times 15 mL). The organic layer was dried (MgSO₄) and solvent removed in vacuo. The crude material was purified by silica gel column chromatography (EtOAc/hexane 1:1) to afford the title product as a brown oil (89 mg, 51%).



R_{f} 0.17 (EtOAc : hexane 3 : 7).

 $\nu_{\rm max}/{\rm cm}^{-1}$ (neat): 3330 (N-H), 2935 (ArCH), 2096 (N=NN), 1638 (C=O), 1531 (C-N).

 $δ_{\rm H}$ (400 MHz, D₆-DMSO): 10.99 (1H, bs, NH^γ), 8.72 (1H, bs, NH^β), 8.55 (2H, s, H¹³), 8.49 (1H, s, H¹⁵), 8.15 (1H, d, J = 7.1 Hz, H⁶), 8.10 (1H, d, J = 7.2 Hz, H⁸), 7.84 (1H, s, H¹⁰), 7.68 (1H, t, J = 7.8 Hz, H⁷), 3.45 (2H, t, J = 6.5 Hz, H³), 3.38 (2H, t, J = 6.5 Hz, H¹), 1.82 (2H, quintet, J = 6.6 Hz, H²).

 $δ_{\rm C}$ (100 MHz, D₆-DMSO): 166.3 (C¹¹), 166.1 (C⁴), 141.4 (C¹²), 135.4 (C⁹), 134.5 (C⁵), 131.3 (q, ²J = 34 Hz, C¹⁴), 131.1 (C⁷), 130.9 (C¹⁵), 129.1 (C⁶), 127.2 (C⁸), 122.3 (C¹³), 120.3 (q, ¹J = 271 Hz, C¹⁶), 117.3 (C¹⁰), 48.9 (C¹), 37.2 (C³), 28.8 (C²).

 $\delta_{\rm F}$ (377 MHz, D₆-DMSO): -61.6.

m/z (ESI): 458.1057 ([M - H]⁻ C₁₉H₁₆F₆N₅O₂ requires 458.1065).

Rotaxane 21. Macrocycle 14 (20 mg, 0.042 mmol) and azide 18 (21 mg, 0.046 mmol) were dissolved in dry CH₂Cl₂ (1 mL) under an argon atmosphere. Then alkyne 20 (12 mg, 0.046 mmol), [Cu(CH₃CN)₄BF₄] (1.4 mg, 0.004 mmol), TBTA (2.4 mg, 0.004 mmol) and dry DIPEA (9 µL, 6.6 mg, 0.051 mmol) were added. The reaction was stirred at RT for 18 hours maintaining the argon atmosphere. Then, the reaction was diluted to 10 mL, washed with 0.02 M EDTA in aq. 1 M NH₃ (2 \times 10 mL) and brine (1 \times 10 mL). The organic layer was dried (MgSO₄), filtered and solvent removed in vacuo. The crude material was purified by preparative TLC (repeated running in 96:2:3 CH₂Cl₂/CH₃OH/CH₃COCH₃) which allowed for isolation of the product with contaminated macrocycle 14. Pure rotaxane was isolated after running another preparative TLC (run twice in 95:2:3 CH₂Cl₂/CH₃OH/CH₃COCH₃) to give the title product as a colourless glassy film (14 mg, 29%).



 $R_{f} 0.59 (CH_{2}Cl_{2}: CH_{3}OH 96: 4).$

*ν*_{max}/cm⁻¹ (neat): 3332 (N–H), 2929 (C–H), 2862 (C–H),1638 (C=O), 1528 (C–N), 1276 (C–O).

$$\begin{split} & \delta_{\rm H} \, (400 \; {\rm MHz}, {\rm CDCl}_3) \!\!: 8.66 \; (1{\rm H}, {\rm s}, {\rm H}^d), 8.31{-}8.21 \; (3{\rm H}, {\rm m}, {\rm H}^b \\ & {\rm H}^{15/17}), \; 8.07 \; (1{\rm H}, \; {\rm bs}, \; {\rm H}^{15/17}), \; 7.84 \; (1{\rm H}, {\rm s}, \; {\rm H}^{25}), \; 7.78 \; (2{\rm H}, \; {\rm bs}, \\ & {\rm H}^{23}), \; 7.73 \; (2{\rm H}, \; {\rm m}, \; {\rm H}^{16} \; {\rm \&} \; {\rm H}^{19}), \; 7.62 \; (1{\rm H}, {\rm t}, J = 7.8 \; {\rm Hz}, \; {\rm H}^a), \; 7.56 \\ (2{\rm H}, \; {\rm bs}, \; {\rm NH}^\alpha), \; 7.32{-}7.16 \; (11{\rm H}, \; {\rm m}, \; {\rm H}^1, \; {\rm H}^2 \; {\rm H}^3 \; {\rm \&} \; {\rm H}^9), \; 6.84 \; (4{\rm H}, \\ {\rm d}, J = 7.9 \; {\rm Hz}, \; {\rm H}^h), \; 6.53 \; (4{\rm H}, \; {\rm d}, J = 8.0 \; {\rm Hz}, \; {\rm H}^i), \; 4.64 \; (2{\rm H}, {\rm s}, \; {\rm H}^7), \\ 4.59{-}4.37 \; (4{\rm H}, \; {\rm m}, \; {\rm H}^f), \; 4.32 \; (1{\rm H}, \; {\rm t}, J = 7.7 \; {\rm Hz}, \; {\rm H}^5), \; 4.14{-}4.02 \\ (8{\rm H}, \; {\rm m}, \; {\rm H}^6, \; {\rm H}^k \; {\rm \&} {\rm H}^{10}), \; 3.97 \; (2{\rm H}, \; {\rm bs}, \; {\rm H}^{21}), \; 3.70{-}3.55 \; (2{\rm H}, \; {\rm m}, \; {\rm H}^1), \\ 3.44{-}3.30 \; (4{\rm H}, \; {\rm m}, \; {\rm H}^m \; {\rm \&} \; {\rm H}^{1'}), \; 3.28{-}3.20 \; (2{\rm H}, \; {\rm m}, \; {\rm H}^m), \; 2.78 \; (2{\rm H}, \\ {\rm bs}, \; {\rm H}^{12}), \; 1.78 \; (2{\rm H}, \; {\rm bs}, \; {\rm H}^{11}). \end{split}$$

 $δ_{\rm C}$ (100 MHz, CDCl₃): 166.9 (C¹³), 166.6 (C²⁰), 166.4 (C^e), 145.2 (C⁸), 141.9 (C⁴), 136.4 (C^g), 134.8 (C^j), 133.8 (C^{14/18/22}), 133.7 (C^{14/18/22}), 133.1 (C^{14/18/22}), 132.1 (C²⁴), 132.0 (C^b), 131.8 $\begin{array}{l} (C^{15/17}), \ 131.6 \ (C^{15/17}), \ 129.9 \ (C^{i}), \ 129.5 \ (C^{a}), \ 129.2 \ (C^{23}), \ 128.8 \\ (C^{16}), \ 128.4 \ (C^{3}), \ 128.3 \ (C^{h}), \ 128.2 \ (C^{2}), \ 126.5 \ (C^{1}), \ 124.5 \ (C^{25}), \\ 123.8 \ (C^{d}), \ 122.6 \ (C^{9}), \ 121.7 \ (C^{19}), \ 73.9 \ (C^{k}), \ 73.7 \ (C^{6}), \ 70.7 \\ (C^{m}), \ 68.8 \ (C^{l}), \ 64.5 \ (C^{7}), \ 50.9 \ (C^{5}), \ 47.6 \ (C^{10}), \ 44.5 \ (C^{f}), \ 43.3 \\ (C^{21}), \ 36.8 \ (C^{12}), \ 29.6 \ (C^{11}). \end{array}$

 $\delta_{\rm F}$ (377 MHz, CDCl₃): -62.6.

m/z (ESI): 1184.4715 ([M + H]⁺ C₆₅H₆₄F₆N₇O₈ requires 1184.4752).

Rotaxane 23. Macrocycle 14 (20 mg, 0.042 mmol) and azide 19 (21 mg, 0.046 mmol) were dissolved in dry CH₂Cl₂ (1 mL) under an argon atmosphere. Then alkyne 20 (12 mg, 0.046 mmol), [Cu(CH₃CN)₄BF₄] (1.4 mg, 0.004 mmol), TBTA (2.4 mg, 0.004 mmol) and dry DIPEA (9 µL, 6.6 mg, 0.051 mmol) were added. The reaction was stirred at RT for 18 hours maintaining the argon atmosphere. Then, the reaction was diluted to 10 mL, washed with 0.02 M EDTA in aq. 1 M NH₃ solution $(2 \times 10 \text{ mL})$ and brine $(1 \times 10 \text{ mL})$. The organic layer was dried (MgSO₄), filtered and solvent removed in vacuo. The crude material was purified by preparative TLC (repeated running in 98:2 CH₂Cl₂/CH₃OH) which allowed for isolation of the product with contaminated macrocycle 14. Pure rotaxane was isolated after running another preparative TLC (repeated running in 96:2:2 CH₂Cl₂/CH₃OH/CH₃COCH₃) to give the *title product* as a colourless glassy film (20 mg, 40%).



 $R_{f} 0.64 (CH_{2}Cl_{2}:CH_{3}OH 96:4).$

 ν_{max} /cm⁻¹ (neat): 3319 (N-H), 2918 (C-H), 1645 (C=O), 1528 (C-N), 1276 (C-O).

δ_H (400 MHz, CDCl₃): 8.55 (1H, s, H^d), 8.32–8.22 (3H, m, H^b & H^{15/17}), 8.06–7.87 (3H, m, H^{15/17} & H¹⁹), 7.84 (1H, bs, H²²), 7.72–7.62 (2H, m, H¹⁶ & H^a), 7.58 (1H, s, H²⁴), 7.43–7.17 (13H, m, H¹, H², H³, H⁹ & NH^α), 6.78 (4H, d, J = 7.9 Hz, H^h), 6.54 (4H, d, J = 8.0 Hz, Hⁱ), 4.67 (2H, s, H⁷), 4.52–4.36 (4H, m, H^f), 4.34–4.20 (3H, m, H⁵ & H¹⁰), 4.18–4.10 (4H, m, H^k), 4.09 (2H, d, 7.3 Hz, H⁶), 3.79–3.72 (2H, m, H^m), 3.68–3.60 (2H, m, H^{m'}), 3.55–3.44 (4H, m, H^l), 3.33 (2H, bs, H¹²), 2.14 (2H, bs, H¹¹).

 $δ_{\rm C}$ (100 MHz, CDCl₃): 166.3 (C^e), 166.2 (C²⁰), 165.7 (C¹³), 145.3 (C⁸), 141.9 (C^{18/14}), 139.4 (C^{21/c}), 137.4 (C^j), 134.8 (C^g), 133.7 (C^{14/18}), 133.7 (C^{21/c}), 132.8 (C^{15/17}), 132.0 (C^{15/17}), 131.9 (C^b), 129.4 (C^a), 129.4 (C⁴), 129.3 (Cⁱ), 129.0 (C¹⁶), 128.5 (C²²), 128.4 (C²), 128.2 (C³), 128.2 (C^h), 126.5 (C¹), 124.8 (C²²), 123.1 (C^d), 122.7 (C⁹), 121.6 (C¹⁹), 117.0 (C²⁴), 73.9 (C^k), 73.7 (C⁶), 70.6 (C^m), 68.8 (C¹), 64.6 (C⁷), 50.9 (C⁵), 47.8 (C¹⁰), 44.4 (C^f), 37.1 (C¹²), 29.7 (C¹¹).

 $\delta_{\rm F}$ (377 MHz, CDCl₃): -63.0.

m/z (ESI): 1170.4559 ([M + H]⁺ C₆₄H₆₂F₆N₇O₈ requires 1170.4595).

Author contributions

NHE proposed the study. NHE conducted initial experiments (see ESI[†]). SRB conducted the synthesis, characterization and analysis of all materials in the main article with assistance from GRA. NHE supervised the work. SRB and NHE wrote the manuscript. All authors discussed and commented on the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

SRB acknowledges PhD funding from EPRSC DTP (EP/ T518037/1) and a Sydney Andrew Scholarship from the Society of Chemical Industry. NHE acknowledges the prize sponsored by Amigo Chem, that was awarded at the EPSRC Dial-a-Molecule and Directed Assembly Networks ECR "Supporting Synthesis and Self-Assembly" event held at the University of Liverpool in 2017, part of which funded the initial experiments reported in the ESI.† We thank Dr David Rochester (Lancaster University) for the recording of mass spectrometry data.

Underlying data for this paper are provided in the Experimental section and ESI.† Electronic copies of NMR spectra (including fid files) will be available upon publication from: https://doi.org/10.17635/lancaster/researchdata/554, https://doi.org/10.17635/lancaster/researchdata/569, https://doi.org/10.17635/lancaster/researchdata/570, https://doi.org/10.17635/lancaster/researchdata/574.

References

- 1 G. Gil-Ramírez, D. A. Leigh and A. J. Stephens, *Angew. Chem., Int. Ed.*, 2015, **54**, 6100–6150.
- 2 M. Xue, Y. Yang, X. Chi, X. Yan and F. Huang, *Chem. Rev.*, 2015, **115**, 7398–7501.
- 3 C. J. Bruns and J. F. Stoddart, *The Nature of the Mechanical Bond: From Molecules to Machines*, Wiley & Sons, 2017.
- 4 Review: (a) J. E. M. Lewis, P. D. Beer, S. J. Loeb and S. M. Goldup, *Chem. Soc. Rev.*, 2017, 46, 2577–2591 Seminal examples: (b) C.-O. Dietrich-Buchecker, J.-P. Sauvage and J.-P. Kintzinger, *Tetrahedron Lett.*, 1983, 34, 5095–5098; (c) V. Aucagne, K. D. Hänni, D. A. Leigh, P. J. Lusby and D. B. Walker, *J. Am. Chem. Soc.*, 2006, 128, 2186–2187 Recent examples: (d) A. W. Heard and S. M. Goldup, *Chem*,

2020, **6**, 994–1006; (e) R. C. Knighton and P. D. Beer, *Org. Chem. Front.*, 2021, **8**, 2468–2472.

- Review: (a) G. Barin, A. Coskin, M. M. G. Fouda and J. F. Stoddart, ChemPlusChem, 2012, 77, 159–185 Seminal examples: (b) P. R. Ashton, T. T. Goodnow, A. E. Kaifer, M. V. Reddington, A. M. Z. Slawin, N. Spencer, J. F. Stoddart, C. Vicent and D. J. Williams, Angew. Chem., Int. Ed. Engl., 1989, 28, 1396–1399; (c) D. G. Hamilton, J. K. M. Sanders, J. E. Davies, W. Clegg and S. J. Teat, Chem. Commun., 1997, 897–898 Recent examples: (d) T.-M. Gianga, E. Audibert, A. Trandafir, G. Kociok-Köhn and G. D. Pantoş, Chem. Sci., 2020, 11, 9685–9690; (e) Y. Jiao, L. Dorđević, H. Mao, R. M. Young, T. Jaynes, H. Chen, Y. Qiu, K. Cai, L. Zhang, X.-Y. Chen, Y. Feng, M. R. Waseielewski, S. I. Stupp and J. F. Stoddart, J. Am. Chem. Soc., 2021, 143, 8000–8100.
- 6 Review: (a) N. H. Evans, Eur. J. Org. Chem., 2019, 3320–3343
 Seminal examples: (b) C. A. Hunter, J. Am. Chem. Soc., 1992, 114, 5303–5311; (c) F. Vögtle, S. Meier and R. Hoss, Angew. Chem., Int. Ed. Engl., 1992, 31, 1619–1622; (d) P. R. Ashton, P. T. Glink, J. F. Stoddart, P. A. Tasker, A. J. P. White and D. J. Williams, Chem. Eur. J., 1996, 2, 729–726; (e) F. G. Gatti, D. A. Leigh, S. A. Nepogodiev, A. M. Z. Slawin, S. J. Teat and J. K. Y. Wong, J. Am. Chem. Soc., 2001, 123, 5983–5989 Recent examples: (f) S. D. P. Fielden, D. A. Leigh, C. T. McTernan, B. Pérez-Saavdra and I. J. Vitorica-Yrezabal, J. Am. Chem. Soc., 2018, 140, 6409–6052; (g) S. Amano, S. D. P. Fielden and D. A. Leigh, Nature, 2021, 594, 529–534.
- 7 Relevant reviews: (a) G. T. Spence and P. D. Beer, Acc. Chem. Res., 2013, 46, 571-586; (b) K. M. Bąk, K. Porfyrakis, J. J. Davis and P. D. Beer, Mater. Chem. Front., 2020, 4, 1052-1073 Seminal examples: (c) G. M. Hübner, J. Gläser, C. Seel and F. Vögtle, Angew. Chem., Int. Ed., 1999, 38, 383-386; (d) J. A. Wisner, P. D. Beer, M. G. B. Drew and M. R. Sambrook, J. Am. Chem. Soc., 2002, 124, 12469-12476; (e) N. L. Kilah, M. D. Wise, C. J. Serpell, A. L. Thompson, N. G. White, K. E. Christensen and P. D. Beer, J. Am. Chem. Soc., 2010, 132, 11893-11895; (f) S. Lee, C. H. Chen and A. H. Flood, Nat. Chem., 2013, 5, 704-710.
- 8 N. H. Evans, C. E. Gell and M. J. G. Peach, *Org. Biomol. Chem.*, 2016, **14**, 792–7981.
- 9 B. E. Fletcher, M. J. G. Peach and N. H. Evans, Org. Biomol. Chem., 2017, 15, 2797–2803.
- 10 C. N. Marrs and N. H. Evans, Org. Biomol. Chem., 2015, 13, 11021–11025.
- 11 Lewis has subsequently reported yields of up 51% for analogous [2]catenane formation by using bis-amines with longer glycol chains: J. E. M. Lewis, *Org. Biomol. Chem.*, 2019, 17, 2442–2447.
- 12 M. J. Langton, L. C. Duckworth and P. D. Beer, *Chem. Commun.*, 2013, **49**, 8608–8610.
- 13 In our hands, we obtained 1 in higher yield than previously reported. See ESI p. S3 for details.[†]
- 14 C. Deng, R. Fang, Y. Guan, J. Jiang, C. Lin and L. Wang, *Chem. Commun.*, 2012, **48**, 7973–7975.
- 15 (*a*) H. M. Branderhorst, R. Ruijtenbeek, R. M. J. Liskamp and R. J. Pieter, *ChemBioChem*, 2008, **9**, 1836–1844.

Paper

- K. Hayashi, Y. Miyaoka, Y. Ohishi, T. Uchinda, M. Iwamura,
 K. Nozaki and M. Inouye, *Chem. Eur. J.*, 2018, 24, 14613– 14616.
- 17 The corresponding author had previously prepared (but not reported) [2]catenane **15**, preparing precursor **11** by a different route. See ESI pp. S48–S67 for details.[†]
- 18 To date it has not been possible to isolate pure samples of the macrocyclic product arising from cyclization of **11–13**.
- 19 By analogous reactions (in the absence of macrocycle 14), non-interlocked axles 22 and 24 were also prepared (in 80% and 79% yield respectively) to aid ¹H NMR spectroscopic interpretation (see ESI pp. S3–S5†).
- 20 (a) N. H. Evans, Chem. Eur. J., 2018, 24, 3101-3112;
 (b) N. Pairault and J. Niemeyer, Synlett, 2018, 689-698;
 (c) E. M. G. Jamieson, F. Modicom and S. M. Goldup, Chem. Soc. Rev., 2018, 47, 5266-5311; (d) K. Nakazono and T. Takata, Symmetry, 2020, 12, 144.