ORGANIC CHEMISTRY

FRONTIERS

RESEARCH ARTICLE

Check for updates

Cite this: Org. Chem. Front., 2022, 9 342

Electrophilic N-trifluoromethylthiophthalimide as a fluorinated reagent in the synthesis of acyl fluorides*

HINESE

CHEMICAL

OCIFT

Chen Zhu, () * Serik Zhumagazy, () * Huifeng Yue * and Magnus Rueping *

Received 2nd November 2021, Accepted 22nd November 2021 DOI: 10.1039/d1qo01633d rsc.li/frontiers-organic

Herein we report the deoxygenated fluorination of readily available carboxylic acids. A series of acyl fluorides have been synthesized using shelf-stable N-trifluoromethylthiophthalimide as a fluorinated reagent for the first time. Scale-up reactions and sequential cross-couplings were performed successfully to demonstrate the practicability of this fluorination protocol.

Fluorine-containing molecules are one class of the most important organic compounds due to the unique character of the fluorine atom in modulating the chemical and biological properties such as metabolic stability, lipophilicity, and bioavailability.¹ Therefore, the development of methods to access such compounds is always a research focus for organic chemists. In this context, acvl fluorides represent intriguing targets that play an important role in organic synthesis. Due to the outstanding balance between stability and reactivity, acyl fluorides were widely employed as versatile synthons in different types of organic transformations (Scheme 1a).² Also, acyl fluorides could be used for the activation of silvl enol ethers or other silicon species in the presence of a Lewis base.³ Conventional methods to access acyl fluorides depend on the halogen exchange reaction of acyl chlorides with "F"" sources.⁴ Recent advances were mainly focused on the deoxyfluorination of readily available carboxylic acids with various fluorinated reagents (e.g., CF₃SO₂OCF₃,⁵ PPh₃/Selectfluor,⁶ Me₄NSCF₃,⁷ cyanuric fluoride,⁸ HF-Pyridine/DCC,⁹ sulphur tetrafluoride derivatives,10 and Carpino'salt TFFH11). Notably, Shibata and coworkers developed a general and practical method to prepare acyl fluorides, in which acids, aldehydes, and alcohols all could be transformed into acyl fluorides by using inexpensive reagents, TCCA and CsF.¹² In addition, acyl fluorides were skillfully accessed via selective C-C bond cleavage using DAST or its derivatives as fluorination reagents.¹³ Although these achievements provide good alternatives, most of them suffered from some drawbacks, mainly including high

342 | Org. Chem. Front., 2022, 9, 342-346

reaction temperature or the use of fluorinated reagents that are toxic, unstable, or expensive. Thus, the development of new approaches for the preparation of acyl fluorides is still highly desirable.

ROYAL SOCIETY

View Article Online

View Journal | View Issue

CHEMISTRY

Shelf-stable N-trifluoromethylthiophthalimide has been developed as an easily synthesized and efficient trifluoromethylthiolation reagent, and an array of valuable compounds were obtained employing this versatile SCF₃-reagent.¹⁴ In 2000, Munavalli and coworkers reported the reaction of SCF₃phthalimide with enamines, affording the α -trifluoromethylthiolated ketones.^{14a} Our group realized the enantioselective trifluoromethylthiolation of β -ketoester^{14b} and oxindoles,^{14c} Cu-catalyzed cross-coupling of boronic acids and alkynes,^{14d} as





(b) The reaction of carboxylic acids and N-trifluoromethylthiophthalimide under different conditions



Scheme 1 Acyl fluorides as synthons in organic synthesis and the reaction of N-trifluoromethylthiophthalimide and carboxylic acids.



KAUST Catalysis Center (KCC), King Abdullah University of Science and Technology (KAUST), Thuwal, 23955-6900, Saudi Arabia. E-mail: huifeng.yue@kaust.edu.sa, magnus.rueping@kaust.edu.sa

[†]Electronic supplementary information (ESI) available. See DOI: 10.1039/ d1qo01633d

[‡]These authors contributed equally to this work.

Organic Chemistry Frontiers

well as the metal-free ring-opening/trifluoromethylthiolation of cycloalkanols¹⁵ with this SCF₃-reagent.

More recently, Hu and coworkers developed a protocol on deoxygenated trifluoromethylthiolation of carboxylic acids in the presence of FeCl₃ and PPh₃, wherein SCF₃-phthalimide acted as a nucleophilic "SCF₃" source.^{14e} In contrast, we herein report that electrophilic *N*-trifluoromethylthiophthalimide acts as a fluorinated reagent in the reaction of carboxylic acids (Scheme 1b).

We started our investigation by evaluating the reaction of 4-(*tert*-butyl)benzoic acid **1a** and *N*-trifluoromethylthiophthalimide **2a** (Table 1). After a series of screening, the optimal reaction conditions were assigned as follows: 10 mol% TBAI as catalyst, 1 equiv. of Et₃N as reductant in 2 mL CH₃CN at 35 °C. The use of DIPEA as base gave lower yield, while the application of other bases including 2,6-lutidine, pyridine, and *n*BnN led to no formation of the desired product. Other solvents such as DMF, DMA, DMSO, THF, acetone were also suitable for this transformation, albeit delivering the acyl fluoride product in low to moderate yields. Using NaI or KI instead of TBAI decreased the yields dramatically. Reversing the ratio of **1a** and **2** from 1:1 to 1:2 also damaged the yield. Control experiments showed that both TBAI and Et₃N are essential for the high efficiency of this transformation.

With the optimized reaction conditions in hand, the generality of the fluorination reaction was first examined. As shown in Table 2, a variety of aryl carboxylic acids could undergo this efficient fluorination transformation smoothly, affording the corresponding products in good to excellent yields. In addition to alkyl (**3a** and **3b**), methoxyl (**3c**), methylthio (**3d**), and phe-

Table 1 Optimization of the reaction of carboxylic acid and
 N-trifluoromethylthiophthalimide to form acyl fluoride

| ^t Bu | O OH ₊ O N-SCF ₃ TBAI (10 mol% Et ₃ N (1 equiv.) CH ₃ CN, 35 °C, 10 | |
|-----------------|---|--------------------------------|
| 1a | 2 | 3a |
| Entry | Variables | $\operatorname{Yield}^{b}(\%)$ |
| 1 | None | 91 |
| 2 | DIPEA as base | 57 |
| 3 | 2,6-Lutidine as base | 0 |
| 4 | Pyridine as base | 0 |
| 5 | ^{<i>n</i>} Bn ₃ N as base | 0 |
| 6 | DMF as solvent | 55 |
| 7 | DMA as solvent | 60 |
| 8 | DMSO as solvent | 35 |
| 9 | THF as solvent | 42 |
| 10 | Acetone as solvent | 33 |
| 11 | 10 mol% NaI as additive | 60 |
| 12 | 10 mol% KI as additive | 42 |
| 13 | 0.5 mL CH ₃ CN | 46 |
| 14 | 1a:2=1:2 | 42 |
| 15 | No TBAI | 50 |
| 16 | NO Et ₂ N | 0 |

 a Reaction conditions: 1a (0.40 mmol), 2 (0.20 mmol), TBAI (10 mol%), Et₃N (1 equiv.) in CH₃CN (2 mL) at 35 °C for 16 h. b GC Yields using dodecane as internal standard.

Table 2 Substrate scope of the fluorination reaction of aryl carboxylic $\operatorname{acids}^{a,b}$



^{*a*} Reaction conditions: Reaction conditions: 1 (0.40 mmol), 2 (0.20 mmol), TBAI (10 mol%), Et₃N (1 equiv.) in CH₃CN (2 mL) at 35 °C for 16 h. ^{*b*} Isolated yield.

noxyl (3e) functional groups, reactive chloride (3f), bromide (3g), iodide (3h) were also tolerated in this fluorination protocol, allowing the further sequential functionalization of the generated acyl fluoride products. Aryl carboxylic acid bearing *N*-Boc protected amine was fluorinated in 99% yield (3i). Also, biphenyl carboxylic acids could undergo this transformation with good to high efficiency (3j and 3k). In addition, bicyclic carboxylic acid bearing acetal group (3l), as well as naphathyl carboxylic acid (3 m and 3n), could all give the corresponding product in high yield. Notably, steric hindrance seems to have no significant influence on the reactivity of the substrates (3b, 3c, 3k, and 3n).

Gratefully, our fluorination protocol could be readily extended to vinyl carboxylic acids. As shown in Table 3, the reaction of cinnamic acid proceeded in good yield (5a). Vinyl carboxylic acids bearing methoxyl (5b and 5c) and fluoride (5d) on the aromatic ring underwent the protocol in good to excellent yields. The reactions of vinyl carboxyl acids containing benzodioxole and naphthyl groups (5e and 5f) with *N*-trifluoromethylthiophthalimide took place in good to high yields. Importantly, heterocycles such as furan and thiophene (5e–5i) were also tolerated in this protocol, providing the possi-

Table 3 Substrate scope of the fluorination reaction of vinyl carboxylic acids^{a,b}



^a Reaction conditions: Reaction conditions: 4 (0.40 mmol), 2 (0.20 mmol), TBAI (10 mol%), Et₃N (1 equiv.) in CH₃CN (2 mL) at 35 °C for 16 h. ^b Isolated yield.

bility for the synthesis of pharmaceutical-related molecules. Moreover, larger π -extended vinyl acid could also be converted to the corresponding acyl fluoride in good yield (5j).

In order to demonstrate the practicality of this newly developed fluorination methodology from carboxylic acids, the gram-scale experiment of phenyl naphthalene-2-carboxylate 1i was conducted, and 98% yield of the desired product 3i was obtained (Scheme 2a). Furthermore, we also achieved the sequential deoxygenated fluorination/decarbonylative alkylation/C-O bond arylation, showing great advantages of this protocol (Scheme 2b).

Based on our results and previous studies,^{13,16} a mechanism for this fluorination protocol is proposed (Scheme 3). First, Et₃N attacks the electrophilic sulfur center with the aid of TBAI, affording the phthalimide anion and intermediate I. Next, the phthalimide anion attacks one of the protons of the ethyl group at Et₃N to generate the nucleophilic trifluoromethylthio group along with phthalimide and the aminium. Then the SCF₃ anion degrades to CSF₂ with the release of one fluoride ion. The rapid reaction of CSF₂ with carboxylic acid delivers the intermediate II that was attacked by a fluoride ion to afford the acyl fluoride product.

In summary, we have developed an efficient deoxygenated fluorination of readily available carboxylic acids. In contrast to previous reports wherein bench-stable N-trifluoromethylthiophthalimide was always used as a trifluoromethylthiolation reagent, this newly developed protocol employed it as a fluorinated reagent for the first time. A series of aryl and vinyl car-

a. Gram-scale fluorination reaction



b. Sequential deoxygenative fluorination/decarbonylative alkylation/C-O bond arylation



Scheme 2 Gram-scale reaction and synthetic application.



Scheme 3 Proposed mechanism for the reaction of carboxylic acids and N-trifluoromethylthiophthalimide.

boxylic acid could be converted to acyl fluorides with good to high efficiency. Gram-scale reaction and sequential synthesis, including deoxygenated fluorination/decarbonyl-ative alkylation/C-O bond arylation, were realized in good yield. This protocol provides a good alternative for the fluorination of carboxylic acids.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was financially supported by the King Abdullah University of Science and Technology (KAUST), Saudi Arabia, Office of Sponsored Research (URF/1/4384).

Notes and references

- 1 (a) P. Kirsch, Modern Fluoroorganic Chemistry: Synthesis, Reactivity, Applications, Wiley-VCH, Weinheim, 2004; (b) D. O'Hagan, Understanding organofluorine chemistry. An introduction to the C-F bond, Chem. Soc. Rev., 2008, 37, 308-319; (c) J. T. Welch, Fluorine in Medicinal Chemistry and Chemical Biology, ed. I. Ojima, Wiley-Blackwell, Chichester, 2009; (d) M. C. Belhomme, T. Besset, T. Poisson and X. Pannecoucke, Recent progress toward the introduction of functionalized difluoromethylated building blocks onto C (sp2) and C (sp) centers, Chem. - Eur. J., 2015, 21, 12836-12865; (e) M. V. Ivanova, A. Bayle, T. Besset, T. Poisson and X. Pannecoucke, Copper-Mediated Formation of Aryl, Heteroarvl, Vinyl and Alkynyl Difluoromethylphosphonates: A General Approach to Fluorinated Phosphate Mimics, Angew. Chem., Int. Ed., 2015, 54, 13406-13410; (f) K. Grollier, A. De Zordo-Banliat, F. Bourdreux, B. Pegot, G. Dagousset, E. Magnier and T. Billard, (Trifluoromethylselenyl) methylchalcogenyl as Emerging Fluorinated Groups: Synthesis under Photoredox Catalysis and Determination of the Lipophilicity, Chem. -Eur. J., 2021, 27, 6028-6033; (g) F. Brüning, C. R. Pitts, J. Kalim, D. Bornemann, C. Ghiazza, J. de Montmollin, N. Trapp, T. Billard and A. Togni, Difluoro (aryl) $(perfluoroalkyl) - \lambda 4 - sulfanes and Selanes: Missing Links of$ Trichloroisocyanuric Acid/Potassium Fluoride Chemistry, Angew. Chem., Int. Ed., 2019, 58, 18937-18941; (h) A. Tlili, E. Ismalaj, Q. Glenadel, C. Ghiazza and T. Billard, Synthetic Approaches to Trifluoromethylselenolated Compounds, Chem. - Eur. J., 2018, 24, 3659-3670; (i) X. Zhao, H.-Y. Tu, L. Guo, S. Zhu, F.-L. Qing and L. Chu, Intermolecular selective carboacylation of alkenes via nickel-catalyzed reductive radical relay, Nat. Commun., 2018, 9, 3488; (j) D. Chen, L. Xu, T. Long, S. Zhu, J. Yang and L. Chu, Metal-free, intermolecular carbopyridylation of alkenes via visible-light-induced reductive radical coupling, Chem. Sci., 2018, 9, 9012-9017.
- 2 (a) Y. Ogiwara and N. Sakai, Acyl Fluorides in Late-Transition-Metal Catalysis, Angew. Chem., Int. Ed., 2020, 59, 574–594; (b) Y. Zhang and T. Rovis, A unique catalyst effects the rapid room-temperature cross-coupling of organozinc reagents with carboxylic acid fluorides, chlorides, anhydrides, and thioesters, J. Am. Chem. Soc., 2004, 126, 15964-15965; (c) Y. Ogiwara, Y. Maegawa, D. Sakino and N. Sakai, Palladium-catalyzed coupling of benzoyl halides with aryltrifluorosilanes leading to diaryl ketones, Chem. Lett., 2016, 45, 790-792; (d) Y. Ogiwara, D. Sakino, Y. Sakurai and N. Sakai, Acid fluorides as acyl electrophiles in Suzuki-Miyaura coupling, Eur. J. Org. Chem., 2017, 4324-4327; (e) Y. Ogiwara, Y. Iino and N. Sakai, Catalytic C- H/C- F coupling of azoles and acyl fluorides, Chem. - Eur. J., 2019, 25, 6513-6516; (f) Y. Ogiwara, Y. Sakurai, H. Hattori and N. Sakai, Palladium-catalyzed reductive conversion of acyl fluorides via ligand-controlled decarbonylation, Org. Lett., 2018, 20, 4204–4208; (g) S. T. Keaveney and F. Schoenebeck,

Palladium-Catalyzed Decarbonylative Trifluoromethylation of Acid Fluorides, Angew. Chem., Int. Ed., 2018, 57, 4073-4077; (h) Y. Okuda, J. Xu, T. Ishida, C.-a. Wang and Y. Nishihara, Nickel-catalyzed decarbonylative alkylation of aroyl fluorides assisted by Lewis-acidic organoboranes, ACS Omega, 2018, 3, 13129-13140; (i) C. A. Malapit, J. R. Bour, C. E. Brigham and M. S. Sanford, Base-free nickel-catalysed decarbonylative Suzuki-Miyaura coupling of acid fluorides, Nature, 2018, 563, 100-104; (j) S. Sakurai, T. Yoshida and M. Tobisu, Iridium-catalyzed decarbonylative coupling of acyl fluorides with arenes and heteroarenes via CH activation, Chem. Lett., 2019, 48, 94-97; (k) Z. Wang, X. Wang and Y. Nishihara, Nickel-catalysed decarbonylative borylation of aroyl fluorides, Chem. Commun., 2018, 54, 13969-13972; (l) J. A. Kalow and A. G. Doyle, Enantioselective ring opening of epoxides by fluoride anion promoted by a cooperative dual-catalyst system, J. Am. Chem. Soc., 2010, 132, 3268-3269; (m) Y. Liang, A. Taya, Z. Zhao, N. Saito and N. Shibata, Deoxyfluorination of acyl fluorides to trifluoromethyl compounds by FLUOLEAD®/Olah's reagent under solvent-free conditions, Beilstein J. Org. Chem., 2020, 16, 3052-3058.

- 3 (a) E. Bappert, P. Müller and G. C. Fu, Asymmetric [3 + 2]annulations catalyzed by a planar-chiral derivative of DMAP, Chem. Commun., 2006, 2604-2606; (b) T. Poisson, V. Dalla, C. Papamicael, G. Dupas, F. Marsais and V. Levacher, DMAP-organocatalyzed O-silyl-O-(or C-)benzoyl interconversions by means of benzoyl fluoride, Synlett, 2007, 0381-0386; (c) T. Poisson, V. Dalla, F. Marsais, G. Dupas, S. Oudeyer and V. Levacher, Organocatalytic enantioselective protonation of silvl enolates mediated by Cinchona alkaloids and a latent source of HF, Angew. Chem., Int. Ed., 2007, 46, 7090-7093; (d) S. J. Ryan, L. Candish and D. W. Lupton, N-heterocyclic carbene-catalyzed generation of α , β -unsaturated acyl imidazoliums: synthesis of dihydropyranones by their reaction with enolates, J. Am. Chem. Soc., 2009, 131, 14176-14177; (e) S. J. Ryan, L. Candish and D. W. Lupton, N-heterocyclic carbene-catalyzed (4 + 2) cycloaddition/decarboxylation of silvl dienol ethers with α , β -unsaturated acid fluorides, J. Am. Chem. Soc., 2011, 133, 4694-4697.
- 4 (a) G. Olah and S. Kuhn, Organic Fluorine Compounds. XXVII. Preparation of Acyl Fluorides with Anhydrous Hydrogen Fluoride. The General Use of the Method of Colson and Fredenhagen, *J. Org. Chem.*, 1961, 26, 237– 238; (b) C. Tullock and D. Coffman, Synthesis of fluorides by metathesis with sodium fluoride, *J. Org. Chem.*, 1960, 25, 2016–2019; (c) A. Pittman and D. Sharp, Fluoro Ketone—Metal Fluoride Adducts as Fluorinating Agents in the Preparation of Fluorosilanes and Fluorinated Acyl Fluorides, *J. Org. Chem.*, 1966, 31, 2316–2318.
- 5 H. X. Song, Z. Y. Tian, J. C. Xiao and C. P. Zhang, Tertiary-Amine–Initiated Synthesis of Acyl Fluorides from Carboxylic Acids and CF3SO2OCF3, *Chem. – Eur. J.*, 2020, **26**, 16261–16265.

- 6 Z. Yang, S. Chen, F. Yang, C. Zhang, Y. Dou, Q. Zhou, Y Yan and L. Tang, PPh3/Selectfluor-Mediated Transformation of Carboxylic Acids into Acid Anhydrides and Acyl Fluorides and Its Application in Amide and Ester Synthesis, Eur. J. Org. Chem., 2019, 5998-6002.
- 7 T. Scattolin, K. Deckers and F. Schoenebeck, Direct synthesis of acyl fluorides from carboxylic acids with the bench-stable solid reagent (Me4N) SCF3, Org. Lett., 2017, 19, 5740-5743.
- 8 S. Groß, S. Laabs, A. Scherrmann, A. Sudau, N. Zhang and U. Nubbemeyer, Improved syntheses of cyanuric fluoride and carboxylic acid fluorides, J. Prakt. Chem., 2000, 342, 711-714.
- 9 C. Chen, C.-T. Chien and C.-H. Su, Preparation of acyl fluorides with hydrogen fluoride-pyridine and 1, 3-dicyclohexylcarbodiimide, J. Fluorine Chem., 2002, 115, 75-77.
- 10 (a) A. L'Heureux, F. Beaulieu, C. Bennett, D. R. Bill, S. Clayton, F. LaFlamme, M. Mirmehrabi, S. Tadayon, D. Tovell and M. Couturier, Aminodifluorosulfinium salts: selective fluorination reagents with enhanced thermal stability and ease of handling, J. Org. Chem., 2010, 75, 3401-3411; (*b*) F. Beaulieu, L.-P. Beauregard, F. LaFlamme and G. Courchesne, М. Couturier, A. L'Heureux, Aminodifluorosulfinium tetrafluoroborate salts as stable and crystalline deoxofluorinating reagents, Org. Lett., 2009, 11, 5050-5053.
- El-Faham, 11(a)L. A. Carpino and A. Tetramethylfluoroformamidinium hexafluorophosphate: a rapid-acting peptide coupling reagent for solution and solid phase peptide synthesis, J. Am. Chem. Soc., 1995, 117, 5401-5402; (b) L. A. Carpino, M. Beyermann, H. Wenschuh and M. Bienert, Peptide synthesis via amino acid halides, Acc. Chem. Res., 1996, 29, 268-274.
- 12 Y. Liang, Z. Zhao, A. Taya and N. Shibata, Acyl Fluorides from Carboxylic Acids, Aldehydes, or Alcohols under Oxidative Fluorination, Org. Lett., 2021, 23, 847-852.
- 13 I. Saidalimu, S. Suzuki, E. Tokunaga and N. Shibata, Successive C-C bond cleavage, fluorination, trifluoromethylthio-and pentafluorophenylthiolation under metalfree conditions to provide compounds with dual fluorofunctionalization, Chem. Sci., 2016, 7, 2106-2110.

- 14 (a) S. Munavalli, D. Rohrbaugh, D. Rossman, F. Berg, G. Wagner and H. Durst, Trifluoromethylsulfenylation of masked carbonyl compounds, Synth. Commun., 2000, 30, 2847-2854; (b) T. Bootwicha, X. Liu, R. Pluta, I. Atodiresei and M. Rueping, N-Trifluoromethylthiophthalimide: A Stable Electrophilic SCF3-Reagent and its Application in the Catalytic Asymmetric Trifluoromethylsulfenylation, Chem., Int. Ed., 2013, 52, 12856-12859; Angew. (c) M. Rueping, X. Liu, T. Bootwicha, R. Pluta and C. Merkens, Catalytic enantioselective trifluoromethylthiolation of oxindoles using shelf-stable N-(trifluoromethylthio) phthalimide and a cinchona alkaloid catalyst, Chem. Commun., 2014, 50, 2508-2511; (d) R. Pluta, P. Nikolaienko and M. Rueping, Direct catalytic trifluoromethylthiolation of boronic acids and alkynes employing electrophilic shelf-stable N-(trifluoromethylthio) phthalimide, Angew. Chem., Int. Ed., 2014, 53, 1650-1653; (e) R. Mao, S. Bera, A. Cheseaux and X. Hu, Deoxygenative trifluoromethylthiolation of carboxylic acids, Chem. Sci., 2019, 10, 9555-9559; (f) W. Xu, J. Ma, X. A. Yuan, J. Dai, J. Xie and C. Zhu, Synergistic catalysis for the umpolung trifluoromethylthiolation of tertiary ethers, Angew. Chem., Int. Ed., 2018, 57, 10357-10361; (g) Q. Xiao, Q. He, J. Li and J. Wang, 1, 4-Diazabicyclo [2.2. 2] octane-Promoted Aminotrifluoromethylthiolation of α , β -Unsaturated Carbonyl Compounds: N-Trifluoromethylthio-4-nitrophthalimide Acts as Both the Nitrogen and SCF3 Sources, Org. Lett., 2015, 17, 6090-6093; (h) S. Mukherjee, T. Patra and F. Glorius, Cooperative catalysis: a strategy to synthesize trifluoromethyl-thioesters from aldehydes, ACS Catal., 2018, 8, 5842-5846.
- 15 T. Ji, X.-Y. Chen, L. Huang and M. Rueping, Remote Trifluoromethylthiolation Enabled by Organophotocatalytic C-C Bond Cleavage, Org. Lett., 2020, 22, 2579-2583.
- 16 (a) R. Honeker, J. B. Ernst and F. Glorius, Transition-Metal-Free Trifluoromethylthiolation of N-Heteroarenes, Chem. - Eur. J., 2015, 21, 8047-8051; (b) H. X. Song, Z. Y. Tian, J. C. Xiao and C. P. Zhang, Tertiary-Amine-Initiated Synthesis of Acyl Fluorides from Carboxylic Acids and CF3SO2OCF3, Chem. - Eur. J., 2020, 26, 16261-16265.