

HIGHLIGHT

View Article Online

View Journal | View Issue

Cite this: *Org. Chem. Front.*, 2022, **9**, 5299

Fluorosulfonyl radicals: new horizons for the synthesis of sulfonyl fluorides

Fu-Sheng He,^{*a} Yuqing Li^a and Jie Wu^{†a,b,c}Received 28th July 2022,
Accepted 17th August 2022

DOI: 10.1039/d2qo01211a

rsc.li/frontiers-organic


Sulfonyl fluorides have found widespread applications in organic synthesis, chemical biology, drug discovery and materials science. Compared with current methods, direct fluorosulfonylation with fluorosulfonyl radicals has emerged as a concise and efficient approach for producing sulfonyl fluorides. This highlight provides an overview of recent advances in the generation of fluorosulfonyl radicals from different precursors and their participation in the synthesis of diverse functionalized sulfonyl fluorides.

Sulfonyl fluoride is a prevalent structural unit found in biologically active molecules and versatile synthetic intermediates in modern organic synthesis (Fig. 1).¹ Since the concept of Sulfur (vi) Fluoride Exchange (SuFEx) was pioneered by Sharpless and co-workers as the next generation of click chemistry in 2014,² sulfonyl fluorides have received considerable attention and continuous interest from the communities of synthetic chemistry and medicinal chemistry. Generally, strategies for the construction of these frameworks mainly focus on the fluoride–chloride exchange from the corresponding sulfonyl chlorides.³ Other methods, including conjugate addition,⁴ electrophilic fluorination of thiols,⁵ anodic oxidative fluorination,⁶ and SO₂ insertion/fluorination,⁷ for the preparation of sulfonyl fluorides have also been reported. Compared with these methods *via* S–F bond formation, direct fluorosulfonylation by using FSO₂-containing reagents represents a concise and effective approach for C–SO₂F bond formation. In this case, the reported fluorosulfonylating reagents such as sulfuryl fluoride gas (SO₂F₂)² and other solid reagents (FDIT and AISF)⁸ have been widely utilized as the synthetic equivalents of electrophilic “FSO₂⁺” synthons to access sulfonyl fluorides. In contrast, fluorosulfonylation of the corresponding fluorosulfonyl radical (FSO₂•) is more challenging and remains less well explored.

As one of the sulfur-centered radicals, the highly active fluorosulfonyl radical has been ignored for a long time owing to its instability and difficulty of generation. It was not until 2013 that Zeng, Beckers, and co-workers first observed the for-

mation of the fluorosulfonyl radical by the flash vacuum pyrolysis of fluorosulfonyl azide.⁹ Inspired by this observation, many efforts have been devoted to the development of suitable fluorosulfonyl radical precursors for the direct radical fluorosulfonylation reactions. In this highlight article, we focus on the most recent advances in the fluorosulfonyl radical-involved transformations, with emphasis on the synthesis of sulfonyl fluorides.


The first radical fluorosulfonylation of alkenes using FSO₂Cl as the fluorosulfonyl radical precursor was reported by Liao and co-workers in 2020 (Scheme 1).^{10,11} This photoredox-catalyzed reaction shows a broad substrate scope and excellent functional group compatibility, providing a novel and facile approach for the synthesis of alkenyl sulfonyl fluorides. Notably, the scope of alkenes could be expanded to cyclic, di- and tri-substituted olefins, giving rise to multi-substituted vinyl sulfonyl fluorides that are challenging to be synthesized using established cross-coupling methods. Moreover, the late-stage fluorosulfonylation of natural products including oleic ester, cinnamate, chromene, estrone, pregnenolone, and cholesterol also demonstrated the superiority of this approach. The

Fig. 1 Selected examples of SO₂F-containing bioactive molecules.

^aSchool of Pharmaceutical and Chemical Engineering & Institute for Advanced Studies, Taizhou University, 1139 Shifu Avenue, Taizhou 318000, China. E-mail: hefs@tzc.edu.cn, jie_wu@fudan.edu.cn

^bState Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Road, Shanghai 200032, China


^cSchool of Chemistry and Chemical Engineering, Henan Normal University, Xinxiang 453007, China



Scheme 1 Radical fluorosulfonylation of alkenes with FSO_2Cl under photoredox conditions.

reaction mechanism for this visible-light-induced radical fluorosulfonylation of alkenes **1** with FSO_2Cl **2** was supported by the experimental and DFT calculation studies. Initially, single-electron reduction of FSO_2Cl by the excited $\text{Ir}(\text{III})^*$ would occur to generate fluorosulfonyl radical **4**, which then attacked the double bond of alkene **1** leading to carbon radical intermediate **5**. In the case of a radical chain process, the radical would react with FSO_2Cl to form chlorinated intermediate **6**. Following the loss of HCl , fluorosulfonylation product **3** would be afforded. On the other hand, a redox pathway that involved a single-electron oxidation/deprotonation process could not be excluded.


However, the above radical fluorosulfonylation of multi-substituted alkenes usually suffered from side reactions, low reaction efficiency, and low *Z/E* selectivity. To address these issues, the same group subsequently developed a photoredox-catalyzed radical chloro-fluorosulfonylation of alkynes for the construction of β -chloro alkenylsulfonyl fluorides (BCASF) (Scheme 2).¹² Under the optimal reaction conditions, both terminal and internal alkynes reacted smoothly in this transformation. More importantly, this protocol provided a novel and powerful class of sulfonyl fluoride hubs, which could be converted to a series of functionalized alkenylsulfonyl fluorides. As a result, BCASF molecules could undergo different types of transformations at the β -chloro site while maintaining the sulfonyl fluoride group intact, such as Suzuki coupling, Sonogashira coupling, selective reduction, and nucleophilic substitution with nitrogen, oxygen, and sulfur nucleophiles. Furthermore, this versatile platform could also be applied in the late-stage modification of peptides and drugs. Similar to their previous work, it was postulated that fluorosulfonyl




Scheme 2 Visible light-induced photoredox-catalyzed chloro-fluorosulfonylation of alkynes.

radical **4** could be first generated from FSO_2Cl **2** by single-electron reduction. Then, addition of fluorosulfonyl radical **4** to alkyne **8** produced vinyl radical intermediate **10**, which could attack FSO_2Cl **2** to deliver the target product **3** and initiate the radical chain process. Meanwhile, vinyl radical **10** might undergo hydrogen-atom transfer from Et_2O leading to alkenylsulfonyl fluoride **11** and radical **12**. A single-electron transfer of the oxidized $\text{Ir}(\text{IV})$ by ether-derived radical **12** would give rise to carbocation intermediate **13** and regenerate the $\text{Ir}(\text{III})$ photocatalyst. Finally, GC-MS detection of α -chlorinated diethyl ether **14** suggested the addition of the chloride anion to the carbocation.


In the same year, Liao, Huang, and co-workers disclosed a practical synthesis of β -keto sulfonyl fluorides by electrochemical oxo-fluorosulfonylation of alkynes with FSO_2Cl (Scheme 3).¹³ For the first time, the fluorosulfonyl radical could be generated under electrochemical conditions. This strategy featured mild reaction conditions, broad substrate scope, air as the oxidant and high chemoselectivity. Remarkably, the divergent synthesis of β -keto sulfonyl fluorides and α -chloro β -keto sulfonyl fluorides from the same starting materials was achieved simply by the variation of solvents. Additionally, some β -keto sulfonyl fluoride derivatives showed potent activities toward *Bursaphelenchus xylophilus* and *Colletotrichum gloeosporioides*. Regarding the reaction mechanism, it was suggested that cathodic reduction of FSO_2Cl **2** would produce fluorosulfonyl radical **4**, which then attacked the triple bond of alkyne **15** leading to vinyl radical **18**. Vinyl radical **18** would react with O_2 to generate peroxy radical **19**, which could be converted to radical **20** according to the Russell mechanism. Subsequently, radical **20** undergoes a



At the same time, Wang and co-workers reported the same air- and bench-stable benzoimidazolium fluorosulfonate reagent from their previous work on the chemistry of imidazolium sulfonate cationic salts,¹⁶ which successfully applied to the radical fluorosulfonylation of unsaturated hydrocarbons (Scheme 6).¹⁷ Under suitable optimized photocatalytic con-



Scheme 5 Photoredox-catalyzed radical fluorosulfonylation of alkenes with a benzoimidazolium-based fluorosulfonyl radical precursor.




Scheme 6 Synthesis of alkenyl and alkyl sulfonate fluorides with a benzoimidazolium-based fluorosulfonyl radical precursor.

ditions, this redox-active fluorosulfonyl radical precursor could react with various alkenes smoothly, providing alkenyl sulfonate fluoride, alkylsulfonate fluoride, and migratory fluorosulfonylation


products in good yields with high stereoselectivity. This operationally simple protocol enabled the efficient introduction of the FSO₂ group into a wide range of complex molecules derived from natural products and drugs. The combination of radical trapping and radical clock experiments demonstrated the generation of the fluorosulfonyl radical during the reaction.

Soon after, using the same 1-fluorosulfonyl benzoimidazolium triflate salt as the fluorosulfonyl radical precursor, the Liao group further developed a radical hydro-fluorosulfonylation of alkenes by employing oxygen-doped anthanthrene (ODA) as the photocatalyst and 1,4-cyclohexadiene (1,4-CHD) as a hydrogen donor (Scheme 7).¹⁸ This method enabled the synthesis of various aliphatic sulfonate fluorides **40** under mild and metal-free conditions, which could be applied to the late-stage functionalization of natural products, peptides and drugs. Moreover, with (TMS)₃SiH as a hydrogen donor, the application of this photocatalytic system to the radical hydro-fluorosulfonylation of alkynes was also demonstrated, allowing the formation of alkenylsulfonate fluorides **41** with high Z-selectivity.

Given the similar bifunctional radical reactivity between alkynyl sulfonate fluorides and alkynyl triflates, Studer and co-workers recently disclosed an efficient transition-metal-free radical 1,2-difunctionalization of unactivated alkenes to access various aliphatic β-alkynylfluorosulfonylalkanes (Scheme 8).¹⁹ Notably, alkynyl sulfonate fluorides could not only be used as highly valuable bifunctional radical trapping reagents but also as the fluorosulfonyl radical precursor. The successful diversification of products with SuFEx click chemistry to sulfonates and sulfonamides also highlighted the practicality of this method. Mechanistically, it was proposed that the reaction was initiated by AIBN to give AIBN-derived cyanopropyl radical **45**. Following the addition to alkyne **43** and radical fragmentation



Scheme 7 Photoredox-catalyzed radical hydro-fluorosulfonylation of unactivated alkenes and alkynes.




Scheme 8 Radical difunctionalization of unactivated alkenes to access β -alkynyl-fluorosulfonylalkanes.

sequence, fluorosulfonyl radical **4** would be generated, which would undergo addition to alkene **42** leading to radical intermediate **46**. This radical intermediate **46** could be trapped by alkynyl sulfonyl fluoride **43** giving rise to vinyl radical **47**, which underwent β -fragmentation to provide the desired product **44** with the regeneration of fluorosulfonyl radical **4**.

Very recently, Li, Su, and co-workers developed a photocatalytic procedure for the preparation of β -keto sulfonyl fluorides from vinyl fluorosulfates (Scheme 9).²⁰ This methodology featured sustainable conditions and a broad substrate scope, affording aromatic β -keto sulfonyl fluorides in moderate to good yields by using $\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})(\text{PF}_6)$ (1 mol%) as a photocatalyst under irradiation of 3 W blue LEDs. According to the mechanistic results, two plausible pathways for this photocatalytic transformation of vinyl fluorosulfates were proposed as shown in Scheme 9. In path A, the $\text{Ir}(\text{III})$ complex was excited under visible light irradiation to form $\text{Ir}(\text{III})^*$, which would then undergo energy transfer with vinyl fluorosulfate **48** to generate enol radical **50** and fluorosulfonyl radical **4**. Subsequently, the corresponding product **49** would be produced through recombination of these two radical species. The authors also suggested another plausible pathway through which the excited $\text{Ir}(\text{III})^*$ and vinyl fluorosulfate **48** would undergo SET to afford $\text{Ir}(\text{IV})$ and a radical anion intermediate **51**. Following radical fragmentation/reconstruction and SET oxidation, product **49** would be formed (path B).

The direct fluorosulfonylation process involving the fluorosulfonyl radical has appeared as an attractive strategy for the rapid assembly of sulfonyl fluorides. In this highlight, the most recent progress in this aspect by utilizing different fluorosulfonyl radical precursors is discussed. Specifically, four types



Scheme 9 Synthesis of β -keto sulfonyl fluorides via photocatalytic transformation of vinyl fluorosulfates.

of radical fluorosulfonylating reagents including FSO_2Cl , benzoimidazolium fluorosulfonates, alkynyl sulfonyl fluorides and vinyl fluorosulfates have been developed and applied to the efficient construction of diverse functionalized sulfonyl fluorides. These achievements would offer new opportunities for SuFEx click chemistry, accelerating the applications of sulfonyl fluorides in chemical biology and drug discovery. There are still some challenges remaining in this field. First, the substrate scope of radical acceptors in the above-mentioned reactions is relatively narrow, which is mostly restricted to alkenes and alkynes. Second, the fluorosulfonyl radical precursors are still limited to the above four types. The development of novel and efficient radical precursors would provide new approaches for the construction of sulfonyl fluorides in a more sustainable way. Therefore, this highly reactive intermediate is worthy of further investigation. Besides that, enantioselective radical fluorosulfonylation involving the fluorosulfonyl radical to deliver chiral sulfonyl fluorides remains undeveloped. The exploration of enantioselective radical fluorosulfonylation is highly in demand. Given these challenges and opportunities, further progress can be expected in the near future.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Financial support from the National Natural Science Foundation of China (No. 22101199), the Natural Science Foundation of Zhejiang Province (LQ21B020002), the Leading

Innovative and Entrepreneur Team Introduction Program of Zhejiang (No. 2019R01005), and the Open Research Fund of School of Chemistry and Chemical Engineering, Henan Normal University (2020ZD04) is gratefully acknowledged.

References

- (a) A. S. Barrow, C. J. Smedley, Q. Zheng, S. Li, J. Dong and J. E. Moses, The growing applications of SuFEx click chemistry, *Chem. Soc. Rev.*, 2019, **48**, 4731–4758; (b) C. Lee, A. J. Cook, J. E. Elisabeth, N. C. Friede, G. M. Sammis and N. D. Ball, The Emerging Applications of Sulfur(VI) Fluorides in Catalysis, *ACS. Catal.*, 2021, **11**, 6578–6589; (c) T. S.-B. Lou and M. C. Willis, Sulfonyl fluorides as targets and substrates in the development of new synthetic methods, *Nat. Rev. Chem.*, 2022, **6**, 146–162.
- J. Dong, L. Krasnova, M. G. Finn and K. B. Sharpless, Sulfur(VI) Fluoride Exchange (SuFEx): Another Good Reaction for Click Chemistry, *Angew. Chem., Int. Ed.*, 2014, **53**, 9430–9448.
- (a) L. Xu and J. Dong, Click Chemistry: Evolving on the Fringe, *Chin. J. Chem.*, 2020, **38**, 414–419; (b) Q. Zheng, J. Dong and K. B. Sharpless, Ethenesulfonyl Fluoride (ESF): An On-Water Procedure for the Kilogram-Scale Preparation, *J. Org. Chem.*, 2016, **81**, 11360–11362.
- (a) J. Chen, B.-Q. Huang, Z.-Q. Wang, X.-J. Zhang and M. Yan, Asymmetric Conjugate Addition of Ethylene Sulfonyl Fluorides to 3-Amido-2-oxindoles: Synthesis of Chiral Spirocyclic Oxindole Sultams, *Org. Lett.*, 2019, **21**, 9742–9746; (b) J. Leng and H.-L. Qin, 1-Bromoethene-1-sulfonyl fluoride (1-Br-ESF), a new SuFEx clickable reagent, and its application for regioselective construction of 5-sulfonylfluoro isoxazoles, *Chem. Commun.*, 2018, **54**, 4477–4480; (c) R. Xu, T. Xu, M. Yang, T. Cao and S. Liao, A rapid access to aliphatic sulfonyl fluorides, *Nat. Commun.*, 2019, **10**, 3752; (d) X. Zhang, W.-Y. Fang, R. Lekkala, W. Tang and H.-L. Qin, An Easy, General and Practical Method for the Construction of Alkyl Sulfonyl Fluorides, *Adv. Synth. Catal.*, 2020, **362**, 3358–3363.
- (a) L. Wang and J. Cornella, A Unified Strategy for Arylsulfur (VI) Fluorides from Aryl Halides: Access to Ar-SOF₃ Compounds, *Angew. Chem., Int. Ed.*, 2020, **59**, 23510–23515; (b) S. W. Wright and K. N. Hallstrom, A Convenient Preparation of Heteroaryl Sulfonamides and Sulfonyl Fluorides from Heteroaryl Thiols, *J. Org. Chem.*, 2006, **71**, 1080–1084; (c) L. Zhang, X. Cheng and Q.-L. Zhou, Electrochemical Synthesis of Sulfonyl Fluorides with Triethylamine Hydrofluoride, *Chin. J. Chem.*, 2022, **40**, 1687–1692.
- (a) G. Laudadio, A. de A. Bartolomeu, L. M. H. M. Verwijlen, Y. Cao, K. T. de Oliveira and T. Noël, Sulfonyl fluoride synthesis through electrochemical oxidative coupling of thiols and potassium fluoride, *J. Am. Chem. Soc.*, 2019, **141**, 11832–11836; (b) A. Shavnya, S. B. Coffey, K. D. Hesp, S. C. Ross and A. S. Tsai, Reaction of Alkyl Halides with Rongalite: One-Pot and Telescoped Syntheses of Aliphatic Sulfonamides, Sulfonyl Fluorides, and Unsymmetrical Sulfones, *Org. Lett.*, 2016, **18**, 5848–5851; (c) A. Shavnya, K. D. Hesp and A. S. Tsai, A Versatile Reagent and Method for Direct Aliphatic Sulfonylation, *Adv. Synth. Catal.*, 2018, **360**, 1768–1774.
- (a) Y. Liu, D. Yu, Y. Guo, J. C. Xiao, Q. Y. Chen and C. Liu, Arenesulfonyl Fluoride Synthesis via Copper-Catalyzed Fluorosulfonylation of Arenediazonium Salts, *Org. Lett.*, 2020, **22**, 2281–2286; (b) T. Zhong, J. Yi, Z. Chen, Q. Zhuang, Y. Li, G. Lu and J. Weng, Photoredox-catalyzed Aminofluorosulfonylation of Unactivated Olefins, *Chem. Sci.*, 2021, **7**, 9359–9365; (c) P. J. Sarver, N. B. Bissonnette and D. W. C. MacMillan, Decatungstate-Catalyzed C(sp³)-H Sulfonylation: Rapid Access to Diverse Organosulfur Functionality, *J. Am. Chem. Soc.*, 2021, **143**, 9737–9743; (d) Z. Ma, Y. Liu, X. Ma, X. Hu, Y. Guo, Q.-Y. Chen and C. Liu, Aliphatic sulfonyl fluoride synthesis via reductive decarboxylative fluorosulfonylation of aliphatic carboxylic acid NHPI esters, *Org. Chem. Front.*, 2022, **9**, 1115–1120; (e) V. T. Nguyen, G. C. Haug, V. D. Nguyen, N. T. H. Vuong, G. B. Karki, H. D. Arman and O. V. Larionov, Functional group divergence and the structural basis of acridine photocatalysis revealed by direct decarboxysulfonylation, *Chem. Sci.*, 2022, **13**, 4170–4179.
- (a) T. Guo, G. Meng, X. Zhan, Q. Yang, T. Ma, L. Xu, K. B. Sharpless and J. Dong, A New Portal to SuFEx Click Chemistry: A Stable Fluorosulfonyl Imidazolium Salt Emerging as an “F-SO₂⁺” Donor of Unprecedented Reactivity, Selectivity, and Scope, *Angew. Chem., Int. Ed.*, 2018, **57**, 2605–2610; (b) H. Zhou, P. Mukherjee, R. Liu, E. Evrard, D. Wang, J. M. Humphrey, T. W. Butler, L. R. Hoth, J. B. Sperry, S. K. Sakata, C. J. Helal and C. W. Am Ende, Introduction of a Crystalline, Shelf-Stable Reagent for the Synthesis of Sulfur (VI) Fluorides, *Org. Lett.*, 2018, **20**, 812–815.
- X. Q. Zeng, H. Beckers and H. Willner, Thermally Persistent Fluorosulfonyl Nitrene and Unexpected Formation of the Fluorosulfonyl Radical, *J. Am. Chem. Soc.*, 2013, **135**, 2096–2099.
- X. Nie, T. Xu, J. Song, A. Devaraj, B. Zhang, Y. Chen and S. Liao, Radical Fluorosulfonylation: Accessing Alkenyl Sulfonyl Fluorides from Alkenes, *Angew. Chem., Int. Ed.*, 2021, **60**, 3956–3960.
- X. Nie and S. Liao, Radical Fluorosulfonylation: Accessing Alkenylsulfonyl Fluorides from Alkenes and Alkynes, *Synlett*, 2022, 401–408.
- X. Nie, T. Xu, Y. Hong, H. Zhang, C. Mao and S. Liao, Introducing A New Class of Sulfonyl Fluoride Hubs via Radical Chloro-Fluorosulfonylation of Alkynes, *Angew. Chem., Int. Ed.*, 2021, **60**, 22035–22042.
- D. Chen, X. Nie, Q. Feng, Y. Zhang, Y. Wang, Q. Wang, L. Huang, S. Huang and S. Liao, Electrochemical Oxo-Fluorosulfonylation of Alkynes under Air: Facile Access to β -Keto Sulfonyl Fluorides, *Angew. Chem., Int. Ed.*, 2021, **60**, 27271–27276.
- Q. Feng, Y. Fu, Y. Zheng, S. Liao and S. Huang, Electrochemical Synthesis of β -Keto Sulfonyl Fluorides via

- Radical Fluorosulfonylation of Vinyl Triflates, *Org. Lett.*, 2022, **24**, 3702–3706.
- 15 P. Wang, H. Zhang, X. Nie, T. Xu and S. Liao, Photoredox catalytic radical fluorosulfonylation of olefins enabled by a bench-stable redox-active fluorosulfonyl radical precursor, *Nat. Commun.*, 2022, **13**, 3370.
 - 16 W. Zhang, Z. Zou, W. Zhao, S. Lu, Z. Wu, M. Huang, X. Wang, Y. Wang, Y. Liang, Y. Zhu, Y. Zheng and Y. Pan, Integrated redox-active reagents for photoinduced regio- and stereoselective fluorocarbonylation, *Nat. Commun.*, 2020, **11**, 2572.
 - 17 W. Zhang, H. Li, X. Li, Z. Zou, M. Huang, J. Liu, X. Wang, S. Ni, Y. Pan and Y. Wang, A practical fluorosulfonylating platform via photocatalytic imidazolium-based SO₂F radical reagent, *Nat. Commun.*, 2022, **13**, 3515.
 - 18 P. Wang, H. Zhang, M. Zhao, S. Ji, L. Lin, N. Yang, X. Nie, J. Song and S. Liao, Radical Hydro-Fluorosulfonylation of Unactivated Alkenes and Alkynes, *Angew. Chem., Int. Ed.*, 2022, **61**, e202207684.
 - 19 N. L. Frye, C. G. Daniliuc and A. Studer, Radical 1-Fluorosulfonyl-2-alkynylation of Unactivated Alkenes, *Angew. Chem., Int. Ed.*, 2022, **61**, e202115593.
 - 20 J. Cui, S. Ke, J. Zhao, S. Wu, W. Luo, S. Xu, X. Su and Y. Li, Photocatalytic access to aromatic keto sulfonyl fluorides from vinyl fluorosulfates, *Org. Chem. Front.*, 2022, **9**, 3540–3545.