


 Cite this: *RSC Adv.*, 2022, 12, 15190

 Received 6th March 2022
 Accepted 11th May 2022

DOI: 10.1039/d2ra01472f

rsc.li/rsc-advances

A facile metal-free one-flask synthesis of multi-substituted furans *via* a $\text{BF}_3 \cdot \text{Et}_2\text{O}$ mediated formal [4 + 1] reaction of 3-chloro-3-phenyldiazirines and α, β -alkenyl ketones†

 Zixin Zhang,^a Aimin Huang,^a Lin Ma,^a Jian-hua Xu^b and Min Zhang^{*,a}

A facile, efficient and metal free one-flask approach to diversely substituted furans from easily accessible 3-chloro-3-phenyldiazirines and α, β -alkenyl ketones is reported. This protocol integrates three steps of cyclopropanation, Cloke–Wilson rearrangement and elimination of HCl in one-flask to give products in moderate to good yields. It provides a metal and oxidant free approach to multi-substituted furans with the advantages of easy operation, mild reaction conditions and a broad scope of substrates.

Introduction


Furan is a five-membered oxygenated heteroaromatic which is widely spread in natural products¹ and plays an important role in both pharmaceutical chemistry^{2,3} and in synthetic organic chemistry as a useful building block.^{2,4–6} Long attracting the interests of chemists, a number of synthetic methods for furan have been developed,^{3,7–12} including the cyclodehydration of dicarbonyl compounds through Paal–Knorr synthesis¹³ and Feist–Bénary cyclocondensation.^{14,15} Nevertheless, more selective approaches to multi-substituted furans under mild conditions still remain a challenging task.

In the past decades, cyclopropane has been widely used as a three-carbon synthon to access variable chemicals due to the readiness of ring-opening from high angle and torsion strain and tunable reactivity by substituent-controlled C–C bond polarization/cleavage.^{16–21} Transition-metal catalyzed intramolecular ring-opening cycloisomerization of cyclopropenyl ketones^{22,23} or alkylidene cyclopropyl ketones^{24–26} has been proved to be a very successful and reliable approach to furans (Scheme 1A). Early in 2003, Ma and Zhang²² developed a regioselective cycloisomerization of cyclopropenyl ketones using copper(i) or Pd catalysts. Later in 2004, they developed a Pd mediated ring-opening cycloisomerization of 2-methylene- or alkylidene cyclopropyl ketone to di- or tri-substituted furans.²⁴ In 2007, Liang group reported a synthesis of trisubstituted furans *via* a Cu(i)-catalyzed formal [4 + 1] cycloaddition of α, β -alkynyl ketones with diazoacetates.²⁷ Xu and co-workers further


developed a Cu–Pd relay catalysis to access tetra-substituted furans from cyclopropanes.²⁸ These elegant transition-metal catalyzed methods are advantageous in both atom economy and efficiency. However, using alternative non-metal catalysts to promote cycloisomerization is very essential in account of economic, environmental and sustainability requirement. Recently, Wang and coworkers^{29–31} have developed $\text{I}_2/\text{K}_2\text{CO}_3$ or DBU mediated ring opening and cyclization of cyano-substituted cyclopropyl ketones to afford furan derivatives.

The Cloke–Wilson rearrangement (CWR) reaction has been intensively used to access dihydrofurans from cyclopropyl ketones.^{32–37} Besides transition-metal catalysis,^{38,39} CWR reaction can also be promoted by Lewis acid,^{40,41} photocatalysis,^{42–44} and organo-catalysis.^{26,32,33} Regrettably, an extra dehydrogenation procedure is a prerequisite to transform dihydrofurans to furans using stoichiometric oxidants such as DDQ.^{45,46} To avoid


A. Ma and Zhang: transition-metal catalyzed cycloisomerization of cyclopropanes.



B. Design for the synthesis of furan from halocyclopropyl ketone:



C. This work: a formal one-flask [1+4] reaction.



Scheme 1 Synthesis of furan from cyclopropanes by literature and this work.

^aSchool of Chemistry and Chemical Engineering, Guangxi University, Nanning, Guangxi 530004, China. E-mail: chemizhang@gxu.edu.cn

^bSchool of Chemistry and Chemical Engineering, Nanjing University, Nanjing, Jiangsu 210093, China

 † Electronic supplementary information (ESI) available. See <https://doi.org/10.1039/d2ra01472f>



the harmful oxidation procedure, we envisioned that halocyclopropyl ketone is an ideal alternative precursor to perform CWR reaction to deliver halogenated dihydrofuran, which is converted to furan *via* elimination of HX (Scheme 1B). However, there are few reports making use of this elimination strategy. As an elegant example, Namboothiri's group reported that acidic Al₂O₃ could be used to promote CWR reaction of dibromocyclopropyl ketone followed by elimination to access 3-bromo furans.^{47,48}

Taking advantage of the fact that the highly reactive halocarbene (RCX) derived from the easily available 3-halodiazirines (RCN₂X) upon the loss of N₂ readily take part in [2 + 1] cycloaddition with alkenes to give halogenated cyclopropanes,^{49–52} we design a formal [4 + 1] approach to furans using this cyclopropanation method to obtain the required halocyclopropyl ketone precursors (Scheme 1C). Firstly, photolysis or thermolysis of 3-halo-3-phenyldiazirine **1** in the presence of α,β -alkenyl ketone **2** gives the halocyclopropyl ketone **3**, which is then subjected to a tandem CWR–elimination reaction sequence to afford the furan **4**. We further succeeded in using the same Lewis acid to promote both the CWR and the elimination reactions to facilitate the reaction procedure. As a result, we herein report a one-flask metal-free synthetic approach to a diversity of di-, tri- or tetra-substituted furans from a series of 3-halo-3-phenyldiazirines and α,β -alkenyl ketones *via* cyclopropanation and BF₃·Et₂O mediated CWR–elimination reactions.

Results and discussions

Initially, the synthesis of halocyclopropyl ketones was investigated (Scheme 2). Photolysis or thermolysis of 3-chloro-3-(4-chlorophenyl)diazirine (**1a**) was used to generate phenylchlorocarbene (PhCCl) *in situ*, which rapidly reacted with chalcone (**2a**) to give the halocyclopropyl ketone diastereomers (**3a/3a'**). After an optimization of solvents and temperatures (for details, please see Table S1†), either photolysis at room temperature or thermolysis at 80 °C in 1,2-dichloroethane (DCE) gave the halocyclopropyl ketones (**3a/3a'**) in highest yield with similar diastereoselectivity.

Next, the transformation of halocyclopropyl ketone **3a** (major isomer) to furan **4a** was investigated and selected results are summarized in Table 1 (for more details, see Table S2†). Lewis acid promoters FeCl₃·6H₂O, TiCl₄ and FeCl₂·4H₂O were the most efficient catalysts for this conversion (0.08–0.5 h), while BF₃·Et₂O, AlCl₃, SnCl₄ and Sc(OTf)₃ promoted the reaction less efficiently (15–45 h). Delightfully, furan **4a** was obtained in excellent yields (85–98%) in the presence of these seven catalysts (entries 1–7). On the contrary, BiCl₃ couldn't complete this transformation in 72 h and gave **4a** in lower yield



Scheme 2 Synthesis of halo-cyclopropyl ketone **3a/3a'**.

Table 1 Screening of Lewis acids for **4a**^a

Entry	Lewis acid	Time, h	Yield, ^b %
1	TiCl ₄	0.16	86
2	FeCl ₂ ·4H ₂ O	0.5	96
3	FeCl ₃ ·6H ₂ O	0.08	86
4	BF ₃ ·Et ₂ O	15	98
5	AlCl ₃	17	90
6	SnCl ₄	21	87
7	Sc(OTf) ₃	45	86
8	BiCl ₃	72	68 ^c
9	None	24	NR
10	PTSA	36	90
11 ^d	FeCl ₂ ·4H ₂ O	1	97
12 ^e	BF ₃ ·Et ₂ O	16	98
13 ^{d,e}	FeCl ₂ ·4H ₂ O	1	97

^a Reagents and conditions: halocyclopropyl ketone **3a** (0.06 mmol), Lewis acid (0.06 mmol, 1 eq.) in 5 mL DCE was heated at 80 °C in a 38 mL reaction tube equipped with a condenser until the reaction was completed by TLC monitoring. NR = no reaction. RSM = recovery of starting material. ^b Isolated yield. ^c Yield is based on consumed halocyclopropyl ketone. RSM was 16%. ^d 0.2 eq. LA was used. ^e **3a'** was used instead of **3a**.

(entry 8). No reaction could take place in the absence of a Lewis acid (entry 9). Brønsted acid *para*-toluene sulfonic acid (PTSA) could also mediate this reaction to give **4a** in 90% yield in 36 h (entry 10). Therefore, among these promoters, BF₃·Et₂O and FeCl₂·4H₂O showed the best catalytic activity to give nearly quantitative yields of **4a**. It is also noted that the halocyclopropyl ketone diastereomer **3a'** was similarly converted to **4a** in the nearly quantitative yield as **3a** in the presence of either BF₃·Et₂O or FeCl₂·4H₂O (entries 12–13). Therefore, there is no need to separate two diastereomers **3a/3a'** for the transformation to **4a**. We then succeeded in implementing these reaction steps in one-flask (for details, please see Table S3†) with BF₃·Et₂O (1 eq.) as the best catalyst, which was added into the flask after the completion of cyclopropanation to avoid side reactions. This one-flask protocol gave **4a** in an overall yield of 68%.


With this optimized one-flask conditions in hand, we investigated the scope of *para*-substituted phenylchlorodiazirines **1** (Table 2). Unsubstituted (R = H) or substituted phenylchlorodiazirines with either electron-donating (R = Me, OMe) or slightly electron withdrawing (R = F, Cl) groups on the phenyl ring gave furans (**4a–4e**) in good yields (57–71%). However, the cyclopropyl ketones with strong electron-withdrawing substituents such as CF₃ or CN (**3f**, **3g**) need to be heated in *n*-octane at 120 °C to give furans in reasonable yields (**4f** 43%; **4g** 42%) due to lower ring opening reactivity for less polarization character of C–C bond. This conversion could also be driven by the powerful FeCl₃·6H₂O and gave furans in better yields (**4f** 57%; **4g** 53%). However, the reaction of 3-benzyl-3-chloro-diazirine (PhCH₂CCLN₂) and



chalcone couldn't afford the expected furan. The scope of chalcones were also investigated: chalcones with substituents on either phenyl ring gave furans (**5a–5i**; **5m–5t**) in good yields (70–80%), no matter they are electron-withdrawing or electron-donating. This one-flask strategy can also be applied to the naphthyl, thiophenyl or pyridinyl substituted chalcones to afford furans (**5j–5l**, **5u**) with moderate to good yields (40–78%).

We further extended this one-flask reaction to a wide range of alkenyl ketones with alkyl groups, and the corresponding furans were obtained in moderate to good yields (Table 3). Alkenyl ketones substituted with a methyl group at R₁, R₂ or R₃ position gave trisubstituted furans **6a–6c** in good yields (60–67%). Methyl vinyl ketone (MVK) gave 2-methyl-5-phenyl furan **6d** in 45% yield. Notably, this protocol enabled an astonishing access to tetra-substituted furans with structural complexity (**6e**, **6f**). For example, 2-benzylidencyclohexan-1-one gave tetra-substituted furan **6f** with a fused ring in good yield (50%). This protocol can also be applied to α,β -unsaturated aldehydes, e.g., cinnamaldehyde was used to synthesize 2,3-disubstituted furan **6g** in 29% yield. Bis(2-phenylvinyl) ketone gave furan **6h** in 42% yield, exemplifying the functional group tolerance for another sensitive C=C double bond. Step-by-step analysis of these two-stage reactions (Table S5†) reveals that the lower yields were owing to the poor cyclopropanation reactivity because of less electronic richness (**6d**, **6g**) or steric hindrance (**6e**) of the C=C double bond, in which a considerable amount of carbene dimer was often generated as side product. Therefore, this one-flask protocol can use a variety of α,β -unsaturated carbonyl substrates to synthesize 2,3- or 2,5-disubstituted, 2,3,5-trisubstituted and even 2,3,4,5-tetrasubstituted furans with moderate to good yields.

To probe the mechanism of these reactions, β -methyl chalcone was subjected to this one-flask reaction (Scheme 3A). Unlike the α -methyl chalcone, the CWR–elimination reaction of the cyclopropyl ketone promoted by BF₃ gave a complicated


Table 3 Scope of alkyl substituted alkenyl ketones^{a,b}

^a Reagents and conditions: using method A as above unless specified.

^b Isolated yield of one-flask reaction. ^c **1a** (0.2 mmol) was reacted with 2 eq. alkenyl ketone (0.4 mmol). ^d Two equivalents of alkenyl ketone were used and reactions were performed at 60 °C in both stages.


mixture without any furan product, indicating the necessity of a β -hydrogen. Reactivities of other 3-halo-3-phenyldiazirines (X = Br, F) were also studied (Scheme 3B). 3-Bromo-3-phenyldiazirine (**1ab**) gave bromocyclopropyl ketone **3ab/3ab'** in 2 h (72%), which gave furan **4a** in 95% yield in 17 h with the similar reactivity as **3a/3a'** (X = Cl). It indicates that BF₃ is supposed to bind with the oxygen in carbonyl group instead of halogen to promote the CWR reaction, leading to no significant difference in the reactivities between **3a** and **3ab** (Scheme 4, path a). On the other hand, 3-fluoro-3-phenyldiazirine (**1ac**) gave the cyclopropyl ketone **3ac** (59%) much slower (48 h) owing to the less electrophilicity and stability of phenylfluorocarbene (PhCF).^{53,54} Moreover, **3ac** is quite ready to give furan **4a** in 90% yield with excellent reactivity (2 h). This efficient transformation is supposed to be attributed to a different pathway because of the high affinity between BF₃ and fluorine (*vide infra*).

Based on these experiments and literature,^{34,47,55} a plausible mechanism is proposed in Scheme 4. Upon thermolysis or photolysis, 3-halo-3-phenyldiazirine (**1**) generates electrophilic singlet phenylhalocarbene (PhCX) with the loss of nitrogen


Table 2 Scope of substrates^{a,b}

^a Reagents and conditions (method A): 3-aryl-3-chlorodiazirine **1** (0.2 mmol), alkenyl ketone **2** (0.2 mmol) in 5 mL DCE was heated at 80 °C in a 38 mL reaction tube with a condenser until the reaction was completed (usually 2 h). BF₃·Et₂O (0.2 mmol, 1 eq.) was added in and kept on heating to complete the transformation. ^b Isolated yield of one-flask reaction. ^c Reacted at 120 °C in *n*-octane and BF₃·Et₂O (5 eq.) was used.

A. β -H of chalcone is required.




B. 3-Halo-3-phenyldiazirine.



Reactant	Time1, h	Yield1, %	Time2, h	Yield2, %	Overall yield, %
1a	2	79	17	98	68
1ab	2	72	17	95	62
1ac	48	59	2	91	40

Scheme 3 Control experiments.





Scheme 4 Plausible mechanism.

(N_2).^{51,56} The PhCX carbene reacts rapidly with α,β -alkenyl ketone (2) to afford halocyclopropyl ketone (3) via a [2 + 1] cycloaddition. Subsequent addition of $BF_3 \cdot Et_2O$ catalyzes the CWR rearrangement of chloro- or bromocyclopropyl ketone 3a/3ab by complexing with the carbonyl oxygen in 3 (path a) to facilitate the heterolytic cleavage of this donor–acceptor cyclopropane to give the key zwitterion intermediate I. Then, an intramolecular cyclization of I by nucleophilic attack of oxyanion to carbocation gives dihydrofuran II, which is converted to furan 4 after the loss of HX with the aid of BF_3 . In the case of fluorocyclopropyl ketone 3ac, the ring-opening might be driven by the loss of tetrafluoroborate (BF_4^-) and proceeds through a cyclopropyl carbocation mechanism in a similar intramolecular cyclization mode (path b).

Conclusions

In conclusion, we have developed a facile one-flask approach to the di-, tri- and even tetra-substituted furans in moderate to good yields from readily available starting materials using inexpensive boron trifluoride as catalyst. This metal and oxidant free method involves the cyclopropanation of α,β -alkenyl ketones with phenylchlorocarbene, BF_3 mediated ring-opening cycloisomerization (Cloke–Wilson rearrangement) and elimination of HCl to give the multi-substituted furans. This method has the advantages of simple operation, mild reaction conditions and a broad scope of substrates, which provides a concise approach to diversified biologically and synthetically useful furans. We believe it will benefit the discovery of new application of furan derivatives.

Author contributions

Z. Zhang: most of the experimental work and writing of ESI.† A. Huang & L. Ma: methodology and discussion. J. Xu: manuscript revision and discussion. M. Zhang: conceptualization, funding acquisition, supervision, and writing, review, and editing of the manuscript.

Conflicts of interest

The authors declare no competing financial interest.

Acknowledgements

We thank the National Natural Science Foundation of China (No. 21861002, 21462003) for financial support.

Notes and references

- Z. Batool, D. Xu, X. Zhang, X. Li, Y. Li, Z. Chen, B. Li and L. Li, A review on furan: formation, analysis, occurrence, carcinogenicity, genotoxicity and reduction methods, *Crit. Rev. Food Sci.*, 2021, **61**, 395–406.
- H.-K. Lee, K.-F. Chan, C.-W. Hui, H.-K. Yim, X.-W. Wu and H. N. C. Wong, Use of furans in synthesis of bioactive compounds, *Pure Appl. Chem.*, 2005, **77**, 139–143.
- A. Blanc, V. Bénéteau, J.-M. Weibel and P. Pale, Silver & gold-catalyzed routes to furans and benzofurans, *Org. Biomol. Chem.*, 2016, **14**, 9184–9205.
- B. H. Lipshutz, Five-membered heteroaromatic rings as intermediates in organic synthesis, *Chem. Rev.*, 1986, **86**, 795–819.
- H. N. C. Wong, P. Yu and C. Y. Yick, The use of furans in natural product syntheses, *Pure Appl. Chem.*, 1999, **71**, 1041–1044.
- A. S. Makarov, M. G. Uchuskin and I. V. Trushkov, Furan Oxidation Reactions in the Total Synthesis of Natural Products, *Synthesis*, 2018, **50**, 3059–3086.
- X. L. Hou, H. Y. Cheung, T. Y. Hon, P. L. Kwan, T. H. Lo, S. Y. Tong and H. N. C. Wong, Regioselective syntheses of substituted furans, *Tetrahedron*, 1998, **54**, 1955–2020.
- R. C. D. Brown, Developments in Furan Syntheses, *Angew. Chem., Int. Ed.*, 2005, **44**, 850–852.
- S. F. Kirsch, Syntheses of polysubstituted furans: recent developments, *Org. Biomol. Chem.*, 2006, **4**, 2076–2080.
- N. T. Patil and Y. Yamamoto, Coinage Metal-Assisted Synthesis of Heterocycles, *Chem. Rev.*, 2008, **108**, 3395–3442.
- A. Deepthi, B. P. Babu and A. L. Balachandran, Synthesis of Furans - Recent Advances, *Org. Prep. Proced. Int.*, 2019, **51**, 409–442.
- D. X. Duc, Recent Progress in the Synthesis of Furan, *Mini-Rev. Org. Chem.*, 2019, **16**, 422–452.
- G. Minetto, L. F. Raveglia, A. Sega and M. Taddei, Microwave-Assisted Paal–Knorr Reaction – Three-Step Regiocontrolled Synthesis of Polysubstituted Furans, Pyrroles and Thiophenes, *Eur. J. Org. Chem.*, 2005, 5277–5288.
- M. A. Calter, C. Zhu and R. J. Lachicotte, Rapid Synthesis of the 7-Deoxy Zalgozic Acid Core, *Org. Lett.*, 2002, **4**, 209–212.
- G. Mross, E. Holtz and P. Langer, Synthesis of 2-Alkenyl-3-(alkoxycarbonyl)furans Based on Feist–Benary Cyclocondensation of (2,4-Dioxobutylidene)phosphoranes with α -Haloketones and α -Chloroacetaldehyde, *J. Org. Chem.*, 2006, **71**, 8045–8049.



- 16 H.-U. Reissig and R. Zimmer, Donor–Acceptor-Substituted Cyclopropane Derivatives and Their Application in Organic Synthesis, *Chem. Rev.*, 2003, **103**, 1151–1196.
- 17 M. A. Cavitt, L. H. Phun and S. France, Intramolecular donor–acceptor cyclopropane ring-opening cyclizations, *Chem. Soc. Rev.*, 2014, **43**, 804–818.
- 18 T. F. Schneider, J. Kaschel and D. B. Werz, A New Golden Age for Donor–Acceptor Cyclopropanes, *Angew. Chem., Int. Ed.*, 2014, **53**, 5504–5523.
- 19 K. R. Babu, X. He and S. Xu, Lewis Base Catalysis Based on Homoconjugate Addition: Rearrangement of Electron-Deficient Cyclopropanes and Their Derivatives, *Synlett*, 2020, **31**, 117–124.
- 20 Y. Xia, X. Liu and X. Feng, Asymmetric Catalytic Reactions of Donor–Acceptor Cyclopropanes, *Angew. Chem., Int. Ed.*, 2021, **60**, 9192–9204.
- 21 T. R. McDonald, L. R. Mills, M. S. West and S. A. L. Rousseaux, Selective Carbon–Carbon Bond Cleavage of Cyclopropanols, *Chem. Rev.*, 2021, **121**, 3–79.
- 22 S. Ma and J. Zhang, 2,3,4- or 2,3,5-Trisubstituted furans: catalyst-controlled highly regioselective ring-opening cycloisomerization reaction of cyclopropenyl ketones, *J. Am. Chem. Soc.*, 2003, **125**, 12386–12387.
- 23 A. Padwa, J. M. Kassir and S. L. Xu, Rhodium-catalyzed ring-opening reaction of cyclopropenes. Control of regioselectivity by the oxidation state of the metal, *J. Org. Chem.*, 2002, **56**, 6971–6972.
- 24 S. Ma and J. Zhang, Tuning the Regioselectivity in the Palladium(II)-Catalyzed Isomerization of Alkylidene Cyclopropyl Ketones: A Dramatic Salt Effect, *Angew. Chem., Int. Ed.*, 2003, **42**, 183–187.
- 25 S. Ma, L. Lu and J. Zhang, Catalytic Regioselectivity Control in Ring-Opening Cycloisomerization of Methylene- or Alkylidenecyclopropyl Ketones, *J. Am. Chem. Soc.*, 2004, **126**, 9645–9660.
- 26 X. He, Y. Tang, Y. Wang, J. B. Chen, S. Xu, J. Dou and Y. Li, Phosphine-Catalyzed Activation of Alkylidenecyclopropanes: Rearrangement to Form Polysubstituted Furans and Dienones, *Angew. Chem., Int. Ed.*, 2019, **58**, 10698–10702.
- 27 L.-B. Zhao, Z.-H. Guan, Y. Han, Y.-X. Xie, S. He and Y.-M. Liang, Copper-Catalyzed [4 + 1] Cycloadditions of α,β -Acetylenic Ketones with Diazoacetates to Form Trisubstituted Furans, *J. Org. Chem.*, 2007, **72**, 10276–10278.
- 28 C. Song, J. Wang and Z. Xu, Tandem metal relay catalysis: from cyclopropene to polysubstituted furan, *Org. Biomol. Chem.*, 2014, **12**, 5802–5806.
- 29 W. Ye, C. Tan, J. Yao, S. Xue, Y. Li and C. Wang, Iodine-Promoted Domino Reactions of 1-Cyanocyclopropane 1-Esters: A Straightforward Approach to Fully Substituted 2-Aminofurans, *Adv. Synth. Catal.*, 2016, **358**, 426–434.
- 30 J.-M. Liu, X.-Y. Liu, X.-S. Qing, T. Wang and C.-D. Wang, I_2/K_2CO_3 -Promoted ring-opening/cyclization/rearrangement/aromatization sequence: a powerful strategy for the synthesis of polysubstituted furans, *Chin. Chem. Lett.*, 2017, **28**, 458–462.
- 31 Z. Su, Z. Xie, S. Wang, N. Luo and C. Wang, Direct synthesis of highly functionalized furans from donor-acceptor cyclopropanes via DBU-mediated ring expansion reactions, *Org. Biomol. Chem.*, 2019, **17**, 7342–7351.
- 32 J. Zhang, Y. Tang, W. Wei, Y. Wu, Y. Li, J. Zhang, Y. Zheng and S. Xu, Organocatalytic Cloke–Wilson Rearrangement: DABCO-Catalyzed Ring Expansion of Cyclopropyl Ketones to 2,3-Dihydrofurans, *Org. Lett.*, 2017, **19**, 3043–3046.
- 33 W. Wei, Y. Tang, Y. Zhou, G. Deng, Z. Liu, J. Wu, Y. Li, J. Zhang and S. Xu, Recycling Catalyst as Reactant: A Sustainable Strategy to Improve Atom Efficiency of Organocatalytic Tandem Reactions, *Org. Lett.*, 2018, **20**, 6559–6563.
- 34 Q. Shi, Y. Wang and D. Wei, Theoretical study on DABCO-catalyzed ring expansion of cyclopropyl ketone: mechanism, chemoselectivity, and role of catalyst, *Comput. Theor. Chem.*, 2018, **1123**, 20–25.
- 35 A. Ortega, R. Manzano, U. Uria, L. Carrillo, E. Reyes, T. Tejero, P. Merino and J. L. Vicario, Catalytic Enantioselective Cloke–Wilson Rearrangement, *Angew. Chem., Int. Ed.*, 2018, **57**, 8225–8229.
- 36 H. Nambu, Y. Onuki, N. Ono and T. Yakura, Iodide-Catalyzed Ring-Opening Cyclization of Cyclohexane-1,3-dione-2-spirocyclopropanes, *Adv. Synth. Catal.*, 2018, **360**, 2938–2944.
- 37 X. Liang, P. Guo, W. Yang, M. Li, C. Jiang, W. Sun, T. P. Loh and Y. Jiang, Stereoselective synthesis of trifluoromethyl-substituted 2H-furan-amines from enaminones, *Chem. Commun.*, 2020, **56**, 2043–2046.
- 38 R. K. Bowman and J. S. Johnson, Nickel-catalyzed rearrangement of 1-acyl-2-vinylcyclopropanes. A mild synthesis of substituted dihydrofurans, *Org. Lett.*, 2006, **8**, 573–576.
- 39 M. L. Piotrowski and M. A. Kerr, Tandem Cyclopropanation/Vinyllogous Cloke–Wilson Rearrangement for the Synthesis of Heterocyclic Scaffolds, *Org. Lett.*, 2018, **20**, 7624–7627.
- 40 V. K. Yadav and R. Balamurugan, Silicon-assisted ring opening of donor-acceptor substituted cyclopropanes. An expedient entry to substituted dihydrofurans, *Org. Lett.*, 2001, **3**, 2717–2719.
- 41 Y. Y. Zhu, M. L. Zhang, T. Li and X. X. Song, $AlCl_3$ -Promoted Stereospecific Cloke–Wilson Rearrangement of Spirocyclopropyl Barbiturates for the Synthesis of Substituted Dihydrofuro[2,3-d]pyrimidines, *ChemistrySelect*, 2019, **4**, 10838–10842.
- 42 D. Kalpogiannaki, C.-I. Martini, A. Nikopoulou, J. A. Nyxas, V. Pantazi and L. P. Hadjarapoglou, Fused dihydrofurans from the one-pot, three-component reaction of 1,3-cyclohexanedione, iodobenzene diacetate and alkenes, *Tetrahedron*, 2013, **69**, 1566–1575.
- 43 C.-H. Lin, D. Pursley, J. E. M. N. Klein, J. Teske, J. A. Allen, F. Rami, A. Köhn and B. Plietker, Non-decarbonylative photochemical versus thermal activation of $Bu_4N[Fe(CO)_3(NO)]$ – the Fe-catalyzed Cloke–Wilson rearrangement of vinyl and arylcyclopropanes, *Chem. Sci.*, 2015, **6**, 7034–7043.
- 44 J. Luis-Barrera, V. Laina-Martín, T. Rigotti, F. Peccati, X. Solans-Monfort, M. Sodupe, R. Mas-Ballesté, M. Liras and J. Alemán, Visible-Light Photocatalytic Intramolecular



- Cyclopropane Ring Expansion, *Angew. Chem., Int. Ed.*, 2017, **56**, 7826–7830.
- 45 N. M. Barl, E. Sansiaume-Dagousset, G. Monzón, A. J. Wagner and P. Knochel, Preparation and Reactions of Heteroarylmethylzinc Reagents, *Org. Lett.*, 2014, **16**, 2422–2425.
- 46 A. Lévesque, T. Maris and J. D. Wuest, ROY Reclaims Its Crown: New Ways to Increase Polymorphic Diversity, *J. Am. Chem. Soc.*, 2020, **142**, 11873–11883.
- 47 N. Satam, S. Nemu, G. N. Gururaja and I. N. N. Namboothiri, Substrate-oriented selectivity in the Mg-mediated conjugate addition of bromoform to electron-deficient alkenes, *Org. Biomol. Chem.*, 2020, **18**, 5697–5707.
- 48 E. Gopi and I. N. N. Namboothiri, Synthesis of Fused Bromofurans via Mg-Mediated Dibromocyclopropanation of Cycloalkanone-Derived Chalcones and Cloke–Wilson Rearrangement, *J. Org. Chem.*, 2013, **78**, 910–919.
- 49 R. A. Moss, Diazirines: Carbene Precursors Par Excellence, *Acc. Chem. Res.*, 2006, **39**, 267–272.
- 50 M. Zhang, R. A. Moss, J. Thompson and K. Krogh-Jespersen, Evolution of Structure and Reactivity in a Series of Ionic Carbenes, *J. Org. Chem.*, 2012, **77**, 843–850.
- 51 L. Wang, K. Krogh-Jespersen and R. A. Moss, Activation Parameters for Additions to Alkenes of Arylchlorocarbenes with Enhanced Electrophilicity, *J. Org. Chem.*, 2015, **80**, 7590–7593.
- 52 E. M. D. Allouche and A. B. Charette, Cyclopropanation Reactions of Semi-stabilized and Non-stabilized Diazo Compounds, *Synthesis*, 2019, **51**, 3947–3963.
- 53 R. A. Moss, Carbenic selectivity in cyclopropanation reactions, *Acc. Chem. Res.*, 1980, **13**, 58–64.
- 54 I. R. Gould, N. J. Turbo, J. Butcher, C. Doubleday, N. P. Hacker, G. F. Lehr, R. A. Moss, D. P. Cox, W. Guo, R. C. Munjal, L. A. Perez and M. Fedorynski, Time-resolved flash spectroscopic investigations of the reactions of singlet arylhalocarbenes, *Tetrahedron*, 1985, **41**, 1587–1600.
- 55 T. Li, D. Yan, C. Cui, X. Song and J. Chang, A thermal decarboxylative Cloke–Wilson rearrangement of dispirocyclopropanes derived from para-quinone methides and bromo-Meldrum's acids: an approach to synthesize spirobutyrolactone para-dienones, *Org. Chem. Front.*, 2020, **7**, 2682–2688.
- 56 G. Richter, E. Mendez-Vega and W. Sander, Singlet Halophenylcarbenes as Strong Hydrogen-Bond Acceptors, *J. Phys. Chem. A*, 2016, **120**, 3524–3532.

