



Cite this: *RSC Adv.*, 2022, **12**, 18615

Heterocycles from cyclopropenones

Ashraf A. Aly,¹ Alaa A. Hassan,¹ Sara M. Mostafa,¹ Asmaa. H. Mohamed,¹ Esraa M. Osman¹ and AbdElAziz A. Nayl²

Great attention has been paid to cyclopropenones as they are present in many natural sources. Various synthesized cyclopropenone derivatives also show a wide range of biological activities. The cyclopropenone derivatives undergo a variety of reactions such as ring-opening reactions, isomerization reactions, C–C coupling reactions, C–H activation, cycloaddition reactions, thermal and photo-irradiation reactions, and acid–base-catalyzed reactions under the influence of various chemical reagents (electrophiles, nucleophiles, radicals, and organometallics) and external forces (heat and light). Many previous reviews have dealt with the chemistry and reactions of cyclopropenones. However and to the best of our knowledge, the utility of cyclopropenones in the synthesis of heterocycles has not been reported before. Therefore, it would be interesting to shed light on this new topic. The present review article provides, for the first time, a comprehensive compilation of synthetic methods for the synthesis of various heterocyclic ring systems, as a significant family in the field of organic chemistry.

Received 12th May 2022
 Accepted 2nd June 2022

DOI: 10.1039/d2ra03011j
rsc.li/rsc-advances

1 Introduction

Cyclopropenone is an important building block for the construction of a large number of skeletons.¹ Very recently, new developments have been made using cyclopropenone and its heteroanalogues, such as [3 + *n*] annulation reactions, acylation, organocatalytic reactions, metal catalytic reactions, base-mediated reactions, nucleophilic substitution reactions, light-induced reactions, σ -bond cross-exchange reactions and C–H activation reactions for the synthesis of diverse heterocycles. Besides, they are used as catalysts in a few reactions. The activation of C–C bonds is a powerful concept for the reorganization or coupling of organic scaffolds, yet it is a relatively challenging process to achieve in the context of synthetic methodology because of their inherent stability.² In order to enable such methods, one can use C–C-strained, often cyclic, building blocks that are consequently spring-loaded for C–C bond activation.^{3–22}

This review highlights the recent applications of cyclopropenone ring-expansion reactions aiming to synthesize various products (essentially various classes of heterocycles). Since there is no extensive comprehensive review concerned with the chemistry of cyclopropenone derivatives in the construction of heterocycles, the present study would be of great interest.

¹Chemistry Department, Faculty of Science, Minia University, El-Minia 61519, Egypt.
 E-mail: ashrafaly63@yahoo.com; ashraf.shehata@mu.edu.eg; alaahassan2001@mu.edu.eg; sara.ahmed@mu.edu.eg; esraamah33@gmail.com; asmaa.hamouda@mu.edu.eg

²Department of Chemistry, College of Science, Jouf University, P. O. Box 2014, Sakaka, Aljouf, Saudi Arabia. E-mail: aanayel@ju.edu.sa; aanayl@yahoo.com

2 Chemistry

Cyclopropenone (**1a**) (Fig. 1) is a cyclic organic ketone with the molecular formula C_3H_2O composed of a cyclopropene with a ketone functional group. Cyclopropenone (**1a**) is an aromatic compound that polymerizes at room temperature.²³

The stability of 2,3-diphenylcyclopropenone (**1b**) increases when the substituents are aryl groups. The possible resonance structures of 2,3-diphenylcyclopropenone (**1b**) (Fig. 2) are shown in **A–C** (equivalent to **D**), which contain a three-membered ring of sp^2 carbon coupled to the electron-donor phenyl group in order to stabilize these structures (Fig. 2).^{24–25}

Since the first synthesis of cyclopropenone (**1**) was done by Breslow,²⁶ organic chemists have been quite interested in it. Cyclopropenones are commonly employed as electrophile-trapping agents.^{27,28} Due to their high strain^{29,30}, cyclopropenones easily participate in cycloaddition,³¹ ring-opening,^{32–34} and ring-enlargement^{35–37} reactions. They are also used as organocatalysts in the conversion of aldoximes to nitriles.³⁸ In addition, they are used as Lewis bases for the organocatalytic transformation of alcohols into alkyl chlorides.³⁹ The effect of UV irradiation on cyclopropenones and their analogues results in efficient decarbonylation and the generation of the corresponding alkynes.^{40–45} Visible light




Fig. 1 Structure of cyclopropenone (**1a**).






Fig. 2 Resonance structures of 2,3-diphenylcyclopropenone (1b).

causes cyclopropenones to undergo decarbonylation and form alkynes.⁴⁶ Cyclopropenones can form metal complexes *via* chelation with transition metals at the oxygen center or at the double bond.⁴⁷

2.1. Natural products containing cyclopropenone (1) and biological investigation of cyclopropenones

Numerous extracted natural products have cyclopropenone moieties such as 2-(hydroxymethyl)-cycloprop-2-enone (penitricin) (**A**),⁴⁸ 2-((8S,8aR)-8,8a-dimethyl-1,2,3,4,6,7,8,8a-octahydronaphthalen-2-yl)cycloprop-2-enone (**B**) and 2-((2R,4aR,8aS)-4a-methyl-8-methylenedecahydronaphthalen-2-yl)cycloprop-2-enone (**C**) (Fig. 3).⁴⁹ Compound **A** was extracted from *Penicillium aculeatum*, whereas compounds **B** and **C** were extracted from plant sources. It was found that penitricin (**A**) is the only one that showed biological activity as an antibiotic agent.⁵⁰

Besides, alutacenoic acid **A** (**D**) and alutacenoic acid **B** (**E**) (Fig. 4) are naturally occurring molecules that contain cyclopropenone structures. They were isolated from common fungi such as *Eupenicillium alutaceum*.⁵¹

Other types of isolated naturally occurring products **F**, **G** and **H** (Fig. 5) having cyclopropenone moieties are 2-(1-hydroxypropyl)cycloprop-2-enone (**F**), 2-(1-hydroxyoctyl)-cycloprop-2-enone (**G**), and 2-(1-hydroxyhexyl)cycloprop-2-enone (**H**) (Scheme 5). Such compounds showed antibacterial activity more than penitricin (**A**).⁵²

Some studies reported in the literature illustrated the utility of diphenylcyclopropenone derivatives for dermatological treatments such as *alopecia areata*, *alopecia totalis*, and *alopecia universalis*.^{53,54} Moreover they were used for the treatment of *recalcitrant warts*,⁵⁵ and others showed antitumor activities in the treatment of B16 melanoma.^{56,57}

2.2. Synthesis of cyclopropenone (1) and its derivatives

2.2.1. From acetals or their formation. Synthesis of cyclopropenone (**1a**) was achieved *via* selective dechlorination using $(C_4H_9)_3SnH$ on perchlorocycloprop-1-ene (**2**) to give 3,3-dichloro-cycloprop-1-ene (**3**). Hydrolysis of **3** led to the formation of compound **1a** in 46% yield (Scheme 1).⁵⁸

The most effective procedure for the synthesis of cyclopropenones was found during the treatment of compounds **4a–e** with equal amounts of boron trifluoride (BF_3) as a Lewis acid in ether to yield substituted cyclopropenones **1c–g** in moderate to excellent yields (Scheme 2).⁵⁹

Another convenient method was known to obtain substituted cyclopropenones **1** in 70–96% yield, *via* the hydrolysis of 6,6-dimethyl-4,8-dioxaspiro[2.5]oct-1-ene derivatives **5** (cyclopropenone acetals) using Amberlyst-15 in acetone or in aqueous tetrahydrofuran (THF) at room temperature (Scheme 3).⁶⁰

Breslow *et al.* also reported on the synthesis of diphenylcyclopropenone (**1b**) by treatment of (2,2-dimethoxyvinyl)benzene (**6**) with (dichloromethyl)benzene in the presence of potassium




Fig. 3 Naturally occurring compounds A–C containing cyclopropenone molecules.




Fig. 4 Naturally occurring acids containing cyclopropenone structures.






Fig. 5 Structures of naturally occurring compounds F–H.




Scheme 1 Synthesis of cyclopropenone (**1a**) from perchlorocycloprop-1-ene (**2**). Reagents and conditions: **A** = $(C_4H_9)_3SnH$, **B** = H_2O .





Scheme 2 Synthesis of substituted cyclopropenones **1c–g** using BF_3 . *Reagents and conditions:* A = BF_3 , Et_2O .



Scheme 3 Hydrolysis of 6,6-dimethyl-4,8-dioxaspiro[2.5]oct-1-ene derivatives **5** into **1**. *Reagents and conditions:* A = Amberlyst-15, MeCOMe or aq. THF , rt.

tert-butoxide ($\text{KO-}t\text{-Bu}$) to obtain 3,3-dimethoxy-1,2-diphenylcyclopropene (**7**) as an intermediate, which was converted by hydrolysis to give **1b** in 80% yield (Scheme 4).²⁵

An alternative method for the synthesis of **1b** was applied during the cycloaddition of phenylmethoxy-acetylene (**8a**) with (dichloromethyl)benzene in the presence of $\text{KO-}t\text{-Bu}$ (Scheme 5).⁶¹

Cyclopropenones **1h,i** were formed in a low yield (4–7%) *via* the reaction between acetylenes **8b,c** and sodium trichloroacetate (Cl_3COONa) using dimethylethane (DME), as shown in Scheme 6.⁶²

McGarry *et al.* used a rapid injection NMR technique to observe the formation of the cyclopropenium intermediate (**9**) *via* hydrolysis of 1,1-diethoxy-2,3-diphenylcycloprop-2-ene (**7**) in slightly acidic aqueous acetone to obtain diphenylcyclopropane (**1b**) (Scheme 7).⁶³

2.2.2. Bromination and/or elimination reaction of bromoketonic compounds. Various substituted cyclopropenones **1j** were synthesized during the reaction of 1-phenyl-2-butanone (**10**) with bromine in DCM and Et_3N . As an example, 2-methyl-3-phenylcycloprop-2-enone (**1**) was obtained in 45% yield (Scheme 8).⁶⁴


The Favorskii reaction was also used to synthesize 2,3-diphenylcyclopropenone (**1b**) in 45% yield during elimination of HBr by the action of Et_3N on dibromodibenzyl ketone (**11a**).

The reaction mechanism was the formation of intermediates **12** and **13** (Scheme 9).⁶⁵


Previously, Curnow *et al.* reported that 2,3-diisopropylcycloprop-2-enone (**1k**) was also obtained in a low yield (18%), *via* dehydrobromination of 3,5-dibromo-2,6-dimethylheptan-4-one (**11b**) using NaH in THF as a solvent, followed by treatment with aqueous HCl (Scheme 10).⁶⁶

2.2.3. Different methods of preparation. 2,3-Bis(methyl-phenyl)-amino)cycloprop-2-enone (**1l**) was formed in 22% yield (Scheme 11) *via* the hydrolysis of *N*-(2,3-bis(methyl(phenyl)amino)cyclopropylidene)-*N*-methylbenzenaminium (**14**) using aqueous KOH. The reaction was proceeded *via* the formation of intermediate **15**.⁶⁷


The reaction mixture of **2** with naphthalene and ferrocene in DCM in the presence of aluminum chloride (AlCl_3) afforded



Scheme 5 Synthesis of **1b** from the cycloaddition of phenylmethoxy-acetylene (**8a**) with (dichloromethyl)benzene. *Reagents and conditions:* A = (i) $\text{KO-}t\text{-Bu}$, (ii) $\text{H}_2\text{O}/\text{H}^+$.




Scheme 6 Formation of cyclopropenone **1h,i** *via* the reaction of acetylenes **8b,c** with Cl_3COONa . *Reagents and conditions:* A = (i) Cl_3CCOONa , DME, reflux, (ii) H_2O .




Scheme 4 Synthesis of 2,3-diphenylcyclopropenone (**1b**). *Reagents and conditions:* A = $\text{KO-}t\text{-Bu}$, B = $\text{H}_2\text{O}/\text{H}^+$.





Scheme 7 Synthesis of cyclopropanone **1b**. Reagents and conditions: A = MeCOMe, H₂O/H₃O⁺.



Scheme 8 Synthesis of 2-methyl-3-phenylcycloprop-2-enone (**1j**). Reagents and conditions: A = Br₂, CH₂Cl₂, Et₃N, 0 °C.

among other products, diferrocenylcyclopropenone (**1m**) in 41% yield (Scheme 12).⁶⁸

During heating of 1,2,3-trichlorocycloprop-2-en-1-ylum aluminum(III) chloride (**16**) in a mixture of benzene/H₂O, 2,3-diphenylcyclopropenone (**1b**) was obtained in 67% yield (Scheme 13).⁶⁹ As compound **16** underwent an electrophilic aromatic substitution (Friedel-Crafts alkylation) during reaction with benzene, which upon heating, *gem*-dichlorodiphenylcyclopropene was obtained. Then, the formed intermediate underwent hydrolysis (during workup) to afford 2,3-diphenylcyclopropenone (**1b**).

Decomposition of 1,3-bis(diazo)-1,3-diphenylpropan-2-one (**17**) in methanol in the presence of Ag₂O yielded compounds **18** and **19**, in addition to 2,3-diphenylcyclopropenone (**1b**), in a 11% yield (Scheme 14).⁷⁰

2.3. Utility of cyclopropenone derivatives in the synthesis of various heterocyclic compounds

2.3.1. Synthesis of four-membered rings with one heteroatom. An example of heterocyclic ring containing an heteroatom was reported.⁷¹ For example, the cyclization reaction of 2-(2-hydroxypropan-2-yl)-3-methylcycloprop-2-enone (**1n**) in CD₃OD and triphenylphosphine (PPh₃) yielded 3-ethylidene-4,4-dimethyloxetan-2-one (**20**) in 60% yield, whereas substituted furanone **21** was also formed as a side product with 39% yield (Scheme 15).⁷¹

2.3.2. Synthesis of five-membered rings with one heteroatom


2.3.2.1. Synthesis of pyrrolones. Under microwave (MW) irradiation, the reaction of primary enaminone derivatives **22a-g** with **1b** catalyzed by Bi(NO₃)₃·5H₂O afforded the corresponding 2-pyrrolinone derivatives **23a-g** in 42–80% yield. The reaction was performed in toluene in the presence of bismuth nitrate Bi(NO₃)₃ as a catalyst (Scheme 16).⁷²

The suggested mechanism involves the formation of pyrrolinones **23a-g** initiated *via* coordination between Bi(III) and the oxygen atom in **1b** (Scheme 17). Then, the nitrogen atom of enaminone **22** would attack on carbonyl carbon of **1b** *via* hard-hard interaction A.⁷² Proton shift to adduct **B** then occurred to yield intermediate **C**. Thereafter, adduct **C** underwent simultaneous ring expansion and Michael reaction, resulting in the formation of enol **D**, which finally gave 2-pyrrolinones **23a-g** (Scheme 17).⁷²


Reaction of (Z)-4-((2,2-dimethoxyethyl)amino)pent-3-en-2-one (**24**) and **1b** in refluxing toluene for 6 d, proceeded to give 1-(2,2-dimethoxyethyl)-5-methyl-5-(2-oxopropyl)-3,4-diphenyl-1*H*-pyrrol-2(5*H*)-one (**25**) in 71% yield (Scheme 18).⁷³

3-Pyrrolinone derivatives **27a-c** were successfully synthesized in a moderate yield of 42–67%, *via* the reaction of diimines **26a-c** with 2,3-diphenylcyclopropenone (**1b**) in dry ethanol for 2–5 h (Scheme 19).⁷⁴

The mechanism of the formation of 3-pyrrolinone **27a-c** is proposed and illustrated in Scheme 20. First, one of the imino






Scheme 10 Synthesis of 2,3-diisopropylcycloprop-2-enone (**1k**). Reagents and conditions: A = (i) NaH, THF, (overnight) and (ii) HCl/H₂O.



Scheme 9 Synthesis of diphenylcyclopropenone (**1b**) from dibromodibenzyl ketone (**11a**).



Scheme 11 Formation of 2,3-bis(methyl(phenyl)-amino)cycloprop-2-enone (11). Reagents and conditions: A = KOH/H₂O, MeOH, rt, 40 h.Scheme 12 Synthesis of diferrocenylcyclopropenone (1m). Reagents and conditions: A = AlCl₃, CH₂Cl₂, H₂O.Scheme 13 Synthesis of 2,3-diphenylcyclopropenone (1b) from (16). Reagents and conditions: A = benzene, H₂O.



nitrogen atoms of **26a–c** was added to C-2 of **1b** to obtain the intermediate (A). The ketene intermediate (B) was then obtained after ring opening, and then imines (C) were formed by cyclization *via* the ketene attack on iminium function, and finally, 3-pyrrolinones **27a–c** were formed.⁷⁴

Haito and Chatani used $[\text{Rh}(\text{OAc})(\text{cod})]_2$ as a catalyst in the reaction between *N*-(pyridin-2-ylmethyl)pentanamide (**28a**) and 2,3-diphenylcyclopropenone (**1b**). The reaction was performed in toluene in the presence of 2-phenylbenzoic acid (2-*Ph*C₆H₄-COOH) to produce 5-butyl-5-hydroxy-1-(pyridin-2-ylmethyl)-1*H*-pyrrol-2(5*H*)-one (**29a**) as a major product and also gave 5-butylidene-1-(pyridin-2-ylmethyl)-1*H*-pyrrol-2(5*H*)-one (**29b**) as a side product (Scheme 21).⁷⁵


In continuation to the methods dealing with the synthesis of five-membered rings with one heteroatom, using cyclopropenone (**1**), Haito and Naoto Chatani reacted *N*-(pyridin-2-ylmethyl)benzamide (**28b**) with diphenylcyclopropenone (**1b**) in the presence of $[\text{Rh}(\text{OAc})(\text{cod})]_2$ as a catalyst. 3,4,5-Triphenyl-1-(pyridin-2-ylmethyl)-1*H*-pyrrol-2(5*H*)-one (**29c**) and 5-hydroxy-


3,4,5-triphenyl-1-(pyridin-2-ylmethyl)-1*H*-pyrrol-2(5*H*)-one (**29d**) were obtained in 80% and 9% yields, respectively (Scheme 22).⁷⁵

In 2020, Nanda *et al.* have reported that the reaction of **1b** with various anilines **30a–q** in the presence of palladium acetate


Scheme 15 Synthesis of 3-ethylidene-4,4-dimethyloxetan-2-one (20). Reagents and conditions: A = PPh₃ (5 mol%), CD₃OD, 25 °C.

22 and 23
a: R = H, R¹ = Me
b: R = H, R¹ = OMe
c: R = H, R¹ = OEt
d: R = n-Bu, R¹ = Me
e: R = C₆H₁₁, R¹ = Me
f: R = CH₂CH₂OH, R¹ = Me
g: R = CH₂CH₂OH, R¹ = OEt


Scheme 16 Synthesis of 2-pyrrolinones **23a–g**. Reagents and conditions: A = MW, toluene, Bi(NO₃)₃·5H₂O, 30–90 min.Scheme 14 Oxidation of **17** into diphenylcyclopropenone (**1b**), **18** and **19**. Reagents and conditions: A = MeOH, Ag₂O.



Scheme 17 Mechanism of the formation of 2-pyrrolinones 23a–g.



Scheme 18 Synthesis of compound 25. Reagents and conditions: A = toluene, reflux, 6 d.



Scheme 19 Synthesis of 3-pyrrolinones 27a–c. Reagents and conditions: A = EtOH, 2–5 h.

as the catalyst, tetrabutylammonium bromide as an additive, and sodium acetate as the base at 120 °C for 12 h in DMF (0.25 M) gave substituted pyrroles 31a–q in 38–80% yield (Scheme 23).⁷⁶

When 1-arylideneamino-2,2,2-trichloroethanols 32a–f were subjected to **1b** at room temperature in methanol, the reaction produced 2,2'-diaryl-4,4',5,5'-tetraphenyl-1,1',2,2'-tetrahydro-3H,3'H-2,2'-bipyrrole-3,3'-diones 35a–f in 23–84% yield (Scheme 24), through the formation of intermediates 33 and 34, respectively.⁷⁷

2.3.2.2. Synthesis of furanones. A series of substituted butenolides were successfully synthesized *via* the reaction between

1b and β -ketoester derivatives **36a–m** in DME and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) as an organocatalyst to produce the substituted 2-furanone derivatives **37a–m** in 41–92% yield (Scheme 25).⁷⁸

Another organocatalyzed synthesis of substituted 2-furanones was achieved by Reitel *et al.* As the chiral compound **39** was used as a catalyst in the reaction between ethyl-3-oxo-3-phenyl-propaneperoxoate (**38**) and **1b** to yield 5-((ethylperoxy)-methyl)-3,4,5-triphenylfuran-2(5H)-one (**40**) in 60% yield (Scheme 26).⁷⁹

In 2018, Matsuda *et al.* succeeded in synthesizing trisubstituted 2-furanones by using another convenient method. The reaction between formamides **41** and **1b** (10–20 equiv.) was performed for refluxing chlorobenzene, which was catalyzed with silver trifluoromethane-sulfonate (AgTOf) to give diphenylfuranone derivatives **42a–d** in 31–98% yield (Scheme 27).⁸⁰


The suggested mechanism is illustrated as follows: diphenylcyclopropane underwent ring-opening [3 + 2] annulation and the corresponding formamides **42** were formed (Scheme 28). The carbonyl oxygen atom of **1** was coordinated to Ag⁺ to form **A**, C-1 atom of **A** was attacked by the carbonyl oxygen atom of **41** and then the intermediate **B** was obtained. The final step afforded the furane ring *via* ring expansion and regenerated Ag⁺.⁸⁰

Ren *et al.* in 2018, used silver catalysis to afford other furanone derivatives **42a–m** in 28–92% yield, *via* the reaction between cyclopropanone derivatives **1b,j,o–s** and formamides **41** using silver hexafluoroantimonate(v) AgSbF₆ at 80 °C (Scheme 29).⁸¹




Triphenylphosphine (TPP) mediated the reaction developed by Nguyen *et al.* The reaction occurred between the substituted cyclopropanones **1t** in methanol and TPP at 23 °C to yield butenolides **44a–l** in 36–91% yield *via* the formation of triphenylphosphine ylidene intermediate **43** (Scheme 30).⁷¹

γ -Alkenylbutenolide **47** was synthesized by Bai *et al.* *via* cycloaddition reaction between **1b** and enones **45** at room






Scheme 20 Mechanism of the formation of 3-pyrrolinone derivatives 27a-c.


Scheme 21 Synthesis of *N*-pyridymethyl-pyrrol-3-ones 29a and 29b. Reagents and conditions: A = $[\text{Rh}(\text{OAc})(\text{cod})]_2$ (5 mol%), toluene, 140 °C, 12 h.Scheme 22 Synthesis of pyrrol-2-ones 29c,d. Reagents and conditions: A = $[\text{Rh}(\text{OAc})(\text{cod})]_2$ (10 mol%), toluene 2 ml, 140 °C, 1 h.Scheme 23 Palladium-catalyzed synthesis of maleimides 31a-q. Reagents and conditions: A = $\text{Pd}(\text{OAc})_2$ (15 mol%), KOAc (4 equiv.), Bu_4NBr (1 equiv.), $\text{K}_2\text{S}_2\text{O}_8$ (1.5 equiv.), DMF (0.25 M), 120 °C, 1 h.

temperature in the presence of Ni complex 46 as an organic catalyst (Scheme 31).⁸²


The plausible reaction mechanism was proposed in Scheme 32, as intermediate A was obtained by oxidative addition of 1b to Ni^0 . The intermediate A was then thought to migrate into the $\text{C}=\text{O}$ bond *via* intermediate B and enantioselective $\text{Ni}-\text{C}(\text{acyl})$ migratory insertion occurred (path a). The Ni^2 -allyl intermediate C that resulted reductively removed the final product 47, allowing the catalyst to be regenerated. Alternatively, or even more probable, the intermediate B may go through a concerted 4,1-insertion of the Ni -acyl into the enone, yielding an 8-membered nickelacycle D that was a direct predecessor of C (path b). The olefin unit is involved in both pathways to allow for allyl stabilization (Scheme 32).⁸²



Scheme 24 Synthesis and mechanism describing the formation of compounds 35a-f. Reagents and conditions: A = MeOH, rt.





Scheme 25 Synthesis of substituted butanolides 37a-m. Reagents and conditions: A = DBU (20 mol%), DME, rt, 24 h.

Scheme 26 Synthesis of 5-((ethylperoxy)-methyl)-3,4,5-triphenylfuran-2(5H)-one (40). Reagents and conditions: A = CH₂Cl₂, 50% aq KOH (1 equiv.), rt, 1 h, 39 (20 mol%).




2.3.2.3. Synthesis of various classes of five-membered rings with one heteroatom. 2-(5-(2,3-Diaryl)cycloprop-2-en-1-ylidene)-2,5-dihydrothiophen-2-yl)malononitriles 49a,b were obtained

in good yield *via* the reaction of 2-(thiophen-2-yl)malononitrile (48) with diarylcyclopropenones 1u,v in refluxing acetic anhydride (Scheme 33).⁸³

Scheme 28 Mechanism of the formation of diphenylfuranones 42a-d. Reagents and conditions: A = AgTOF (10 mol%), C₆H₅Cl, 130 °C, 2 h.

Scheme 27 Synthesis of substituted diphenylfuranones 42a-d. Reagents and conditions: A = AgTOF (10 mol%), PhCl, 130 °C, 2 h.



Scheme 29 Synthesis of compound 42a-m. Reagent and conditions: A = AgSbF_6 (10 mol%), 80 °C, 20 h.Scheme 30 TPP mediated the synthesis of 44a-l. Reagents and conditions: A = PPh_3 (5 mol%), MeOH, 23 °C.Scheme 31 Synthesis of γ -alkenylbutenolide 47 from enones 45 and 1b. Reagents and conditions: A = $\text{Ni}(\text{cod})_2$ (1–2 mol%), $\text{L}^* 46$ (2–3 mol%), toluene, rt, 1–24 h.



2.3.2.4. Spirocyclic heterocycles with one heteroatom. Spirocyclic heterocycles with one heteroatom were established *via* the reaction of cyclopropenones **1** with organic compounds having various heteroatoms.

Matsuda and Sakurai used gold catalysis in their reaction between 4-methyl-N-(3-methylbut-2-en-1-yl)-N-(prop-2-yn-1-yl)benzene-sulfonamide (**50**) and cyclopropenone **1j,s,w,x**. The former reaction was carried out in DCM, at room temperature, and in the presence of $(\text{IPr})\text{AuNTf}_2$ to give the spiro compounds **51a-d** in 85–97% yield, as shown in Scheme 34.⁸⁴

Xu *et al.* succeeded to synthesize the spiro heterocyclic compounds **53** and **54** from the reaction between cyclopropenone derivatives **1b,o,p,q,y** and isatin **52** (Scheme 35). The reaction was carried out in toluene and 4-dimethylaminopyridine (DMAP) as a catalyst at 50 °C. The two isomers of spiro furano-indolinone **53a-h** and **54a-h** were formed in 5–50% and 10–99% yield, respectively.⁸⁵

Previously, Cunha and Rocha had reported that 1,6-diisopropyl-2,7-diphenyl-4-oxaspiro-[2.4]hepta-1,6-dien-5-one (**55**) was synthesized in 48% yield, *via* refluxing 2-isopropyl-3-phenyl-cycloprop-2-en-1-one (**1z**) in dioxane in the presence of copper chloride (CuCl) for 12 h, as shown in Scheme 36.⁸⁶

In 2014, Rivero *et al.* have reported that the reaction between substituted cyclopropenones **1b,a'** and trisubstituted cyclopropanes **56a-j** in DCM catalyzed by scandium(III) trifluoromethane-sulfonate $\text{Sc}(\text{OTf})_3$ for 4 h gave the spiro heterocyclic compounds **57a-j** (Scheme 37).⁸⁷


Scheme 32 Mechanism describing the synthesis of γ -alkenylbutenolide 47.Scheme 33 Reaction of 2-(thiophen-2-yl)malononitrile (48) with diarylcyclopropenones 1u,v. Reagents and conditions: A = Ac_2O , reflux.


1,2-Bis(2-methoxy-5-methylphenyl)-6,7-diphenyl-4-oxaspiro[2.4]hepta-1,6-dien-5-one (59) was obtained in 70% yield *via* the reaction of two derivatives of cyclopropenones **1b** and **1b'** together with compound 2 in the presence of CuBr as a catalyst. One of these cyclopropenones was synthesized from the

reaction of **58** with 2 in $\text{AlCl}_3/\text{CH}_2\text{Cl}_2$ at $-20\text{ }^\circ\text{C}$ (Scheme 38).⁸⁸ The reaction was due to the formation of **1c'** (Scheme 38).⁸⁸


Then, we proceeded to the formation of fused compounds such as indoles **61a–g**, which were obtained in 45–85% yield, *via* the reaction between *N*-nitrosoanilines **60** and cyclopropenones **1b,r,d'** in the presence of $[\text{RhCp}^*(\text{OAc})_2]$ and AgNTf_2 in DCE at $120\text{ }^\circ\text{C}$ for 24 h (Scheme 39).⁸⁹

The interesting approach to prepare compounds containing indole moieties is outlined in Scheme 40. The strategy started with treatment compound **62** with trimethylsilyl trifluoromethanesulfonate (TMS-OTf), Et_3N and TiCl_4 (method A). Oxidation of **63** with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) in CHCl_3 (method B) afforded compound **64**. Rearrangement of **64** using methyl triflate (MeOTf) (method C) gave the corresponding compound **65** (Scheme 40).⁹⁰ Upon heating **65** with cyclopropenone (**1a**) in MeCN (method D), the reaction gave the target products **66a** in 89% yield together with its

Scheme 34 (IPr)AuNTf₂-catalyzed synthesis of compounds 51a–d. Reagents and conditions: A = (IPr)AuNTf₂ (2 mol), DCM, rt.



Scheme 35 Synthesis of spiro furano-indolinone 53a-h and 54a-h. Reagents and conditions: A = DMAP (20 mol%), toluene, 50 °C, 6 h.




Scheme 36 Synthesis of the spiro compound 55. Reagents and conditions: A = dioxane, CuCl, 12 h.


diastereomer **66b** (Scheme 40). The two diastereomers **66a,b** were used as precursors in the synthesis of (\pm) Aspergilline A.⁹⁰

Cunha *et al.* reported on a direct path to obtain pyrrolizidine **68** and indolizidine **69** by the reaction of **67** with **1b** in refluxing toluene (Scheme 41).⁹¹


Recently, in 2021, Yao *et al.* reacted dimethyl 2-(1,11-diphenylundeca-1,3,8,10-tetraen-6-yl)malonate (**70**) with cyclopropane derivatives **1j,o,p,r** under the stream of O₂ to give the




Scheme 37 Sc(OTf)₃ mediated the synthesis of 57a-j. Reagents and conditions: A = Sc(OTf)₃ (10 mol%), DCM, 25 °C 4 h.




Scheme 38 Synthesis of 1,2-bis(2-methoxy-5-methylphenyl)-6,7-diphenyl-4-oxaspiro[2.4]hepta-1,6-dien-5-one (59). Reagents and conditions: A = (i) AlCl₃, CH₂Cl₂, -20 °C, 45 min; (ii) H₂O; B = CuBr, CH₂Cl₂, 70 °C, 12 h.




Scheme 39 Synthesis of indoles 61a–g. Reagents and conditions: A = $[\text{RhCp}^*(\text{OAc})_2]$, AgNTf_2 , DCE, $120\text{ }^\circ\text{C}$, 24 h.



Scheme 40 Synthesis of compounds 66a, b. Reagents and conditions: A = TMS-OTf , Et_3N , DCM, TiCl_4 , $0\text{--}35\text{ }^\circ\text{C}$; B = CHCl_3 , DDQ, H_2O , $70\text{ }^\circ\text{C}$; C = MeOTf , CH_2Cl_2 ; D = MeCN , $50\text{ }^\circ\text{C}$.





Scheme 41 Synthesis of pyrrolizidine 68 and indolizidine 69. Reagents and conditions: A = toluene, reflux.




Scheme 42 Synthesis of benzo[b]furane derivatives 71a–d. Reagents and conditions: A = O_2 , MeCN, reflux, 12 h.



Scheme 43 Synthesis of spiro oxazoles 73a-g. Reagents and conditions: A = K₂CO₃ (0.8 mmol), HFIP (2 ml), 50 °C, 12 h.

Scheme 44 Mechanism illustrating the formation of spiro oxazoles 73a-g.



Scheme 45 Synthesis of spiro compound 75. Reagents and conditions: A = n pentane, -40 °C, overnight.

benzo[b]furan derivatives 71a-d in 69–82% yield, as shown in Scheme 42.⁹²

2.3.2.5. Five-membered rings with two heteroatoms. Using hexafluoroisopropanol (HFIP) and potassium carbonate (K₂CO₃) in the reaction between cyclopropenone derivatives 1b, e', f' and α -halohydroxamate 72 afforded the spiro oxazoles 73a-g in 68–86% yield (Scheme 43).⁹³

The mechanism described the formation of 73a-g as an initiation step using K₂CO₃, and 72 was converted *in situ* into azaoxyallyl cation intermediate A. Thereafter, the azaoxyallyl cation A gave the zwitterionic intermediate B after addition of

carbonyl oxygen in the cyclopropenone 1. Finally, the spirocyclic oxazoles 73 were obtained by intramolecular nucleophilic addition of B, as illustrated in Scheme 44.⁹³

When (E)-di-*t*-butyl(1-(dimethylphosphino)-2-phenylvinyl) aluminum (74) reacted with 1b in *n*-pentane at -40 °C, the spiro product 75 was produced in 51% yield (Scheme 45).⁹⁴

In 2020, Wu *et al.* had reported that the reaction between diaryl cyclopropenones 1j, g'-o' and elemental sulfur in dimethylformamide (DMF) at room temperature for 12 h provided disubstituted dithiolone derivatives 76a-j in 25–98% yield, as shown in Scheme 46.⁹⁵


[1,2]Dithiolo[5,1-*e*][1,2]dithiole derivatives 78a-d were successfully synthesized in 40–51% yield, by refluxing two moles of cyclopropenones 1 in DCM at 70 °C in the presence of CuBr, the spironolactone intermediate 77 was formed and then reacted with elemental sulfur in DMF at 50 °C for 5 h to give the targeted compounds (Scheme 47).⁹⁵

Similarly, Wu *et al.* also succeeded to obtain diselenolone derivatives 79a-f in 76–85% yield (Scheme 48), *via* the reaction between cyclopropenones 1 and elemental selenium in dimethylsulfoxide (DMSO) at 120 °C under N₂ flow for 12 h.⁹⁵


2.3.2.6. Five-membered rings with three heteroatoms. When 4-phenyl-1-tosyl-1*H*-1,2,3-triazole (80) reacted with 2,3-diphenylcyclopropenone (1b) in DCE for 2.5 h, 2,2'-(2,3-diphenylcycloprop-2-ene-1,1-diyl)bis(4-phenyl-2*H*-1,2,3-triazole) (81) was obtained in 97% yield (Scheme 49).⁹⁶

Hassan *et al.* reported that a series of thiadiazoles 83a-p were obtained in 69–84% yield, *via* a catalyst-free reaction between 1b and alkenylidene hydrazine carbothioamides 82a-p in dry ethanol.⁹⁷


The reaction mechanism was explained by the attack of the azomethine (CH=N) nitrogen atom of compound 82 to the carbonyl group of 1b. The mechanism of the reaction was explained by the formation of intermediate B *via* a spontaneous



Scheme 46 Synthesis of dithiolone derivatives **76a–j**. Reagents and conditions: A = KF, DMF, air, rt, 12 h.




Scheme 47 Synthesis of [1,2]dithiolo[5,1-e][1,2]dithioles **78a–d**. Reagents and conditions: A = CuBr (5 mol%), DCM, 75 °C, N₂, 12 h, B = KF (2 equiv.), S₈, DMF, 50 °C, air, 5 h.



Scheme 48 Synthesis of diselenolone derivatives **79a–f**. Reagents and conditions: A = DMSO, N₂, 120 °C, 12 h.

intramolecular nucleophilic attack of the sulfur atom lone pair on the CH=N group. That was followed by cyclization, *via* the formation of intermediate **B**, which rearranged to the final product **83** (Scheme 50 and Table 1).⁹⁷




Scheme 49 Synthesis of 2,2'-(2,3-diphenylcycloprop-2-ene-1,1-diy)bis(4-phenyl-2H-1,2,3-triazole) (**81**). Reagents and conditions: A = DCE, 80 °C, 2.5 h.

Aly *et al.* reported that the reaction mixture of **1b** and ylidene-N-phenylhydrazine-carbothioamides **84a–e** in glacial acetic acid at room temperature afforded 2,5,6,7-tetrasubstituted-pyrrolo [2,1-*b*][1,3,5-oxadiazolyl]-2-amines **85a–e** in 60–76% yield (Scheme 51).⁹⁸

The suggested mechanism is explained as follows: the afforded product structures supported the formal [2 + 3] cyclo-addition pathway proposed by Eicher to generate the adducts (**A**). This was followed by cyclization of intermediate (**B**) with aromatization of the pyrrole ring. Eventually, intermediate **B** lost a molecule of hydrogen sulfide and the final product **85** was formed, as shown in Scheme 52.⁹⁸







Scheme 50 Synthesis and mechanism of thiadiazoles 83a–p. Reagents and conditions: A = EtOH, 4–8 h.

Table 1 Substituents and the yield of thiadiazoles 83a–p


	R	R ¹	Yield (%)
a	Ph-	C ₆ H ₅ –CH=CH (E)	82
b	Bn-	C ₆ H ₅ –CH=CH (E)	75
c	CH ₂ =CH–CH ₂ –	C ₆ H ₅ –CH=CH (E)	70
d	Ph-	2- MeO–C ₆ H ₄ –CH=CH (E)	72
e	Bn-	2- MeO–C ₆ H ₄ –CH=CH (E)	76
f	CH ₂ =CH–CH ₂ –	2- MeO–C ₆ H ₄ –CH=CH (E)	68
g	Ph-	C ₆ H ₅ –CH=C–Me ₂ (E)	84
h	Ph-	Me–(CH ₂) ₂ –CH=CH– (E)	83
i	Bn-	Me–(CH ₂) ₂ –CH=CH– (E)	65
j	CH ₂ =CH–CH ₂ –	Me–(CH ₂) ₂ –CH=CH– (E)	71
k	Ph-	Me–CH=CH– (E)	75
l	Bn-	Me–CH=CH– (E)	71
m	CH ₂ =CH–CH ₂ –	Me–CH=CH– (E)	67
n	Ph-	Me ₂ –CH–	76
o	Bn-	Me ₂ –CH–	74
p	CH ₂ =CH–CH ₂ –	Me ₂ –CH–	69



Scheme 51 Synthesis of pyrrolo[2,1-b](1,3,5-oxadiazolyl)-2-amines 85a–e. Reagents and conditions: A = AcOH, 4–8 h.



Scheme 52 Mechanism of the formation of pyrrolo[2,1-b](1,3,5-oxadiazolyl)-2-amines 85a–e.



Scheme 53 Synthesis of 4-isopropyl-3,6-diphenyl-2H-pyran-2-one (**87a**). Reagents and conditions: A = benzene, reflux, 5 h.

2.3.3. Synthesis of six-membered rings

2.3.3.1. Six-membered rings with one heteroatom. On reacting 2-isopropyl-3-phenylcycloprop-2-en-1-one (**1z**) with 1-phenyl-2-(triphenyl-phosphoranylidene)ethanone (**86**) in benzene for 5 h, 4-isopropyl-3,6-diphenyl-2H-pyran-2-one (**87a**) was obtained in 35% yield (Scheme 53).⁸⁶

Furthermore, Zhou *et al.* reported on the synthesis of a series of 2-pyranone derivatives **87b–j** in 52–93% yield, that was achieved by the reaction between β -ketosulfoxonium ylides **1b** in MeCN at 100 °C for 12 h, and dichloro(pentamethyl-cyclopentadienyl)-rhodium(III) dimers (Cp^*RhCl_2)₂ (Scheme 54).⁹⁹

The reaction between 2-aryl(cycloprop-2-enones **1** and substituted 2-indolones **89** at 25 °C in acetonitrile for 1 h and DABCO was used as an organic catalyst to form compound **90a–n** in 67–92% yield (Scheme 55).¹⁰⁰

The mechanism was described as follows. First, DABCO subtracts one proton from two to form the nucleophile **A**, on the


less sterically hindered side. Thereafter, 1,4-addition of the C=C bond of **1** was initiated *via* nucleophile **A**, yielding the enolate intermediate **B**. The intermediate **B** was then added intermolecularly to another molecule of **1** to form intermediate **C**, which was converted to intermediate **D** by a concerted process after the ring-opening process and intramolecular nucleophilic addition; after that, it was passed through another ring-opening step to produce intermediate **E**, which after protonation gave the final product **90**, and the catalyst was regenerated (Scheme 56).¹⁰⁰

A series of substituted diaryl spiro[cycloprop[2]ene-1,9'-xanthene] derivatives **92a–j** were successfully synthesized in 40–80% yield, *via* the reaction between compound **91a–d** (2.5 equiv.) and diaryl cyclopropenones **1** (1 equiv.) (Scheme 57). The reaction was performed in MeCN using cesium fluoride (CsF) as a catalyst at 30 °C for 24 h.¹⁰¹


The rhodium-catalyzed reaction occurred between substituted *N*-nitrosoanilines **60** and **1b** in the presence of AgBF_4 and DCE at 100 °C for 12 h. The quinoline-4-one derivatives **93a–f** were produced in 46–70% yield, as shown in Scheme 58.¹⁰²

Moreover, in 2020, Shi *et al.* reacted with **1b** with *N*-nitrosoanilines **60** using $[\text{RhCp}^*(\text{OAc})_2]$, $[\text{Rh}(\text{COD})\text{Cl}]_2$ and AgBF_6 in DCM and yielded 4-quinolones **93a–h** in 51–85% yield (Scheme 59).⁹⁹

(3*S*,4*a**S*,9*b**S*)-Diethyl-3-methyl-4-oxo-4*a*-phenyl-1,4,4*a*,9*b*-tetrahydrobenzofuro[3,2-*b*]pyridine-2,2(3*H*)-dicarboxylates **96a–**




Scheme 54 Synthesis of 2-pyranone derivatives **87b–j**. Reagents and conditions: A = $(\text{Cp}^*\text{RhCl}_2)_2$, NaOAc, MeCN, 100 °C, 12 h.




Scheme 55 Synthesis of compound **90a–n**. Reagents and conditions: A = DABCO (20 mol%), MeCN, 25 °C, 1 h.





Scheme 56 Mechanism describing the formation of 90a-n.




Scheme 57 Synthesis of substituted diaryl spiro[cycloprop[2]ene-1,9'-xanthene] derivatives 92a-j. Reagents and conditions: A = CsF, MeCN, 30 °C, 24 h.

d were synthesized *via* the reaction of (*E*)-diethyl 2-((2-hydroxybenzylidene)amino)malonate derivatives (**94**) with **1j** at room temperature in the presence of the catalyst **95** (Scheme 60).¹⁰³


In 2020, it was reported that phenanthridine derivatives **98a-d** were synthesized in 67–82% yield, *via* refluxing [1,1'-biphenyl]-2-amine (**97**) with substituted cyclopropenones **1** in DCM for 48 h and in the presence of [Ru(*p*-cymene)Cl₂]₂ and Ag₂CO₃ (Scheme 61).¹⁰⁴

Substituted pyrido[3,4-*b*]indoles **99a-j** could react easily with cyclopropenone derivatives **1** at room temperature in absolute ethanol for 5–50 h. The reaction yielded compounds **100a-j** in 39–93% yield (Scheme 62).¹⁰⁵


Refluxing diarylcyclopropenones **1** in pyridine at 100 °C in the presence of cupric acetate (Cu(OAc)₂) afforded biindolizines **101a-b**, as shown in Scheme 63.¹⁰⁶



Scheme 58 Rhodium(III)-catalyzed synthesis of quinoline-4-one derivatives **93a–f**. Reagents and conditions: A = $(Cp^*\text{RhCl}_2)_2$ (2 mol%), AgBF_4 (1.2 equiv.), NaF , DCE, $100\text{ }^\circ\text{C}$, 12 h.



Scheme 59 Synthesis of 4-quinolones **93a–h**. Reagents and conditions: A = $[\text{RhCp}^*(\text{OAc})_2]$ (0.5 mol%), $[\text{Rh}(\text{COD})\text{Cl}]_2$, AgBF_6 (0.02 mmol), DCM, $120\text{ }^\circ\text{C}$, 36 h.



Scheme 60 Synthesis of compounds **95a–d**. Reagents and conditions: A = mesitylene, rt, cat 96.


7-Methyl-1-phenyl-3,4-dihydrobenzo[*b*][1,7]naphthyridine (**102**) easily reacted with cyclopropenones **1** without using any catalyst at room temperature in EtOH to afford 2,3-disubstituted-9-methyl-12*b*-phenyl-5,6-dihydrobenzo[*b*]pyrrolo[1,2-*h*][1,7]naphthyridin-1(12*b*H)-ones **103a–b** in 45–56% yield, as shown in Scheme 64.¹⁰⁵

Carbonylation of symmetrical cyclopropenones **1** with alkynes **8c–e** occurred in refluxing toluene for 20 h. The reaction


was mediated by triruthenium dodecacarbonyl $\text{Ru}_3(\text{CO})_{12}$ and yielded pyranopyrandiones **104a–c** in 54–82% yield (Scheme 65).¹⁰⁷

Zhu *et al.* in 2021, have developed a palladium-catalyzed three-component reaction of substituted iodochromen-4-ones **105a–h**, (1*R*,4*S*)-bicyclo[2.2.1]hept-2-ene (**106**) and **1b** in fluorobenzene in the presence of tris-(4-trifluoromethyl-phenyl) phosphine $\text{P}(4\text{CF}_3\text{-C}_6\text{H}_4)_3$ at $100\text{ }^\circ\text{C}$ for 24 h.







Scheme 61 Synthesis of phenanthridine derivatives **98a–d**. *Reagents and conditions:* A = $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$, Ag_2CO_3 , H_3BO_3 , DCM, reflux, 48 h.



Scheme 62 Synthesis of **100a–j**. *Reagents and conditions:* A = EtOH, rt, 5–50 h.



Scheme 63 Synthesis of biindolizines **101a,b**. *Reagents and conditions:* A = pyridine, $\text{Cu}(\text{OAc})_2$, 15 min.



Scheme 64 Synthesis of the fused heterocyclic compounds **103a,b**. *Reagents and conditions:* A = EtOH, rt.

(*6aR,7S,10R,10aS*)-*6a*,*7,8,9,10,10a*-Hexahydro-7,10-methanoindeno[2,1-*b*]chromene-6,11-diones **107a–h** were obtained in 65–70% yield (Scheme 66).¹⁰⁸




In 2021, Chen *et al.* successfully synthesized a series of tetrasubstituted pyrano[2,3-*b*]indol-2(9*H*)-ones **109a–l** in 65–91% yield *via* the reaction between substituted isatines **40** and diaryl cyclopropenones **1** at 110 °C in toluene and in the presence of lanthanide amides $[(\text{Me}_3\text{Si})_2\text{N}]_3\text{La}(\mu\text{-Cl})\text{Li}(\text{THF})_3$ and ligand **108** as a catalyst (Scheme 67).¹⁰⁹

2.3.3.2. Six-membered rings with two heteroatoms. (5*S*,6*R*)-5,6-Diphenyl-2-((*R*)-1-phenylethyl)amino)-5,6-dihydropyrimidin-4(1*H*)-one (**113**) was successfully synthesized by Ahmed *et al.* *via* the reaction between (*R*)-1-(1-phenylethyl)guanidine (**112**) and **1b**. The reaction was performed at room temperature in benzene and EtOH (Scheme 68).¹¹⁰

A catalyst-free reaction between *N*-carbamoyl sulfilimines **114a–f** and **1b** in toluene at 110 °C for 13 h afforded the two isomers of diphenyl pyrimidinediones **115a–f** in 51–94% yields, as shown in Scheme 69.¹¹¹


Aly *et al.* in 2016, successfully synthesized pyrimidines **117a–f** in 67–87% yield, *via* the reaction between **116a–f** and **1b** in EtOH and Et₃N (Scheme 70).¹¹²


The proposed reaction mechanism is illustrated in Scheme 71; the carbonyl group of cyclopropenone was attacked by the hydrazine nitrogen atom yielding intermediate **B**, following which an amidine-like reaction of N-3 to carbonyl may occur to obtain salt **C**. Nucleophilic addition to positively charged

Scheme 65 Synthesis of pyranopyrandiones 102a-c. Reagents and conditions: A = toluene, Ru₃(CO)₁₂, Et₃N, 150 °C, 20 h.Scheme 66 Synthesis of compounds 107a-h. Reagents and conditions: A = PdCl₂, P(4-CF₃-C₆H₄)₃ (20 mol%), Cs₂CO₃, C₆H₅F, 100 °C, 24 h.Scheme 67 Synthesis of tetrasubstituted pyrano[2,3-b]indol-2(9H)-ones (109a-l). Reagents and conditions: A = [(Me₃Si)₂N]₃La(μ-Cl)Li(THF)₃, toluene, 110 °C, 2.5 h, (108), HOP(OEt)₂.



nitrogen *via* ring opening followed by proton transfer would afford **D**, and the final product **117** was obtained by losing ammonia from **D** (Scheme 71).¹¹²

Aly *et al.* also reported that the reaction between (E)-N'-aryl-N-(phenylcarbamothioyl)-benzimidamides **118** and **1b** in EtOH produced substituted 3-aryl-2,5,6-triphenylpyrimidin-4(3*H*)-ones **119a-e**, as shown in Scheme 72.¹¹³

Scheme 68 Synthesis of 5,6-dihydropyrimidine 113. Reagents and conditions: A = conc. HCl, 1,4 dioxane, 20 °C; B = NH₂CN, pH 8-9, H₂O, reflux, 5 h; C = passed through Amberlite IRA-401 (hydroxide form); D = benzene : EtOH (1 : 1), rt, 18 h.



Scheme 69 Synthesis of diphenyl pyrimidinediones 115a-f. Reagents and conditions: A = toluene, 110 °C, 13 h.

Scheme 70 Synthesis of pyrimidines 117a-f. Reagents and conditions: A = EtOH, Et₃N, reflux 6–10 h.

Scheme 71 Mechanism of the formation of pyrimidines 117a-f.


Mohan and Jose, in 2017, have reported that reaction between (*E*)-disubstituted diazene-1,2-dicarboxylates 120a-j and diarylcyclopropane 1 in DCM at room temperature. The reaction was catalyzed by PPh₃ to afford substituted 1,3-oxazin-6-one 121a-j in 50–70% yield (Scheme 73).¹¹⁴

The reaction was initiated *via* PPh₃ attack on 120 to give A and Huisgen zwitterion's nucleophilic attack on


cyclopropane (1) in step B; the intermediate (C) was formed followed by internal cyclization of D to generate 1,3-oxazin-6-ones 121a-j, as illustrated in Scheme 74.¹¹⁴

Via the reaction of equal amounts of amide derivatives 124a-i with 2,3-diphenylcyclopropanes (1) in DCE using CsOAc as a catalyst, the substituted 1,3-oxazin-6-ones 121k-q were obtained in 50–93% yield (Scheme 75).¹¹⁵






Scheme 72 Synthesis of substituted 3-aryl-2,5,6-triphenylpyrimidin-4(3H)-ones (119a–e). Reagents and conditions: A = EtOH, reflux, 10–16 h.



Scheme 73 Synthesis of substituted 1,3-oxazin-6-ones 121a–j. Reagents and conditions: A = PPh₃ (1 equiv.), DCM, rt, 15 min.



Scheme 74 Mechanism of the formation of substituted 1,3-oxazin-6-ones 121a–j.


Recently, it has been reported that the oxime derivatives **123a–g** reacted with 2,3-diphenyl-cyclopropenones (**1**) to give substituted 1,3-oxazine-4-ones **124a–g** in 52–91% yield (Scheme 76). The reaction was performed at 80 °C in cyclohexane for 18 h, and Ag₂O was also used as a catalyst.¹¹⁶

The reaction mechanism is illustrated in Scheme 77. At first, the resonance structure of cyclopropenone was nucleophilically attacked by oximes, yielding intermediate **A**. Subsequently the intermediate **A** was fragmented into intermediates **B** and **C**. Accordingly, the intermediate **D** was obtained *via* a [4 + 2] cycloaddition between **B** and **C**. Then, a reaction occurred between **D** and another cyclopropenone to generate intermediate **E**, and the intermediate **F** was subjected to a rearrangement/protonation process to form compound **124**.¹¹⁶

In 2021, Sizhan, *et al.* successfully synthesized substituted 1,3-oxazin-6-ones **121r–y** in 55–93% yield, *via* the reaction between α -halogenated hydroxamates **125a–g** and **1b** in DCM and Et₃N (Scheme 78).¹¹⁷


2,4,5-Triphenyl-6H-1,3-oxazin-6-one (**121z**) was obtained in 95% yield by reacting *N*-(pivaloyloxy)benzamide (**126**) with 2,3-diphenylcyclopropenone (**1**) at 60 °C in THF using K₂CO₃. The mechanism of the reaction was explained by the nucleophilic attack of the nitrogen atom of amide **26** on **1b** and intermediate **B** was then formed in the presence of base. During the ring opening, **B** lost a pivalate anion to give ketene **C**, which rearranged to form the final structure (**121z**), *via* a 6 π electrocyclization process, as shown in Scheme 79.¹¹⁸

In 2019, substituted thiazinane derivatives **128a–e** were successfully synthesized by Hassan *et al.* in 2019. It was achieved during the reaction of hydrazinecarbothioamide derivatives **127a–e** with **1b** in refluxing EtOH (Scheme 80). The




Scheme 75 Scandium-catalyzed synthesis of substituted 1,3-oxazin-6-ones 121k–q. Reagent and conditions: A = CsOAc (1 equiv.), DCE, rt, 6 h.






Scheme 76 Synthesis of 1,3-oxazine-4-ones 124a-g. Reagents and conditions: A = Ag₂O, cyclohexane, 80 °C, 18 h.



Scheme 77 Mechanism describing the formation of 1,3-oxazine-4-ones 124a-g.



Scheme 78 Synthesis of substituted 1,3-oxazin-6-ones 123r-y. Reagents and conditions: A = DCM, reflux, Et₃N, 2 h.

reaction afforded compounds 128a-e, as well as a side product 129.¹¹⁹



The reaction mechanism could be simply described as the conjugate double bond of 1b was attacked by the sulfur atom generating intermediate A. The intermediate B was formed via the intramolecular nucleophilic attack of N4-H on C=O, which rearranged to generate 128. However, the carbonyl group of 1

was attacked by N4-H, resulting in the formation of the product 128 via intermediate D (Scheme 81).¹¹⁹


When pyrazolylthioureas 130a-c were subjected to 1b, the reaction proceeded to give compounds 131a-c in 75-90% yield (Scheme 82). The reaction was performed in ethanol for 4-7 h in the presence of DDQ as the oxidizing agent.¹²⁰

Moreover, in 2021, Shi *et al.* succeeded to synthesize the heterocyclic substituted 2-phenyl-10'H-spiro[indene-1,12'-isoindolo[1,2-b]quinazolin]-10'-ones 133a-j in 58-87% yield by reacting 2-phenylquinazolin-4-ones 132a-j with cyclopropenones 1 in refluxing DCE for 24 h, and a (cymene)-ruthenium dichloride dimer [Ru(p-cymene)Cl]₂ was used as a metal catalyst (Scheme 83).¹²¹

Shi *et al.* also reacted substituted 2-phenylquinazolin-4-ones 134a-f with 1b to obtain compounds 135a-f in 62-82% yield. The reaction was performed in trifluoroethanol (TFE) at 110 °C for 48 h. Subsequently, treatment of compound 135a-f with trifluoroacetic acid (TFA) at 140 °C and in the presence of Rh [Cp*(OAc)] and Ag₂O yielded compounds 136a-f (Scheme 84).¹²¹


Scheme 79 Synthesis of 2,4,5-triphenyl-6H-1,3-oxazin-6-one (146z). Reagents and conditions: A = K_2CO_3 (0.5 equiv.), THF, 60 °C, 2 h.

Scheme 80 Synthesis of thiazinanes 128a-e. Reagents and conditions: A = EtOH, reflux.







Scheme 81 Mechanism of the formation of thiazinanes 128a-e.





Scheme 82 Synthesis of thiazines 131a–c. Reagents and conditions: A = DDQ, EtOH, reflux, 4–7 h.

Scheme 83 Synthesis of substituted quinazolin-ones 133a–j. Reagents and conditions: A = $[\text{Ru}(p\text{-cymene})\text{Cl}]_2$, AgSbF_6 , AdCOOH, DCE, reflux, 24 h.Scheme 84 Synthesis of compounds 135a–f and 136a–f. Reagents and conditions: A = $\text{Rh}[\text{Cp}^*(\text{OAc})]_2$, TFE, 110 °C, 48 h; B = $\text{Pd}(\text{OAc})_2$, Ag_2O , TFA, 140 °C, 24 h.Scheme 85 Synthesis of quinazoline derivatives 138a–i. Reagents and conditions: A = $(\text{Cp}^*\text{RhCl})_2$, AgSbF_6 , DCM, reflux, 36 h, O_2 .




Scheme 86 Synthesis of 2-benzyl-3-isopropylquinoxaline (**140**). *Reagents and conditions:* A = Et₂O, 8 h.

Refluxing **137** with 2,3-substituted cyclopropenones **1** in DCM yielded quinazoline derivatives **138a–i** in 44–85% yield. The reaction was catalyzed using (Cp^{*}RhCl₂)₂ and AgSbF₆, as shown in Scheme 85.¹²²

2-Benzyl-3-isopropylquinoxaline (**140**) was synthesized *via* reacting 2-isopropyl-3-phenylcycloprop-2-en-1-one (**1z**) with benzene-1,2-diamine (**139**) in diethyl ether for 8 h and in the presence of 1*H*-pyrazole as a catalyst (Scheme 86).⁸⁶

6,7-(Diphenylpyrrolo[1,2-*b*]pyridazin-5-yl)acetate (**143**) was successfully synthesized in 55% yield *via* a cyclization reaction between pyridazine (**141a**) and **1b** in DCE at the reflux temperature. Subjecting the formed intermediate **142** with acetic anhydride and 4-(*N,N*-dimethylamino)pyridine (DMAP) as a base catalyst gave the final product **143** (Scheme 87).¹²²

The reaction between pyridazine-4,5-dicarbonitrile (**141b**) and **1b** in acetone at 110 °C afforded 1-oxo-2,3-diphenyl-1*H*-pyrazolo[1,2-*a*]pyridazine-6,7-dicarbonitrile (**145**) in 38% yield *via* the formation of intermediate **144** (Scheme 88). In addition to the formation of **145**, [(*E*)-3,4-dicyano-6,7-diphenylpyrrolo[1,2-*b*]pyridazin-5-yl]-2,3-diphenylacrylate (**147**) was also obtained in 47% yield. The formation of **147** can be explained as due to the cyclization process of intermediate **144** into **146**,



Scheme 89 Synthesis of pyridazines **149a,b**. *Reagents and conditions:* A = *p*-xylene, reflux, 138 °C, 20 min.


which tautomerized and reacted with another molecule of **1** to give **147** (Scheme 88).¹²³

Molchanov *et al.* have developed on the synthesis of pyridazines **149a,b** during refluxing a mixture of 6-aryl-3,3-dimethyl-1,5-diazabicyclo[3.1.0]hexanes **148a,b** with 2-methyl-3-phenylcycloprop-2-enone (**1j**) in *p*-xylene for 20 min (Scheme 89).¹²⁴


Aly's group reacted **1b** with hydrazinecarbothioamides **138a–c** in MeOH.¹²⁵

The reaction yielded substituted triazolopyridazines **150a–d**. However, when 1,2-diphenyl-hydrazinecarbothioamide (**129h**) was used, the reaction gave only (*Z*)-1,4,5-triphenyl-6-(phenylimino)-1,6-dihydropyridazine-3(2*H*)-thione (**151**) (Scheme 90).¹²⁵

2.3.3.3. Six-membered rings with three heteroatoms. When cyclopropenone **1** was treated with Et₃OB₄ and CH₂Cl₂ *in situ*, the intermediate **152** was suggested to be formed (Scheme 91). Accordingly, upon addition of Et₂NH *in situ* to the salt **152**, the intermediate **153** was suggested to be formed. Finally, after *in situ* addition of sodium azide (NaN₃) in DMF to **153**, the reaction




Scheme 87 Synthesis of 6,7-diphenylpyrrolo[1,2-*b*]pyridazin-5-yl acetate (**143**). *Reagents and conditions:* A = DCE, reflux, B = Ac₂O, DMAP.



Scheme 88 Synthesis of pyridazines **145** and **147**. *Reagents and conditions:* A = MeCOMe, 110 °C, 48 h.





Scheme 90 Synthesis of pyridazines 150, 151. Reagents and conditions: A = MeOH, reflux, 48 h, B = MeOH, reflux, 6–12 h.

proceeded to yield the triazines 154a–d in 11–45% yield (Scheme 91).¹²⁶

2.3.4. Synthesis of seven-membered-ring heterocycles. Tetrasubstituted-6,7-dihydro-1*H*-silepin-4(5*H*)-ones 156a–h were obtained in 65–89% yield by refluxing 1,1-disubstituted


siletanes 155a–h with cyclopropenone derivatives 1 in toluene at room temperature for 48 h, and Pd(OAc)₂ (Scheme 92).¹²⁷ The reaction was initiated *via* bond cleavage of silacyclobutane (155): the C–Si bond of silacyclobutane was broken through an oxidative addition, which was proposed to give palladacycle A as the first step. Then, after a series of oxidative additions to the C–C bond of cyclopropenone 1b, intermediate C was formed (Scheme 93). The eight-membered palladacycle D was formed *via* the reductive elimination of intermediate C, which might go through a second reductive elimination step to produce the final product 156 and form the palladium (0) catalyst again. The possibility of path b could not be completely excluded, which would first involve the cleavage of the C–C σ -bond of cyclopropenone (1) (Scheme 93).¹²⁷

2.3.5. Synthesis of nine-membered-ring heterocycles.


When compounds 157a–c reacted with 2,3-diphenylcyclopropenone (1b) in absolute ethanol, intermediate 158 was formed followed by the formation of nine-membered-ring bicyclic products 159a–c in 51–62% yield (Scheme 94).¹²⁸

3 Conclusion

In summary, the use of cyclopropane derivatives for the construction of various heterocycles can provide a practical alternative to traditional methods for the preparation of such compounds. Since cyclopropanes have high strain, they easily participated in various reactions and, therefore, in the construction of various organic molecules. In that review, we consider the utility of cyclopropenones in the synthesis of heterocycles, especially those of prospective biologically active compounds. The reactions of cyclopropenones are described as highly regioselective, often stereoselective, and permit the synthesis of a variety of heterocyclic systems, both saturated




Scheme 91 Synthesis of triazines (154a–d). Reagents and conditions: A = Et₃OBF₄, CH₂Cl₂; B = Et₂NH; C = NaN₃, CH₂Cl₂, DMF.




Scheme 92 Synthesis of tetrasubstituted-6,7-dihydro-1*H*-silepin-4(5*H*)-ones 156a–h. Reagents and conditions: A = (1 mol%), Pd(OAc)₂ toluene, rt, 48 h.





Scheme 93 Mechanism of the formation of 6,7-dihydro-1H-silepin-4(5H)-ones 157a-h.



Scheme 94 Synthesis of compounds 159a-c. Reagents and conditions: A = EtOH, reflux.

and unsaturated. The substrates employed in most cases are simple (and often commercially available), making the methods amenable to the rapid construction of diverse collections of compounds.

Author contributions

A. A. Aly (conceptualization, writing, editing, and submitting), Alaa A. Hassan, Sara M. Mostafa (supervision), Asmaa. H. Mohamed (editing of revision), Esraa M. Osman (methodology and writing a draft). A. A. Nayl (editing of revision). All authors have read and agreed to the published version of the manuscript.

Conflicts of interest

The authors declare no conflicts of interest.

References

- 1 K. Komatsu and T. Kitagawa, *Chem. Rev.*, 2003, **103**, 1371–1427, DOI: [10.1021/cr010011q](https://doi.org/10.1021/cr010011q).
- 2 G. Dong, *C-C Bond Activation*, Springer, Berlin, 1st edn, 2014, vol. 346, pp. 1–258.
- 3 W. D. Jones, *Nature*, 1993, **364**, 676–677, DOI: [10.1038/364676a0](https://doi.org/10.1038/364676a0).
- 4 B. Rybtchinski and D. Milstein, *Angew. Chem., Int. Ed.*, 1999, **38**, 870–883, DOI: [10.1002/\(SICI\)1521-3773\(19990401\)38:7<870::AID-ANIE870>3.0.CO;2-3](https://doi.org/10.1002/(SICI)1521-3773(19990401)38:7<870::AID-ANIE870>3.0.CO;2-3).
- 5 C. H. Jun, *Chem. Soc. Rev.*, 2004, **33**, 610–618, DOI: [10.1039/B308864M](https://doi.org/10.1039/B308864M).
- 6 C. Aissa, *Synthesis*, 2011, **21**, 3389–3407, DOI: [10.1055/s-0030-1260233](https://doi.org/10.1055/s-0030-1260233).
- 7 N. Cramer and T. Seiser, *Synlett*, 2011, **4**, 449–460, DOI: [10.1055/s-0030-1259536](https://doi.org/10.1055/s-0030-1259536).
- 8 M. Murakami and T. Matsuda, *Chem. Commun.*, 2011, **47**, 1100–1105, DOI: [10.1039/C0CC02566F](https://doi.org/10.1039/C0CC02566F).
- 9 A. Koroticka, D. Necas and M. Kotora, *Curr. Org. Chem.*, 2012, **16**, 1170–1214, DOI: [10.2174/138527212800564213](https://doi.org/10.2174/138527212800564213).
- 10 A. Dermenci, J. W. Coe and G. Dong, *Org. Chem. Front.*, 2014, **1**, 567–581, DOI: [10.1039/C4QO00053F](https://doi.org/10.1039/C4QO00053F).
- 11 I. Marek, A. Masarwa, P. O. Delaye and M. Leibeling, *Angew. Chem., Int. Ed.*, 2015, **54**, 414–429, DOI: [10.1002/anie.201405067](https://doi.org/10.1002/anie.201405067).
- 12 L. Souillart and N. Cramer, *Chem. Rev.*, 2015, **115**, 9410–9464, DOI: [10.1021/acs.chemrev.5b00138](https://doi.org/10.1021/acs.chemrev.5b00138).
- 13 M. Murakami and N. J. Ishida, *Am. Chem. Soc.*, 2016, **138**, 13759–13769, DOI: [10.1021/jacs.6b01656](https://doi.org/10.1021/jacs.6b01656).
- 14 D. S. Kim, W. J. Park and C.-H. Jun, *Chem. Rev.*, 2017, **117**, 8977–9015, DOI: [10.1021/acs.chemrev.6b00554](https://doi.org/10.1021/acs.chemrev.6b00554).
- 15 P. H. Chen, B. A. Billett, T. Tsukamoto and G. Dong, *ACS Catal.*, 2017, **7**, 1340–1360, DOI: [10.1021/acscatal.6b03210](https://doi.org/10.1021/acscatal.6b03210).



16 Q. Z. Zheng and N. Jiao, *Chem. Soc. Rev.*, 2016, **45**, 4590–4627, DOI: [10.1039/C6CS00107F](https://doi.org/10.1039/C6CS00107F).

17 B. Zhao, T. Rogge, L. Ackermann and Z. Shi, *Chem. Soc. Rev.*, 2021, **50**, 8903–8953, DOI: [10.1039/C9CS00571D](https://doi.org/10.1039/C9CS00571D).

18 R. Vicente, *Chem. Rev.*, 2021, **121**, 162–226, DOI: [10.1021/acs.chemrev.0c00151](https://doi.org/10.1021/acs.chemrev.0c00151).

19 J. Wang, S. A. Blaszczyk, X. Li and W. Tang, *Chem. Rev.*, 2021, **121**, 110–139, DOI: [10.1021/acs.chemrev.0c00160](https://doi.org/10.1021/acs.chemrev.0c00160).

20 M. Murakami and N. Ishida, *Chem. Rev.*, 2021, **121**, 264–299, DOI: [10.1021/acs.chemrev.0c00569](https://doi.org/10.1021/acs.chemrev.0c00569).

21 X. Wang, C. Yu, I. L. Atodiresei and F. W. Patureau, *Org. Lett.*, 2022, **24**, 1127–1131, DOI: [10.1021/acs.orglett.1c04045](https://doi.org/10.1021/acs.orglett.1c04045).

22 H. Huo and Y. Gong, *Org. Biomol. Chem.*, 2022, **20**, 3847–3869, DOI: [10.1039/D1OB02450G](https://doi.org/10.1039/D1OB02450G).

23 L. Qua, Y. Wu, P. Sun, K. Zhang and Z. Liu, *Polymer*, 2017, **114**, 36–43, DOI: [10.1016/j.polymer.2017.02.071](https://doi.org/10.1016/j.polymer.2017.02.071).

24 H. L. Ammon, *J. Am. Chem. Soc.*, 1973, **95**, 7093–7101, DOI: [10.1021/ja00802a033](https://doi.org/10.1021/ja00802a033).

25 R. Breslow, T. Eicher, A. Krebs, R. A. Peterson and J. Posner, *J. Am. Chem. Soc.*, 1965, **87**, 1320–1325, DOI: [10.1021/ja01084a029](https://doi.org/10.1021/ja01084a029).

26 R. Breslow, R. Haynie and J. Mirra, *J. Am. Chem. Soc.*, 1959, **81**(1), 247–248.

27 O. Körner, R. Gleiter and F. Rominger, *Synthesis*, 2009, **2009**, 3259–3262, DOI: [10.1055/s-0029-1216934](https://doi.org/10.1055/s-0029-1216934).

28 K. Köhler and W. E. Piers, *Can. J. Chem.*, 1998, **76**, 1249–1255, DOI: [10.1139/v98-170](https://doi.org/10.1139/v98-170).

29 A. Greenberg, R. P. T. Tomkins, M. Dobrovolny and J. F. Liebman, *J. Am. Chem. Soc.*, 1983, **105**, 6855–6858, DOI: [10.1021/ja00361a018](https://doi.org/10.1021/ja00361a018).

30 A. Kascheres, A. C. Joussef and H. C. Duarte, *Tetrahedron Lett.*, 1983, **24**, 1837–1840, DOI: [10.1016/S0040-4039\(00\)81785-2](https://doi.org/10.1016/S0040-4039(00)81785-2).

31 P. A. Wender, T. J. Paxton and T. J. Williams, *J. Am. Chem. Soc.*, 2006, **128**, 14814–14815, DOI: [10.1021/ja065868p](https://doi.org/10.1021/ja065868p).

32 T. Matsuda and Y. Sakurai, *Eur. J. Org. Chem.*, 2013, **2013**, 4219–4222, DOI: [10.1002/ejoc.201300220](https://doi.org/10.1002/ejoc.201300220).

33 L. Kong, X. Zhou, Y. Xu and X. Li, *Org. Lett.*, 2017, **19**, 3644–3647, DOI: [10.1021/acs.orglett.7b01650](https://doi.org/10.1021/acs.orglett.7b01650).

34 C. Jenny, D. Obrecht and H. Heimgartner, *Tetrahedron Lett.*, 1982, **23**, 3059–3060, DOI: [10.1016/S0040-4039\(00\)87532-2](https://doi.org/10.1016/S0040-4039(00)87532-2).

35 K. T. Potts and J. S. Baum, *Chem. Rev.*, 1974, **74**, 189–213, DOI: [10.1021/cr60288a003](https://doi.org/10.1021/cr60288a003).

36 P. C. Venneri and J. Warkentin, *Can. J. Chem.*, 2000, **78**, 1194–1203, DOI: [10.1139/v00-118](https://doi.org/10.1139/v00-118).

37 R. Noyori, I. Umeda and H. Takaya, *Chem. Lett.*, 1972, **1**, 1189–1190, DOI: [10.1246/cl.1972.1189](https://doi.org/10.1246/cl.1972.1189).

38 A. Rai and L. D. S. Yadav, *Eur. J. Org. Chem.*, 2013, **2013**, 1889–1893, DOI: [10.1002/ejoc.201300059](https://doi.org/10.1002/ejoc.201300059).

39 T. Stach, J. Dräger and P. H. Huy, *Org. Lett.*, 2018, **20**, 2980–2983, DOI: [10.1021/acs.orglett.8b01023](https://doi.org/10.1021/acs.orglett.8b01023).

40 N. K. Urdabayev, A. Poloukhine and V. V. Popik, *Chem. Commun.*, 2006, **4**, 454–456, DOI: [10.1039/B513248G](https://doi.org/10.1039/B513248G).

41 A. G. Lvov, N. A. Milevsky, V. Z. Shirinian and M. M. Krayushkin, *Chem. Heterocycl. Compd.*, 2015, **51**, 933–935, DOI: [10.1007/s10593-015-1800-8](https://doi.org/10.1007/s10593-015-1800-8).

42 C. D. McNitt, H. Cheng, S. Ullrich, V. V. Popik and M. Bjerknes, *J. Am. Chem. Soc.*, 2017, **139**, 14029–14032, DOI: [10.1021/jacs.7b08472](https://doi.org/10.1021/jacs.7b08472).

43 Y. Chiang, A. Kresge and V. Popik, *J. Am. Chem. Soc.*, 1995, **117**, 9165–9171, DOI: [10.1021/ja00141a007](https://doi.org/10.1021/ja00141a007).

44 A. Poloukhine and V. V. Popik, *J. Org. Chem.*, 2003, **68**, 7833–7840, DOI: [10.1021/jo034869m](https://doi.org/10.1021/jo034869m).

45 A. Poloukhine and V. V. Popik, *J. Org. Chem.*, 2006, **71**, 7417–7421, DOI: [10.1021/jo061285m](https://doi.org/10.1021/jo061285m).

46 K. Mishiro, T. Kimura, T. Furuyama and M. Kunishima, *Org. Lett.*, 2019, **21**, 4101–4105, DOI: [10.1021/acs.orglett.9b01280](https://doi.org/10.1021/acs.orglett.9b01280).

47 A. Schuster-Haberhauer, R. Gleiter, O. Körner, A. Leskovar, D. B. Werz, F. R. Fischer and F. Rominger, *Organometallics*, 2008, **27**, 1361–1366, DOI: [10.1021/om701115w](https://doi.org/10.1021/om701115w).

48 T. Okuda, K. Yokose, T. Furumai and H. B. Maruyama, *J. Antibiot.*, 1984, **37**, 718–722, DOI: [10.7164/antibiotics.37.718](https://doi.org/10.7164/antibiotics.37.718).

49 F. Bohlmann, J. Jakupovic, L. Müller and A. Schuster, *Angew. Chem., Int. Ed.*, 1981, **20**, 292–293, DOI: [10.1002/anie.198102921](https://doi.org/10.1002/anie.198102921).

50 T. Okuda, N. Shimma and T. Furumai, *J. Antibiot.*, 1984, **37**, 723–727, DOI: [10.7164/antibiotics.37.723](https://doi.org/10.7164/antibiotics.37.723).

51 H. Kogen, T. Kiho, K. Tago, S. Miyamoto, T. Fujioka, N. Otsuka, K. Suzuki-Konagai and T. Ogita, *J. Am. Chem. Soc.*, 2000, **122**, 1842–1843, DOI: [10.1021/ja992355s](https://doi.org/10.1021/ja992355s).

52 H. Tokuyama, M. Isaka, E. Nakamura, R. Ando and Y. Morinaka, *J. Antibiot.*, 1992, **45**, 1148–1154, DOI: [10.7164/antibiotics.45.1148](https://doi.org/10.7164/antibiotics.45.1148).

53 R. Thuangtong, S. Varothai, D. Triwongwaranat and C. Rujitharanawong, *J. Med. Assoc. Thailand*, 2017, **100**, 86–92, PMID: 29911776.

54 G. Singh and M. Lavanya, *Int. J. Trichol.*, 2010, **2**, 36, DOI: [10.4103/0974-7753.66911](https://doi.org/10.4103/0974-7753.66911).

55 J. A. Upitis and A. Krol, *J. Cutaneous Med. Surg.*, 2002, **6**, 214–217, DOI: [10.1007/s10227-001-0050-9](https://doi.org/10.1007/s10227-001-0050-9).

56 K. K. Veverka, J. W. Jakub and C. L. Baum, *Dermatol. Surg.*, 2018, **44**, 1501–1508, DOI: [10.1097/DSS.0000000000001603](https://doi.org/10.1097/DSS.0000000000001603).

57 C. Robles-Planells, S. A. Michelson, J. Mena, D. Escrig, J. L. Rojas, G. Sanchez-Guerrero, R. Hernández, C. Barrera-Avalos, L. E. Rojo and D. Sauma, *Front. Pharmacol.*, 2019, **10**, 1201, DOI: [10.3389/fphar.2019.01201](https://doi.org/10.3389/fphar.2019.01201).

58 R. Breslow and G. Ryan, *J. Am. Chem. Soc.*, 1967, **89**, 3073, DOI: [10.1021/ja00988a063](https://doi.org/10.1021/ja00988a063).

59 Z. L. Cheng and Q. Y. Chen, *Chin. J. Chem.*, 2006, **24**, 1219–1224, DOI: [10.1002/cjoc.200690227](https://doi.org/10.1002/cjoc.200690227).

60 M. Nakamura, H. Isobe and F. Nakamura, *Chem. Rev.*, 2003, **103**, 1295–1326, DOI: [10.1021/cr0100244](https://doi.org/10.1021/cr0100244).

61 R. Breslow, L. J. Altman, A. Krebs, E. Mohacs, I. Murata, R. A. Peterson and J. Posner, *J. Am. Chem. Soc.*, 1965, **87**, 1326–1331, DOI: [10.1021/ja01084a030](https://doi.org/10.1021/ja01084a030).

62 D. B. Werz, R. Gleiter and F. Rominger, *Eur. J. Org. Chem.*, 2003, **2003**, 151–154, DOI: [10.1002/1099-0690\(200301\)2003:1<151::AID-EJOC151>3.0.CO;2-7](https://doi.org/10.1002/1099-0690(200301)2003:1<151::AID-EJOC151>3.0.CO;2-7).

63 J. F. McGarry, J. Prodollet and T. Smyth, *Org. Magn. Reson.*, 1981, **17**, 59–65, DOI: [10.1002/mrc.1270170114](https://doi.org/10.1002/mrc.1270170114).



64 P. A. Wender, T. J. Paxton and T. J. Williams, *J. Am. Chem. Soc.*, 2006, **128**, 14814–14815, DOI: [10.1021/ja065868p](https://doi.org/10.1021/ja065868p).

65 Z. X. Zhao, H. Y. Wang, C. Xu and Y. L. Guo, *Rapid Commun. Mass Spectrom.*, 2010, **24**, 2665–2672, DOI: [10.1002/rcm.4694](https://doi.org/10.1002/rcm.4694).

66 O. J. Curnow, G. M. Fern and R. J. Pipal, *ARKIVOC*, 2006, (iii), 43–47.

67 Z. Yoshida, H. Konishi, Y. Tawara, K. Nishikawa and H. Ogoshi, *Tetrahedron Lett.*, 1973, **14**, 2619–2622, DOI: [10.1016/S0040-4039\(00\)87412-2](https://doi.org/10.1016/S0040-4039(00)87412-2).

68 K. Elena, K. Tatiana, L. Irina, D. C. Jorge, H. O. Simon, M. I. Daniel and M. G. Marcos, *Open Org. Chem. J.*, 2008, **2**, 35–40, DOI: [10.2174/1874095200801020035](https://doi.org/10.2174/1874095200801020035).

69 S. W. Tobey and R. West, *J. Am. Chem. Soc.*, 1964, **86**, 4215–4216, DOI: [10.1021/ja01073a075](https://doi.org/10.1021/ja01073a075).

70 P. J. Whitman and B. M. Trost, *J. Am. Chem. Soc.*, 1969, **91**, 7534–7535, DOI: [10.1021/ja01054a067](https://doi.org/10.1021/ja01054a067).

71 S. S. Nguyen, A. Ferreira, Z. G. Long, T. K. Heiss, R. S. Dorn, R. D. Row and J. A. Prescher, *Org. Lett.*, 2019, **21**, 8695–8699, DOI: [10.1021/acs.orglett.9b03298](https://doi.org/10.1021/acs.orglett.9b03298).

72 S. Cunha, J. C. Serafim, L. L. B. de Santana, F. Damasceno, J. T. M. Correia, A. O. Santos, M. Oliveira, J. Ribeiro, J. Amparo and S. L. Costa, *J. Heterocycl. Chem.*, 2017, **54**, 3700–3710, DOI: [10.1002/jhet.2921](https://doi.org/10.1002/jhet.2921).

73 A. Kascheres, H. C. Schumacher and R. A. Rodrigues, *J. Heterocycl. Chem.*, 1997, **34**, 757–759, DOI: [10.1002/jhet.5570340309](https://doi.org/10.1002/jhet.5570340309).

74 M. A. M. Gomaa, *J. Chem. Soc., Perkin Trans. 1*, 2002, **3**, 341–344, DOI: [10.1039/B109711N](https://doi.org/10.1039/B109711N).

75 A. Haito and N. Chatani, *Chem. Commun.*, 2019, **55**, 5740–5742, DOI: [10.1039/C9CC02397F](https://doi.org/10.1039/C9CC02397F).

76 T. Nanda and P. Ravikumar, *Org. Lett.*, 2020, **22**, 1368–1374, DOI: [10.1021/acs.orglett.9b04656](https://doi.org/10.1021/acs.orglett.9b04656).

77 N. Lozinskaya, S. Sosonyuk, Y. N. Firsova, M. Proskurnina and N. Zefirov, *Russ. Chem. Bull.*, 2011, **60**, 1989–1994, DOI: [10.1007/s11172-011-0301-x](https://doi.org/10.1007/s11172-011-0301-x).

78 X. Li, C. Han, H. Yao and A. Lin, *Org. Lett.*, 2017, **19**, 778–781, DOI: [10.1021/acs.orglett.6b03737](https://doi.org/10.1021/acs.orglett.6b03737).

79 K. Reitel, K. Kriis, I. Järving and T. Kanger, *Chem. Heterocycl. Compd.*, 2018, **54**, 929–933, DOI: [10.1007/s10593-018-2372-1](https://doi.org/10.1007/s10593-018-2372-1).

80 T. Matsuda, Y. Tabata and H. Suzuki, *New J. Chem.*, 2018, **42**, 19178–19182, DOI: [10.1039/C8NJ04579H](https://doi.org/10.1039/C8NJ04579H).

81 J. T. Ren, J. X. Wang, H. Tian, J. L. Xu, H. Hu, M. Aslam and M. Sun, *Org. Lett.*, 2018, **20**, 6636–6639, DOI: [10.1021/acs.orglett.8b02612](https://doi.org/10.1021/acs.orglett.8b02612).

82 D. Bai, Y. Yu, H. Guo, J. Chang and X. Li, *Angew. Chem., Int. Ed.*, 2020, **59**, 2740–2744, DOI: [10.1002/anie.201913130](https://doi.org/10.1002/anie.201913130).

83 K. Takahashi and S. Tarutani, *J. Chem. Soc., Chem. Commun.*, 1994, **4**, 519–520, DOI: [10.1039/C39940000519](https://doi.org/10.1039/C39940000519).

84 T. Matsuda and Y. Sakurai, *J. Org. Chem.*, 2014, **79**, 2739–2745, DOI: [10.1021/jo500045n](https://doi.org/10.1021/jo500045n).

85 J. Xu, J. Cao, C. Fang, T. Lu and D. Du, *Org. Chem. Front.*, 2017, **4**, 560–564, DOI: [10.1039/C6QO00734A](https://doi.org/10.1039/C6QO00734A).

86 S. Cunha and Z. N. da Rocha, *Quim. Nova*, 2008, **31**, 788–792, DOI: [10.1590/S0100-40422008000400015](https://doi.org/10.1590/S0100-40422008000400015).

87 A. R. Rivero, I. Fernandez, C. R. de Arellano and M. A. Sierra, *J. Org. Chem.*, 2015, **80**, 1207–1213, DOI: [10.1021/jo502292y](https://doi.org/10.1021/jo502292y).

88 J. R. Syu, C. H. Lin, C. W. Kuo and D. Y. Yang, *Tetrahedron Lett.*, 2014, **55**, 1207–1211, DOI: [10.1016/j.tetlet.2013.12.115](https://doi.org/10.1016/j.tetlet.2013.12.115).

89 Y. Shi, H. Xing, T. Huang, X. Liu, J. Chen, X. Guo, G. B. Li and Y. Wu, *Chem. Commun.*, 2020, **56**, 1585–1588, DOI: [10.1039/C9CC08926H](https://doi.org/10.1039/C9CC08926H).

90 M. C. Nakhla and J. L. Wood, *J. Am. Chem. Soc.*, 2017, **139**, 18504–18507, DOI: [10.1021/jacs.7b12570](https://doi.org/10.1021/jacs.7b12570).

91 S. Cunha, F. Damasceno and J. Ferrari, *Tetrahedron Lett.*, 2007, **48**, 5795–5798, DOI: [10.1016/j.tetlet.2007.06.087](https://doi.org/10.1016/j.tetlet.2007.06.087).

92 L. Yao, Q. Hu, L. Bao, W. Zhu and Y. Hu, *Org. Lett.*, 2021, **23**, 4971–4975, DOI: [10.1021/acs.orglett.1c01304](https://doi.org/10.1021/acs.orglett.1c01304).

93 S. J. Zhou, X. Cheng and J. Xuan, *Asian J. Org. Chem.*, 2019, **8**, 1376–1379, DOI: [10.1002/ajoc.201900272](https://doi.org/10.1002/ajoc.201900272).

94 M. Lange, J. C. Tendyck, P. Wegener, A. Hepp, E. U. Würthwein and W. Uhl, *Chem.-Eur. J.*, 2018, **24**, 12856–12868, DOI: [10.1002/chem.201706089](https://doi.org/10.1002/chem.201706089).

95 J. Wu, W. X. Gao, X. B. Huang, Y. B. Zhou, M. C. Liu and H. Y. Wu, *Org. Lett.*, 2020, **22**, 5555–5560, DOI: [10.1021/acs.orglett.0c01914](https://doi.org/10.1021/acs.orglett.0c01914).

96 L. H. Li, Y. Jiang, J. Hao, Y. Wei and M. Shi, *Adv. Synth. Catal.*, 2017, **359**, 3304–3310, DOI: [10.1002/adsc.201700936](https://doi.org/10.1002/adsc.201700936).

97 A. A. Hassan, F. F. Abdel-Latif, A. M. N. El-Din, S. M. Mostafa, M. Nieger and S. Braese, *Tetrahedron*, 2012, **68**, 8487–8492, DOI: [10.1016/j.tet.2012.07.063](https://doi.org/10.1016/j.tet.2012.07.063).

98 A. A. Aly, A. A. Hassan, M. A. Ameen and A. B. Brown, *Tetrahedron Lett.*, 2008, **49**, 4060–4062, DOI: [10.1016/j.tetlet.2008.04.066](https://doi.org/10.1016/j.tetlet.2008.04.066).

99 P. Zhou, W. T. Yang, A. U. Rahman, G. Li and B. Jiang, *J. Org. Chem.*, 2019, **85**, 360–366, DOI: [10.1021/acs.joc.9b02253](https://doi.org/10.1021/acs.joc.9b02253).

100 W. T. Zhao, X. Y. Tang and M. Shi, *Eur. J. Org. Chem.*, 2014, **2014**, 2672–2676, DOI: [10.1002/ejoc.201400077](https://doi.org/10.1002/ejoc.201400077).

101 J. Wallbaum, P. G. Jones and D. B. Werz, *J. Org. Chem.*, 2015, **80**, 3730–3734, DOI: [10.1021/acs.joc.5b00330](https://doi.org/10.1021/acs.joc.5b00330).

102 L. Liu, J. Li, W. Dai, F. Gao, K. Chen, Y. Zhou and H. Liu, *Molecules*, 2020, **25**, 268, DOI: [10.3390/molecules25020268](https://doi.org/10.3390/molecules25020268).

103 J. Cao, R. Fang, J. Y. Liu, H. Lu, Y. C. Luo and P. F. Xu, *Chem.-Eur. J.*, 2018, **24**, 18863–18867, DOI: [10.1002/chem.201803861](https://doi.org/10.1002/chem.201803861).

104 J. Chen, B. Tang, X. Liu, G. Lv, Y. Shi, T. Huang, H. Xing, X. Guo, L. Hai and Y. Wu, *Org. Chem. Front.*, 2020, **7**, 2944–2949, DOI: [10.1039/D0QO00769B](https://doi.org/10.1039/D0QO00769B).

105 S. W. Liu, C. Yuan, X. F. Jiang, X. X. Wang and H. L. Cui, *Asian J. Org. Chem.*, 2020, **9**, 82–85, DOI: [10.1002/ajoc.201900630](https://doi.org/10.1002/ajoc.201900630).

106 D. H. Wadsworth, C. H. Weidner, S. L. Bender, R. H. Nuttall and H. R. Luss, *J. Org. Chem.*, 1989, **54**, 3652–3660, DOI: [10.1021/jo00276a028](https://doi.org/10.1021/jo00276a028).

107 T. Kondo, Y. Kaneko, Y. Taguchi, A. Nakamura, T. Okada, M. Shiotsuki, Y. Ura, K. Wada and T.-a. Mitsudo, *J. Am. Chem. Soc.*, 2002, **124**, 6824–6825, DOI: [10.1021/ja0260521](https://doi.org/10.1021/ja0260521).

108 W.-Q. Zhu, Y. C. Fang, W. Y. Han, F. Li, M. G. Yang and Y.-Z. Chen, *Org. Chem. Front.*, 2021, **8**, 3082–3090, DOI: [10.1039/D1QO00222H](https://doi.org/10.1039/D1QO00222H).



109 Q. Chen, Y. Teng and F. Xu, *Org. Lett.*, 2021, **23**, 4785–4790, DOI: [10.1021/acs.orglett.1c01506](https://doi.org/10.1021/acs.orglett.1c01506).

110 S. Ahmad, L. Shukla, J. Szawkalo, P. Roszkowski, J. K Maurin and Z. Czarnocki, *Catalysis Communications*, 2017, **89**, 44–47, DOI: [10.1016/j.catcom.2016.10.008](https://doi.org/10.1016/j.catcom.2016.10.008).

111 M. Takahashi, Y. Kadokawa, Y. Uno and Y. Nakano, *Heterocycles*, 1999, **51**(9), 637–639, DOI: [10.3987/COM-99-8652](https://doi.org/10.3987/COM-99-8652).

112 A. A. Aly, M. Ramadan, M. A. Al-Aziz, H. M. Fathy, S. Bräse, A. B. Brown and M. Nieger, *J. Chem. Res.*, 2016, **40**(10), 637–639, DOI: [10.3184/174751916X14743924874916](https://doi.org/10.3184/174751916X14743924874916).

113 A. A. Aly, A. M. Nour El-Din, M. A.-M. Gomaa, A. B. Brown and A. S. Fahmi, Unusual Reactivity of 2,3-diphenylcyclopropenone towards *N*-imidoylthioureas; Facile Synthesis of 3-aryl-2,5,6-triphenylpyrimidin-4(3*H*)-one (PART III), *J. Chem. Res.*, 2007, **2007**(8), 439–441, DOI: [10.3184/030823407X234563](https://doi.org/10.3184/030823407X234563).

114 R. D. Mohan and A. Jose, *Asian J. Chem.*, 2018, **30**, 1075–1077.

115 B. Niu, B. Jiang, L. Z. Yu and M. Shi, *Org. Chem. Front.*, 2018, **5**, 1267–1271, DOI: [10.1039/C8QO00091C](https://doi.org/10.1039/C8QO00091C).

116 Y. F. Yang, X. B. Huang, W. X. Gao, Y. B. Zhou, M. C. Liu and H. Y. Wu, *Org. Biomol. Chem.*, 2020, **18**, 5822–5825, DOI: [10.1039/D0OB00601G](https://doi.org/10.1039/D0OB00601G).

117 L. Sizhan, C. Mingyue, W. Bowen, H. Chunmei, Z. Yingying, L. Jing, X. Xuetao, W. Zhen and W. Shaohua, *Chin. J. Org. Chem.*, 2021, **41**, 1622–1630, DOI: [10.6023/cjoc202010041](https://doi.org/10.6023/cjoc202010041).

118 T. Matsuda, K. Yamanaka, Y. Tabata and T. Shiomi, *Tetrahedron Lett.*, 2018, **59**, 1458–1460, DOI: [10.1016/j.tetlet.2018.02.084](https://doi.org/10.1016/j.tetlet.2018.02.084).

119 A. A. Hassan, N. K. Mohamed, A. A. Aly, H. N. Tawfeek, H. Hopf, S. Bräse and M. Nieger, *Mol. Diversity*, 2019, **23**, 821–828, DOI: [10.1007/s11030-018-09912-5](https://doi.org/10.1007/s11030-018-09912-5).

120 E. M. El-Sheref, *J. Sulfur Chem.*, 2017, **38**, 625–634, DOI: [10.1080/17415993.2017.1337121](https://doi.org/10.1080/17415993.2017.1337121).

121 Y. Shi, T. Huang, T. Wang, J. Chen, X. Liu, Z. Wu, X. Huang, Y. Zheng, Z. Yang and Y. Wu, *Chem.-Eur. J.*, 2021, **27**, 13346–13351, DOI: [10.1002/chem.202101839](https://doi.org/10.1002/chem.202101839).

122 S. Ahmad, L. Shukla, J. Szawkalo, P. Roszkowski, J. K. Maurin and Z. Czarnocki, *Catal. Commun.*, 2017, **89**, 44–47, DOI: [10.1016/j.catcom.2016.10.008](https://doi.org/10.1016/j.catcom.2016.10.008).

123 O. B. Østby, L. L. Gundersen, F. Rise, Ø. Antonsen, K. Fosnes, V. Larsen, A. Bast, I. Custers and G. R. Haenen, *Arch. Pharm.*, 2001, **334**, 21–24, DOI: [10.1002/1521-4184\(200101\)334:1<21::AID-ARDP21>3.0.CO;2-W](https://doi.org/10.1002/1521-4184(200101)334:1<21::AID-ARDP21>3.0.CO;2-W).

124 A. Molchanov, D. Sipkin, Y. B. Koptelov and R. Kostikov, *Russ. J. Org. Chem.*, 2005, **41**, 567–574, DOI: [10.1007/s11178-005-0205-z](https://doi.org/10.1007/s11178-005-0205-z).

125 A. A. Aly, A. A. Hassan, M. A.-M. Gomaa and E. M. El-Sheref, *ARKIVOC*, 2007, **xiv**, 1–11, DOI: [10.3998/ark.5550190.0008.e01](https://doi.org/10.3998/ark.5550190.0008.e01).

126 K. Matsumoto, A. Okada, T. Girek, Y. Ikemi, J. C. Kim, N. Hayashi, H. Yoshida and A. Kakehi, *Heterocycl. Commun.*, 2002, **8**, 325–328, DOI: [10.1515/HC.2002.8.4.325](https://doi.org/10.1515/HC.2002.8.4.325).

127 W. T. Zhao, F. Gao and D. Zhao, *Angew. Chem.*, 2018, **130**, 6437–6440, DOI: [10.1002/ange.201803156](https://doi.org/10.1002/ange.201803156).

128 K. Hemming, P. A. O'Gorman and M. I. Page, *Tetrahedron Lett.*, 2006, **47**, 425–428, DOI: [10.1016/j.tetlet.2005.11.081](https://doi.org/10.1016/j.tetlet.2005.11.081).

