



Cite this: *RSC Adv.*, 2022, **12**, 23680

Correction: Chrysomycins A–C, antileukemic naphthocoumarins from *Streptomyces sporoverrucosus*

Shreyans K. Jain,^{ab} Anup S. Pathania,^{bc} Rajinder Parshad,^d Chandji Raina,^d Asif Ali,^{ab} Ajai P. Gupta,^e Manoj Kushwaha,^e Subrayashastry Aravinda,^f Shashi Bhushan,^{bc} Sandip B. Bharate^{*bf} and Ram A. Vishwakarma^{*abf}

DOI: 10.1039/d2ra90078e

rsc.li/rsc-advances

Correction for 'Chrysomycins A–C, antileukemic naphthocoumarins from *Streptomyces sporoverrucosus*' by Shreyans K. Jain *et al.*, *RSC Adv.*, 2013, **3**, 21046–21053, <https://doi.org/10.1039/c3ra42884b>.

The authors regret that incorrect versions of Fig. 6 and Fig. 7 were included in the original article. The correct versions of Fig. 6 and 7 are presented below.

^aNatural Products Chemistry Division, Indian Institute of Integrative Medicine (CSIR), Canal Road, Jammu, 180001, India. E-mail: ram@iiim.ac.in; Fax: +91-191-2569333; Tel: +91-191-2569111

^bAcademy of Scientific & Innovative Research (AcSIR), Anusandhan Bhawan, 2 Rafi Marg, New Delhi, 110001, India

^cCancer Pharmacology Division, Indian Institute of Integrative Medicine (CSIR), Canal Road, Jammu, 180001, India

^dFermentation Division, Indian Institute of Integrative Medicine (CSIR), Canal Road, Jammu, 180001, India

^eQuality Control and Quality Assurance Division, Indian Institute of Integrative Medicine (CSIR), Canal Road, Jammu, 180001, India

^fMedicinal Chemistry Division, Indian Institute of Integrative Medicine (CSIR), Canal Road, Jammu, 180001, India. E-mail: sbharate@iiim.ac.in






Fig. 6 Influence of compounds 1–3 on the nuclear morphology of human leukaemia HL-60 cells. The cells were treated with 1, 3 and 5 μM concentrations of these compounds for 24 h and stained with Hoechst 33258 for 40 min. The altered nuclear morphology and apoptotic bodies indicated by white arrows are seen in treated cells while the nuclei of the untreated cells were round and intact.






Fig. 7 Phase contrast microscopy of compound-treated leukaemia HL-60 cells. Cells were treated with compounds 1–3 at 1, 3 and 5 μM for 24 h and visualized using a phase contrast microscope (Olympus1X72). The morphology of treated cells altered in a concentration-dependent manner, while the untreated cells remained healthy.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

