Chemical Science



EDGE ARTICLE

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2022, 13, 11607

d All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 24th March 2022 Accepted 11th July 2022

DOI: 10.1039/d2sc01716d

rsc.li/chemical-science

Nickel-catalyzed arylative substitution of homoallylic alcohols†

Hai N. Tran, (D) Chau M. Nguyen, Mason T. Koeritz, Dustin D. Youmans and Levi M. Stanley (D)*

Direct coupling of unactivated alcohols remains a challenge in synthetic chemistry. Current approaches to cross-coupling of alcohol-derived electrophiles often involve activated alcohols such as tosylates or carbonates. We report the direct arylative substitution of homoallylic alcohols catalyzed by a nickel-bisphosphine complex as a facile method to generate allylic arenes. These reactions proceed *via* formation of an allylic alcohol intermediate. Subsequent allylic substitution with arylboroxine nucleophiles enables the formation of a variety of allylic arenes. The presence of *p*-methoxyphenylboronic acid is crucial to activate the allylic alcohol to achieve high product yields.


Introduction

Alkanols are among the most common and accessible functional groups in organic synthesis and are attractive as potential C(sp³) coupling partners to form C(sp³)–C bonds. However, due to the relatively strong bond dissociation energy of the C–O bond and poor leaving group ability of the OH group,¹ alcohols are rarely used directly as alkylating agents in cross-coupling chemistry.² Although numerous alcohol derivatives (*i.e.* tosylates, carbonates, *etc.*) have been extensively explored in redoxneutral cross-coupling reactions,³,⁴ these derivatives typically require one or more preparation steps from the corresponding alcohol precursors. Therefore, processes that form C(sp³)–C bonds directly from unactivated alcohols have the potential to streamline synthetic routes.

Transition metal-catalyzed hydroarylations of 1,3-dienes are privileged synthetic methods to generate allylic arenes and allylic heteroarenes. Traditionally, hydro(hetero)arylation of 1,3-dienes involves activation of an aryl C–H bond (Scheme 1a, top). Alternatively, hydro(hetero)arylations of 1,3-dienes have been developed by employing alcohols or silanes as external hydride sources (Scheme 1a, bottom). In the past decade, approaches to tandem transition metal-catalyzed hydro(hetero) arylation of 1,3-dienes generated *in situ* from homoallylic electrophiles have been reported. Sigman and co-workers developed Pd-catalyzed arylative substitutions of primary and secondary homoallylic tosylates with arylboronic acids (Scheme 1b). In 2020, Kawatsura and co-workers developed the arylative substitution of secondary homoallylic carbonates with

Department of Chemistry, Iowa State University, Ames, IA 50011, USA. E-mail: lstanley@iastate.edu

[†] Electronic supplementary information (ESI) available: Full experimental details, characterization data, and NMR spectra for all new compounds. See https://doi.org/10.1039/d2sc01716d



Scheme 1 Transition metal-catalyzed formation of allylic arenes *via* hydroarylation and arylative substitutions.

arylboronic acids catalyzed by a nickel complex of a monophosphine ligand (Scheme 1c). 10 These reactions proceed via in situ formation of a diene by a sequence of oxidative addition of the C-O bond to generate a metal-alkyl intermediate and subsequent β-hydride elimination. Alkene insertion to the metal-hydride bond, transmetallation with an arylboronic acid, and reductive elimination generate the allylic arene product. These formal hydroarylations of 1,3-dienes are attractive examples of coupling reactions involving secondary sp³ electrophiles. However, the opportunity exists to improve these reactions by eliminating the need for an activated homoallylic electrophile. To this end, we now report the development of nickel-catalyzed arylative substitution of unactivated secondary homoallylic alcohols with arylboroxine nucleophiles (Scheme 1d). These arylative substitution reactions occur via in situ formation of allylic alcohols through alkene isomerization, followed by allylic arylation12-14 to generate a variety of allylic arenes in short reaction times.

Results and discussion

Initially, reactions of 1-(*p*-methoxyphenyl)but-3-en-1-ol **1a** with common arylboron nucleophiles were evaluated to identify a suitable aryl coupling partner. Reactions did not occur when *p*-MeOC₆H₄B(OH)₂, *p*-MeOC₆H₄Bpin, and *p*-MeOC₆H₄Bneop

were used as the coupling partners (Table 1, entries 1-3). However, the reaction of 1 equiv. of p-methoxyphenylboroxine 2a (12:1 mixture of arylboroxine: arylboronic acid) with 1a formed the product 3aa in 95% yield (entry 4). The yield of the reaction remained high upon lowering the loading of 2a to 0.6 equiv. (entry 5). Interestingly, the reaction of 1a with arylboroxine 2a containing a higher ratio of boroxine to its corresponding boronic acid (arylboroxine : arylboronic acid = 16:1, entry 6) led to a drop in the yield of 3aa to 71%. We hypothesized that the catalytic amount of p-methoxyphenylboronic acid could play a role in activating the homoallylic alcohol 1a.15 Indeed, adding 0.25 equiv. of p-MeOC₆H₄B(OH)₂ to the reaction of 0.6 equiv. of boroxine 2a with 1a led to the formation of the desired product 3aa in 98% yield (entry 7). The addition of pmethoxyphenylboronic acid allowed the amount of arylboroxine 2a to be lowered to 0.5, 0.4 or 0.33 equiv. without significant impact on the yield of 3aa (entries 8-10). Since electrondeficient arylboronic acids are expected to activate the alcohol effectively,15a more we attempted to use p-trifluoromethylphenylboronic acid p-methoxand yearbonyphenylboronic acid as additives in the model reaction to activate the homoallylic alcohol. However, due to high content of corresponding arylboroxines in these commercially available, electron-deficient arylboronic acids, we observed the formation of the allylic arene byproducts derived from these

Table 1 Identification of reaction conditions^a

Entry	Deviation from standard conditions	Yield 3aa ^b (%)
1	2 equiv. p -MeOC ₆ H ₄ B(OH) ₂ instead of 2a	0
2	2 equiv. p-MeOC ₆ H ₄ Bpin instead of 2a	0
3	2 equiv. p-MeOC ₆ H ₄ Bneop instead of 2a	0
4	1 equiv. 2a (arylboroxine : arylboronic acid = 12 : 1)	95
5	None (arylboroxine : arylboronic acid $= 12:1$)	93
6	None (arylboroxine : arylboronic acid $= 16 : 1$)	71
7 ^c	0.25 equiv. p-MeOC ₆ H ₄ B(OH) ₂ added	98 (93) ^d
8 ^c	0.5 equiv. 2a, 0.25 equiv. p -MeOC ₆ H ₄ B(OH) ₂ added	90
9^c	0.4 equiv. 2a, 0.25 equiv. p -MeOC ₆ H ₄ B(OH) ₂ added	88
10^c	0.33 equiv. 2a, 0.25 equiv. p -MeOC ₆ H ₄ B(OH) ₂ added	84
11 ^e	2 equiv. p-MeOC ₆ H ₄ Bneop instead of 2a	0
12^e	2 equiv. p-OMeC ₆ H₄Bpin instead of 2a	0

^a Reaction conditions: **1a** (0.25 mmol), **2a** (0.15 mmol), Ni(cod)₂ (0.0188 mmol), Xantphos (0.0188 mmol), K₃PO₄ (0.162 mmol), THF (1.1 mL) at 65 °C for 2 h under a N₂ atmosphere. E/Z ratios were determined by ¹H NMR spectroscopy of crude mixtures. ^b Yields were determined by ¹H NMR spectroscopy of the crude reaction mixture using CH_2Br_2 as an internal standard. ^c **2a** with arylboroxine : arylboronic acid ratio = 16 : 1. ^d Isolated yield. ^e 0.25 equiv. p-MeOC₆H₄B(OH)₂ added.

electron-deficient arylboroxines. In contrast, using p-methoxyphenylboronic acid as an additive, we did not observe the allylic arene byproduct derived from the p-methoxyphenylboroxine when evaluating the arylboroxine scope (see Scheme 4, below). Therefore, we chose p-methoxyphenylboronic acid as an additive in our arylative coupling reactions. In addition we reevaluated the arylboronic ester nucleophiles with p-methoxyphenylboronic acid as an additive. Reactions did not occur when the arylboronic acid pinacol ester and arylboronic acid neopentylglycol ester were used as the coupling partner in the presence of 0.25 equiv. of p-MeOC₆H₄B(OH)₂ (entries 11–12). We chose reaction conditions using 0.6 equiv. of 2a with the addition of 0.25 equiv. of p-MeOC₆H₄B(OH)₂ to evaluate the scope of the arylative coupling reaction.

With reaction conditions identified that lead to high yields in our model reaction, we sought to establish the scope of the reaction by evaluating arylative substitutions of a variety of homoallylic alcohols (Scheme 2). An array of α -aryl-substituted homoallylic alcohols 1 react with p-methoxyphenylboroxine 2a to generate allylic arene products 3 in modest-to-high yields (33–95%). The reaction of chlorinated homoallylic alcohol (1d) furnished the product 3da in 33% yield with isomerization of 1d to a mixture of allylic alcohols accounting for the balance of the mass. It is likely that oxidative addition of the Ni catalyst to the C_{aryl} -Cl bond is competitive with the desired arylate substitution. However, Suzuki-Miyaura coupling products are not observed. Notably, a homoallylic alcohol bearing an ester

MeC₆H₄BO)₃ **2a** (0.6 equiv) Ni(cod)₂ (7.5 mol%) Xantphos (7.5 mol%) OMeC₆H₄B(OH)₂ (0.25 equiv) K₂PO₄ (0.65 equiv) 65°C 2 h 0.25 mmol (E/Z = >98:2)3aa, R = OMe, 93% 3ca, R = F, 91% 3da, R = Cl, 33% 3fa, R = CO₂Me, 82% 3ga, 86% OMe, 88% R = Me. 80% 3na. 86% 3la, 95% 3ma, 86% 3oa, 48% 3pa, 39% 3qa, 66%b Challenging Substrates 1r, 0% 1s. 0% 1t, 0% 1u.0%

Scheme 2 Homoallylic alcohol substrate scope. Reactions were carried out on a 0.25 mmol scale. 2a with arylboroxine : arylboronic acid = 16:1. E/Z ratios were determined by 1H NMR spectroscopy of crude mixtures. Yields of isolated products are shown. a Homoallylic alcohol 1p (syn:anti=1:1). b Isolated yield of a mixture of 3,4- and 1.4-isomers.

substituent at the aryl moiety (1f) was tolerated under the standard reaction conditions, furnishing the allyic arene 3fa in 82% yield. Homoallylic alcohols with polysubstituted α-aryl groups 1k-m and α-naphthyl homoallylic alcohol 1n react to form allylic arenes 3ka-3na in good-to-high yields (72-95%). Moreover, the reaction of α -thienyl homoallylic alcohol 10 generated the corresponding heteroarene-containing product 30a in 48%. Additionally, the reaction of homoallylic alcohol 1p bearing a β-substituent furnished the corresponding allylic arene **3pa** in 39% yield with 3:1 E/Z ratio. The reaction of α alkyl-substituted homoallylic alcohol 1q with boroxine 2a formed the corresponding product 3qa in 63% yield with 2:1 rr. However, reactions of tertiary homoallylic alcohol 1r, homoallylic alcohols containing 1,2- and 1,1-disubstituted alkenes (1s and 1t), and bishomoallylic alcohol 1u did not occur under our reaction conditions.

We attempted to extend the arylative substitution reaction to encompass α -styrenyl homoallylic alcohols (Scheme 3). Interestingly, the reaction of α -styrenyl homoallylic alcohol 1v with boroxine 2a did not generate the desired allylic arene 3va. Instead, product 4va was isolated in 53% yield. The observation of the aryl nucleophile replacing the OH at the α -position in product 4va in lieu of the γ -position like products 3aa-3qa suggests our arylative substitution reactions might proceed via a different mechanism compared to the previously reported formal hydroarylations of homoallylic alcohol derivatives (Scheme 1b and c).

Next, we examined the scope of the arylative substitution reactions with respect to arylboroxine coupling partners (Scheme 4). Reactions of 1-(p-methoxyphenyl)but-3-en-1-ol 1a with a variety of substituted boroxines 2 generated allylic arenes 3 in modest-to-high yields (43–94%). The scope of these arylative substitution reactions encompasses a variety of para-, meta-, ortho-, and disubstituted arylboroxines. Notably, arylboroxines containing fluoride (2d), ester (2e), chloride (2h), and acetal (2j) substituents were tolerated under our reaction conditions. Moreover, furanyl-3-boroxine 2m reacts with homoallylic alcohol 1a to form the corresponding allylic heteroarene in 53% yield.

We examined the scalability of the arylative substitution in a 5 mmol scale reaction of homoallylic alcohol **1a** (Scheme 5). This reaction formed the desired product **3aa** in 84% yield when conducted in the presence of 5 mol% nickel precatalyst and 5 mol% Xantphos ligand. In addition, the synthetic utility was

Scheme 3 Arylative substitution of α -styrenyl homoallylic alcohol. Reaction was carried out on a 0.25 mmol scale. Yield of isolated product 4va is shown.


(A/BO)₃ 2 (0.6 equiv)
Ni(cod)₂ (7.5 mol%)
Nantphos (7.5 mol%)
Nantphos (7.5 mol%)
Nantphos (7.5 mol%)
Nantphos (9.6 equiv)
NsPo₄ (9.6 equiv)
NsPo₅ (9.6 equiv

Chemical Science


Scheme 4 Arylboroxine substrate scope. Reactions were carried out on a 0.25 mmol scale. 2 with arylboroxine : arylboronic acid > 10 : 1. E/Z ratios were determined by 1H NMR spectroscopy of crude mixtures. Yields of isolated products are shown.

demonstrated by subjecting the allylic arene 3aa to alkene functionalization transformations. The reaction of 3aa with mCPBA, followed by workup with saturated aq. NaHCO $_3$ furnished the hydroxyester 5aa in 65% yield as a 1:1 ratio of separable diastereomers. Transition metal-free sulfuration/annulation¹⁷ of 3aa formed thiophene 6aa in 53% yield.

To gain insights into the reaction mechanism, we conducted a series of control experiments. In previous studies reported by the Sigman (Scheme 1b)⁹ and Kawatsura groups (Scheme 1c),¹⁰ a 1,3-diene is proposed to be generated *in situ*. To probe whether our arylative substitution proceeds *via* a similar mechanism, we subjected the diene 7a to the standard reaction conditions. Compound 7a was recovered in 97% yield, and the allylic arene 3aa was not detected (Scheme 6a, top). In the presence of MeOH as a hydride source, ^{7b,8c,18} the hydroarylation of diene 7a formed the allylic arene 3aa in 40% yield. We also conducted the hydroarylation of 1,4-diphenyl-1,3-diene 7b, furnishing the allylic arene 3sa in 10% (Scheme 6a, bottom).^{7b} In contrast, the reaction of homoallylic alcohol 1s with boroxine 2a did not generate 3sa (Scheme 2). These observations suggest the



Scheme 5 Large-scale reaction and synthetic transformations. Yields of isolated products are shown. Reaction conditions for (a): 3aa (0.3 mmol), mCPBA (0.45 mmol), DCM (3 mL), rt, 24 h then sat. NaHCO $_3$ (2 mL), rt, 5 min. Reaction conditions for (b): 3aa (0.3 mmol), K_2S (0.9 mmol), DMSO (2 mL), 140 °C, 24 h under air.



Scheme 6 Control experiments. Yields were determined by ¹H NMR spectroscopy of the crude reaction mixture using CH₂Br₂ as an internal standard.

mechanism involving *in situ* generation of a 1,3-diene is not likely in our arylative substitution reactions.

We then turned our focus to the potential for in situ formation of an allylic alcohol via isomerization of the homoallylic alcohol, followed by allylic arylation to form the allylic arene products 3. The reaction of allylic alcohol (E)-8a and boroxine 2a under the standard conditions produced allylic arene 3aa in 90% (with p-MeOC₆H₄B(OH)₂) and 55% (without p-MeOC₆H₄-B(OH)₂) yields, suggesting the arylative substitution may occur via an allylic alcohol intermediate (Scheme 6b). In addition, allylic alcohol 9a reacts with boroxine 2a under standard reaction conditions to form allylic arene 3ba in 90% yield (Scheme 6c). These observations suggest the arylative substitutions of allylic alcohols (E)-8a and 9a proceed via π -allylnickel(II) intermediates.13,19 Furthermore, the reactions of these allylic alcohols produce the corresponding allylic arenes in higher yields in the presence of p-methoxyphenylboronic acid, suggesting pmethoxyphenylboronic acid is involved in activation of the allylic alcohols.15


We continued our mechanistic studies by probing the *in situ* formation of the allylic alcohol. We envisioned four potential pathways to form the allylic alcohol. Pathway 1 involves the intramolecular hydronickelation of homoallylic alcohol 1 to form the oxanickelacycle, followed by β -hydride elimination (Scheme 7a).²⁰ Pathway 2 consists of the oxidative addition of nickel catalyst to the O–H bond of p-MeOC₆H₄B(OH)₂, subsequent migratory insertion of the alkene present in 1, and β -hydride elimination (Scheme 7b). Pathway 3 comprises an intramolecular hydride shift involving the formation of a π -allyl-Ni(II) to enable the migration of a hydride from the

Scheme 7 Possible pathways for the in situ formation of allylic alcohol

β position to the terminal position (Scheme 7c). ^{21,22} Pathway 4 comprises an intermolecular hydride shift involving a sequential formation of a π -allyl-Ni(II), migratory insertion, and β hydride elimination (Scheme 7d).22

conducted experiments with deuterium-labelled substrates to evaluate these pathways. The reaction of deuterated homoallylic alcohol 1a-OD with boroxine 2a produced allylic arene 3aa in 90% yield without any deuterium incorporation (Scheme 8a). Similarly, homoallylic alcohol 1a reacted with boroxine 2a in the presence of deuterated p-methoxyphenylboronic acid to form 3aa in 91% yield without any deuterium incorporation (Scheme 8b). These observations rule out the *in situ* formation of allylic alcohol *via* pathways 1 and 2. Notably, in the reaction of β -deuterated homoallylic alcohol **1b**-D2, we observed deuterium incorporation (88% D) to the terminal position of product 3ba-D2 (Scheme 8c), suggesting that the pathways 3 and 4 are feasible in our arylative substitution reaction. To further distinguish these pathways, we performed the cross-over experiment using β -deuterated homoallylic alcohol 1b-D2 and nondeuterated homoallylic alcohol 1m (Scheme 8d). We observed deuterium incorporation into products arising from both homoallylic alcohol substrates. Higher amounts of deuterium were incorporated into the terminal position (31% D) of product 3ma-D2 derived from the nondeuterated homoallylic alcohol 1m compared to the product 3ba-D2 derived from the deuterated homoallylic alcohol 1b-D2 (12% D). This result indicates that the in situ formation of allylic alcohol can proceed through the intermolecular hydride shift mechanism (pathway 4, Scheme 7d), but we cannot rule out formation of the allylic alcohol via the intramolecular hydride shift mechanism (pathway 3, Scheme 7c) as a competing reaction pathway.

We also subjected homoallylic alcohol 1a to the standard reaction conditions in the absence of arylboroxine nucleophile,



Scheme 8 Deuterium-labelling experiments. Yields and deuterium ratios were determined by ¹H NMR spectroscopy of the crude reaction mixture. a Isolated yields.

p-methoxyphenylboronic acid additive, and K₃PO₄ (Scheme 9). The reaction formed the allylic alcohol 8a in 35% yield with 1.7:1 dr. This experiment, in combination with the allylic arylation of 8a (Scheme 6b) and the reactions of deuteriumlabelled homoallylic alcohol 1b-D2 (Schemes 8c and d), is consistent with in situ formation of the allylic alcohol intermediate by pathways 3 and/or 4 (Schemes 7c and d). In addition, the 1,3-diene was not observed in the isomerization reaction of homoallylic alcohol (Scheme 9), which further suggests the mechanism involving the in situ generation of 1,3-diene is not active in our arylative substitutions of homoallylic alcohols.

Scheme 9 Isomerization of homoallylic alcohol 1a. Yield was determined by ¹H NMR spectroscopy of the crude reaction mixture using CH₂Br₂ as an internal standard.

Scheme 10 Proposed mechanism.

Based on these control experiments, we propose the tandem catalytic cycles in Scheme 10. In cycle I, the isomerization of the homoallylic alcohol is proposed to occur via formation of a π allyl-Ni(II) **B** from LNi(0) catalyst and homoallylic alcohol **1**. ^{21,22} Intermediate B undergoes migratory insertion to another homoallylic alcohol 1 to form intermediate C, followed by βhydride elimination to generate allylic alcohol 8. Intermediate B can also undergo reductive elimination to form allylic alcohol 8. Next, the allylic alcohol 8 generated in cycle I is activated in situ (intermediate **D**) by coordination of the oxygen atom to the boron atom of *p*-methoxyphenylboronic acid.¹⁵ Intermediate **D** undergoes oxidative addition with Ni(0) catalyst in cycle II to produce π -allylnickel(π) intermediate F. Subsequent transmetallation with arylboroxine 2 facilitated by K₃PO₄²³ and reductive elimination forms the desired product 3 and regenerates the Ni(0) catalyst.12a In a recent study, Fang and coworkers proposed a related isomerization/substitution mechanism for nickel-catalyzed hydrocyanation of alkenyl alcohols.24

Conclusions

We have developed an arylative substitution of homoallylic alcohols catalyzed by a nickel complex prepared $in\ situ$ from Ni(cod)₂ and Xantphos ligand to generate allylic arenes under mild reaction conditions. These reactions are fast and proceed via a tandem isomerization/allylic arylation mechanism. The use of p-methoxyphenylboronic acid to activate the $in\ situ$ generated allylic alcohol leads to the formation of products in higher yields. These arylative substitutions demonstrate the utility of unactivated alcohols in transition metal-catalyzed

cross-coupling chemistry, and the requirement of preactivated alcohols is not necessary. Additional studies on the use of unactivated alcohols in nickel-catalyzed cross-coupling reactions are ongoing.

Data availability

Experimental data associated with this article are provided in the ESI.†

Author contributions

H. N. T. and L. M. S. conceived the project. H. N. T. optimized the reactions conditions, performed the experimental mechanistic studies, and completed the synthetic applications. H. N. T. and C. M. N. synthesized starting materials and evaluated the scope. H. N. T. and L. M. S. wrote the manuscript with input from C. M. N. M. T. K. and D. D. Y. performed the deuterium-labelling experiments during the revision of the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was financially supported by the National Science Foundation (CHE-1955529).

Notes and references

- 1 Handbook of Bond Dissociation Energies in Organic Compounds, ed. Y.-R. Luo, CRC Press, Boca Raton, FL, 2002.
- 2 (a) J. Jin and D. W. C. MacMillan, *Nature*, 2015, 525, 87–90;
 (b) D.-H. Lee, K.-H. Kwon and C. S. Yi, *J. Am. Chem. Soc.*,
 2012, 134, 7325–7328; (c) D.-H. Lee, K.-H. Kwon and C. S. Yi, *Science*, 2011, 333, 1613–1616.
- 3 For nickel-catalyzed stereospecific cross-coupling reactions using alcohol derivatives, see: (a) J. Xu, O. P. Bercher and M. P. Watson, *J. Am. Chem. Soc.*, 2021, **143**, 8608–8613; (b) J. Xu, O. P. Bercher, M. R. Talley and M. P. Watson, *ACS Catal.*, 2021, **11**, 1604–1612; (c) E. J. Tollefson, L. E. Hanna and E. R. Jarvo, *Acc. Chem. Res.*, 2015, **48**, 2344–2353.
- 4 For selected racemic transition metal-catalyzed cross-coupling reactions using alcohol derivatives, see: (a) M. Ohsumi, A. Ito and N. Nishiwaki, RSC Adv., 2018, 8, 35056–35061; (b) M. Ohsumi and N. Nishiwak, ACS Omega, 2017, 2, 7767–7771; (c) X.-X. Wang, M.-J. Luo and J.-M. Lu, Org. Biomol. Chem., 2015, 13, 11438–11444; (d) X.-X. Wang, B.-B. Xu, W.-T. Song, K.-X. Suna and J.-M. Lu, Org. Biomol. Chem., 2015, 13, 4925–4930; (e) Q. Chen, X.-H. Fan, L.-P. Zhang and L.-M. Yang, RSC Adv., 2015, 5, 15338–15340; (f) R. Kuwano and M. Yokogi, Chem. Commun., 2005, 5899–5901; (g) M. McLaughlin, Org. Lett., 2005, 7, 4875–4878; (h) R. Kuwano and M. Yokogi, Org. Lett., 2005, 7, 945–947.

Edge Article

(a) M. Nagamoto, H. Yorimitsu and T. Nishimura, *Org. Lett.*,
 2018, 20, 828–831; (b) W.-C. Lee, W.-C. Shih, T.-H. Wang,
 Y. Liu, G. P. A. Yap and T.-G. Ong, *Tetrahedron*, 2015, 71,
 4460–4464; (c) Y. Nakao, N. Kashihara, K. S. Kanyiva and
 T. Hiyama, *J. Am. Chem. Soc.*, 2008, 130, 16170–16171.

- 6 For related rhodium-catalyzed hydroarylation of diene electron-rich hetero- and homoarenes, see: (a) L. Gu, L. M. Wolf, A. Zielinski, W. Thiel and M. Alcarazo, J. Am. Chem. Soc., 2017, 139, 4948–4953; (b) C. C. Roberts, D. M. Matías, M. J. Goldfogel and S. J. Meek, J. Am. Chem. Soc., 2015, 137, 6488–6491.
- 7 (a) L. Liao and M. S. Sigman, J. Am. Chem. Soc., 2010, 132, 10209–10211; (b) L.-J. Xiao, L. Cheng, W.-M. Feng, M.-L. Li, J.-H. Xie and Q.-L. Zhou, Angew. Chem., Int. Ed., 2018, 57, 461–464; (c) C. Wang, Y. Guo, X. Wang, Z. Wang and K. Ding, Chem.–Eur. J., 2021, 27, 15903–15907.
- 8 For transition metal-catalyzed asymmetric hydro(hetero) arylation of 1,3-diene, see: (a) J. S. Marcum, T. R. Taylor and S. J. Meek, Angew. Chem., Int. Ed., 2020, 59, 14070–14075; (b) X.-Y. Lv, C. Fan, L.-J. Xiao, J.-H. Xie and Q.-L. Zhou, CCS Chem., 2019, 1, 328–334; (c) Y.-G. Chen, B. Shuai, X.-T. Xu, Y.-Q. Li, Q.-L. Yang, H. Qiu, K. Zhang, P. Fang and T.-S. Mei, J. Am. Chem. Soc., 2019, 141, 3395–3399; (d) J. S. Marcum, C. C. Roberts, R. S. Manan, T. N. Cervarich and S. J. Meek, J. Am. Chem. Soc., 2017, 139, 15580–15583; (e) S. M. Podhajsky, Y. Iwai, A. Cook-Sneathen and M. S. Sigman, Tetrahedron, 2011, 67, 4435–4441.
- (a) B. J. Stokes, A. J. Bischoff and M. S. Sigman, *Chem. Sci.*,
 2014, 5, 2336–2339; (b) B. J. Stokes, S. M. Opra and M. S. Sigman, *J. Am. Chem. Soc.*, 2012, 134, 11408–11411.
- 10 T. Hamaguchi, Y. Takahashi, H. Tsuji and M. Kawatsura, *Org. Lett.*, 2020, **22**, 1124–1129.
- 11 For a related arylative substitution of homoallylic alcohol derivatives to form linear products, see: K. Muto, T. Kumagai, F. Kakiuchi and T. Kochi, *Angew. Chem., Int. Ed.*, 2021, **60**, 24500–24504.
- 12 For reviews of transition metal-catalyzed allylic substitution of allylic alcohols, see: (a) D. Ghorai, À. Cristòfol and A. W. Kleij, Eur. J. Inorg. Chem., 2022, e202100820; (b) J. Qu and G. Helmchen, Acc. Chem. Res., 2017, 50, 2539–2555; (c) N. A. Butt and W. Zhang, Chem. Soc. Rev., 2015, 44, 7929–7967; (d) B. Sundararaju, M. Achard and C. Bruneau, Chem. Soc. Rev., 2012, 41, 4467–4483.
- 13 For selected direct nickel-catalyzed allylic arylation of allylic alcohols, see: (a) H. Yua and Z.-X. Wang, Org. Biomol. Chem., 2021, 19, 9723–9731; (b) Y.-B. Wang, B.-Y. Liu, Q. Bu, B. Dai and N. Liu, Adv. Synth. Catal., 2020, 362, 2930–2940; (c) S. H. Nazari, J. E. Bourdeau, M. R. Talley, G. A. Valdivia-Berroeta, S. J. Smith and D. J. Michaelis, ACS Catal., 2018, 8, 86–89; (d) X.-G. Jia, P. Guo, J. Duan and X.-Z. Shu, Chem. Sci., 2018, 9, 640–645.
- 14 For selected nickel-catalyzed allylic arylations of allylic alcohol derivatives, see: (a) X. Li, Y. Li, Z. Zhang, X. Shi, R. Liu, Z. Wang, X. Li and D. Shi, *Org. Lett.*, 2021, 23,

- 6612–6616; (*b*) J.-L. Tao, B. Yang and Z.-X. Wang, *J. Org. Chem.*, 2015, **80**, 12627–12634; (*c*) L. L. Anka-Lufford, M. R. Prinsell and D. J. Weix, *J. Org. Chem.*, 2012, 77, 9989–10000; (*d*) S. Wang, Q. Qian and H. Gong, *Org. Lett.*, 2012, **14**, 3352–3355; (*e*) Y. Dai, F. Wu, Z. Zang, H. You and H. Gong, *Chem.-Eur. J.*, 2012, **18**, 808–812.
- 15 For *in situ* activation of allylic alcohols using arylboronic acids, see: (a) D. G. Hall, *Chem. Soc. Rev.*, 2019, 48, 3475–3496; (b) H.-B. Wu, X.-T. Maa and S.-K. Tian, *Chem. Commun.*, 2014, 50, 219–221; (c) J. Ye, J. Zhao, J. Xu, Y. Mao and Y. J. Zhang, *Chem. Commun.*, 2013, 49, 9761–9763; (d) H. Tsukamoto, T. Uchiyama, T. Suzuki and Y. Kondo, *Org. Biomol. Chem.*, 2008, 6, 3005–3013.
- 16 The superiority of arylboroxines over other arylboron sources has been observed previously, see: R. Li, F. Liu and G. Dong, *Chem*, 2019, 5, 929–939.
- 17 L. Chen, H. Min, W. Zeng, X. Zhu, Y. Liang, G. Deng and Y. Yang, *Org. Lett.*, 2018, **20**, 7392–7395.
- 18 For examples of MeOH as a hydride source, see: (a) H. N. Tran and L. M. Stanley, *Org. Lett.*, 2022, 24, 395–399; (b) H. N. Tran, R. W. Burgett and L. M. Stanley, *J. Org. Chem.*, 2021, 86, 3836–3849; (c) Y. He, C. Liu, L. Yu and S. Zhu, *Angew. Chem., Int. Ed.*, 2020, 59, 9186–9191; (d) J.-F. Li, Z.-Z. Wei, Y.-Q. Wang and M. Ye, *Green Chem.*, 2017, 19, 4498–4502.
- 19 For examples of formations π-allyl-nickel(II) intermediates from allylic alcohols, see: (a) Y. Gan, H. Hu and Y. Liu, Org. Lett., 2020, 22, 4418–4423; (b) Y. Bernhard, B. Thomson, V. Ferey and M. Sauthier, Angew. Chem., Int. Ed., 2017, 56, 7460–7464; (c) M. S. Azizi, Y. Edder, A. Karim and M. Sauthier, Eur. J. Org. Chem., 2016, 3796–3803; (d) Y. Kita, H. Sakaguchi, Y. Hoshimoto, D. Nakauchi, Y. Nakahara, J.-F. Carpentier, S. Ogoshi and K. Mashima, Chem.–Eur. J., 2015, 21, 14571–14578; (e) H. Bricouta, E. Monflier, J.-F. Carpentier and A. Mortreux, Eur. J. Inorg. Chem., 1998, 1739–1744.
- 20 For examples of β-hydride elimination of metallacylces, see: (a) D. Jin, T. J. Schmeier, P. G. Williard, N. Hazari and W. H. Bernskoetter, *Organometallics*, 2013, 32, 2152–2159; (b) X. Huang, J. Zhu and Z. Lin, *Organometallics*, 2004, 23, 4154–4159.
- 21 (a) K. Farshadfar, A. Chipman, M. Hosseini, B. F. Yates and A. Ariafard, *Organometallics*, 2019, 38, 2953–2962; (b) J. Tan, Z. Zhang and Z. Wang, *Org. Biomol. Chem.*, 2008, 6, 1344–1348.
- 22 H. Iwamoto, T. Tsuruta and S. Ogoshi, ACS Catal., 2021, 11, 6741–6749.
- 23 For selected examples on the use of K₃PO₄ to facilitate the transmetallation of arylboroxines, see: (a) J. Zhu, L. Huang, W. Dong, N. Li, X. Yu, W.-P. Deng and W. Tang, *Angew. Chem., Int. Ed.*, 2019, 58, 16119–16123; (b) F. Pan, H. Wang, P.-X. Shen, J. Zhao and Z.-J. Shi, *Chem. Sci.*, 2013, 4, 1573–1577.
- 24 Y. Ding, J. Long, F. Sun and X. Fang, *Org. Lett.*, 2021, 23, 6073–6078.