

Cite this: *Chem. Sci.*, 2022, 13, 7007

All publication charges for this article have been paid for by the Royal Society of Chemistry

Hydroxy-directed fluorination of remote unactivated C(sp³)-H bonds: a new age of diastereoselective radical fluorination†

Stefan Andrew Harry,  Michael Richard Xiang, Eric Holt, Andrea Zhu, Fereshte Ghorbani, Dhaval Patel and Thomas Lectka*

We report a photochemically induced, hydroxy-directed fluorination that addresses the prevailing challenge of high diastereoselectivity in this burgeoning field. Numerous simple and complex motifs showcase a spectrum of regio- and stereochemical outcomes based on the configuration of the hydroxy group. Notable examples include a long-sought switch in the selectivity of the refractory sclareolide core, an override of benzylic fluorination, and a rare case of 3,3'-difluorination. Furthermore, calculations illuminate a low barrier transition state for fluorination, supporting our notion that alcohols are engaged in coordinated reagent direction. A hydrogen bonding interaction between the innate hydroxy directing group and fluorine is also highlighted for several substrates with ¹⁹F-¹H HOESY experiments, calculations, and more.

Received 2nd April 2022

Accepted 2nd May 2022

DOI: 10.1039/d2sc01907h

rsc.li/chemical-science

The hydroxy (OH) group is treasured and versatile in chemistry and biology.¹ Its ubiquity in nature and broad spectrum of chemical properties make it an attractive source as a potential directing group.² The exploitation of the mild Lewis basicity exhibited by alcohols has afforded several elegant pathways for selective functionalization (*e.g.*, Sharpless epoxidation,³ homogeneous hydrogenation,⁴ cross-coupling reactions,⁵ among others⁶). Recently, we reported a photochemically promoted carbonyl-directed aliphatic fluorination, and most notably, established the key role that C-H...O hydrogen bonds play in the success of the reaction.⁷ Our detailed mechanistic investigations prompt us to postulate that other Lewis basic functional groups (such as -OH) can direct fluorination in highly complementary ways.⁸ In this communication, we report a hydroxy-directed aliphatic fluorination method that exhibits unique directing properties and greatly expands the domain of radical fluorination into the less established realm governing high diastereoselectivity.⁹

Our first inclination that functional groups other than carbonyls may influence fluorination regiochemical outcomes was obtained while screening substrates for our published ketone-directed radical-based method (Scheme 1).^{8a} In this example, we surmised that oxidation of the tertiary hydroxy group on substrate **1** cannot occur and would demonstrate functional group tolerance (directing to C11, compound **2**). Surprisingly, the two major regioisomers (products **3** and **4**) are



derivatized by Selectfluor (SF) on C12 and C16 – indicative of the freely rotating hydroxyl directing fluorination. Without an obvious explanation of how these groups could be involved in dictating regiochemistry, we continued the mechanistic study of carbonyl-directed fluorination (Scheme 2A). We established that the regioselective coordinated hydrogen atom abstraction occurs by hydrogen bonding between a strategically placed carbonyl and Selectfluor radical dication (SRD).⁷ However, we noted that the subsequent radical fluorination is not diastereoselective due to the locally planar nature of carbonyl groups. Thus, we posed the question: are there other directing groups that can provide both regio- and diastereoselectivity? Such a group would optimally be attached to a sp³ hybridized carbon; thus the “three dimensional” hydroxy carbon logically comes to mind as an attractive choice, and Scheme 1 illustrates the first positive hint.

We began our detailed study with a simple substrate that contains a tertiary hydroxyl group. Alcohol **5** was synthesized stereoselectively by the reaction of 3-methylcyclohexanone, FeCl₃, and 4-chlorophenylmagnesium bromide;¹⁰ the 4-chlorophenyl substituent allows for an uncomplicated product identification and isolation (aromatic chromophore). We sought to determine optimal reaction conditions by examination of numerous photosensitizers, bases, solvents, and light sources (Table 1). To our satisfaction, fluorination not only provides the intended regioisomer but only a single diastereomer is formed (compound **6**). A photosensitizer screen shows that benzil (in MeCN solvent) generally affords the highest yield (83%).⁷ Although we utilize cool blue LEDs (sharp cutoff *ca.* 400 nm), CFLs (small amount of UVB (280–315 nm)

Department of Chemistry, Johns Hopkins University, 3400 N. Charles St., Baltimore, MD, 21218, USA. E-mail: lectka@jhu.edu

† Electronic supplementary information (ESI) available. See <https://doi.org/10.1039/d2sc01907h>



Table 2 Substrate scope^a





Fig. 1 Example of an intramolecular competition (secondary vs. tertiary C–H abstraction/fluorination) and calculated C–H...O distances of compound 5 (B3LYP/6-311++G**).

in coordinated hydrogen atom transfer (HAT) (55% and 40% for molecules 9 and 8) – a change in regiochemistry based on the stereochemistry of the alcohol. Additionally, only a single stereoisomer is produced for both (d.r. 99 : 1) and reinforce this study as a salient example of diastereoselective radical fluorination.

In the midst of characterizing compound 9, we uncovered a noteworthy hydrogen bonding interaction. Firstly, our plan was to identify the –OH peak within the ^1H NMR spectrum and determine if there is a through-space interaction with fluorine in the ^{19}F – ^1H HOESY NMR spectrum (ultimately aiding in assigning the stereochemistry of the fluorine).¹⁵ At first glance, no peaks were immediately discernible as the –OH; however, when a stoichiometric amount of H_2O is added, it becomes apparent that the –OH group and geminal proton to the hydroxy peaks broaden by rapid proton exchange (Fig. 2A). Upon closer examination of the dry ^1H NMR spectrum, the –OH peak appears to be a sharp doublet of doublets: one bond coupling to the geminal C–H proton of 9 Hz and one of the largest reported through-space couplings to fluorine of 20 Hz. The ^{19}F – ^1H HOESY spectrum also supports our regio- and stereochemical assignment – a strong interaction between fluorine and H_a , H_b , and H_d , as well as no apparent interaction with H_c and H_e (Fig. 2B). Consequently, we postulate that intramolecular hydrogen bonding is responsible for the considerable coupling constant. This conclusion is also supported by calculations at

alcohol directionality switch



–OH stereochemistry predicts regio- and stereoselectivity

Fig. 3 Examples of hydroxy group stereochemical switches.

B3LYP/6-311++G** (Fig. 2C): the O–H–F angle is given as 140° and $\text{F} \cdots \text{H} - \text{O}$ bond distance is 1.97 \AA .

Appreciating the complexity and biological significance of steroids,¹⁶ we derivatized dehydroepiandrosterone to afford fluorinated substrate 10 (42%; d.r. 99 : 1). Computational modeling assisted in verifying that the β -hydroxy group targets the C12 position (B3LYP/6-311++G**); furthermore, the β -fluoro isomer is the major product (validated by NOESY, ^1H , and ^{19}F NMR). Additionally, we subjected 17α -hydroxyprogesterone (endogenous progestogen steroid hormone¹⁷) to fluorination conditions and found the α -fluoro product (11) as the major diastereomer in 55% yield (99 : 1 d.r.). To investigate further the notion of coordinated fluorination and explanation of the observed stereoisomers (e.g., β -hydroxy/ β -fluoro and α -hydroxy/ α -fluoro), we calculated a simplified system comparing the fluorination of 1-propyl radical and γ -propanol radical (Scheme 2C). The reaction can be distilled into two key steps: a site-selective HAT, followed by a diastereoselective fluorination reaction. The following isodesmic relation ($\omega\text{B97xd}/6-31+\text{G}^*$, $-7.63 \text{ kcal mol}^{-1}$) illustrates the stabilizing energetic role that the hydroxy group plays in commanding diastereoselectivity. The transition states represent low barrier processes; a solvent dielectric was necessary to find saddle points.

Additionally, a simple Protein Data Bank (PDB) survey showed numerous intermolecular close contacts between hydroxy groups and $\text{H}-\text{C}^+-\text{NR}_3$ moieties.¹⁸ What is more,





Fig. 2 (A) Top spectrum (pink) has broadened peaks due to adventitious H_2O in solution. (B) Strong interaction observed between the installed fluorine and designated hydroxy proton in the ^{19}F – ^1H HOESY NMR spectrum. (C) Calculated structure for compound 9 at B3LYP/6-311++G** revealing the hydroxy proton aiming toward the fluorine.





Scheme 3 Isodesmic equation comparing carbonyl *versus* hydroxy group Selectfluor coordination.

solutions of Selectfluor with various alcohols at different concentrations reveal characteristic H-C-⁺NR₃ downfield chemical shifts in the ¹H NMR spectra (Scheme 2D).¹⁹ Both of these observations buttress the claim of a putative hydrogen bonding interaction between Selectfluor and the hydroxy group.

We theorize that the regioselective HAT step proceeds similarly to the reported carbonyl-directed pathway (Scheme 2A) involving Selectfluor radical cation coordination (considering the likenesses in conditions and aforementioned Lewis basicity logic). Alternatively, one can imagine the reaction proceeding through a Barton²⁰ or Hofmann-Löffler-Freytag²¹ style mechanism. To probe this possibility, we employed a β-caryophyllene alcohol hypochlorite derivative to form the alkoxy radical directly, and found that under standard conditions there is complex fragmentation and nonselective fluorination (Scheme 2B). Lastly, we compared the hydroxy *versus* carbonyl group SF coordination computationally. The carbonyl group is preferred to bind to SF through nonclassical C-H...O hydrogen bonds preferentially over the hydroxy group, as the following isodesmic relation shows (acetone and *t*-BuOH as models; $\omega\text{B97xd/6-31+G}^*$, -3.81 kcal mol⁻¹), but, once again, rigidity and propinquity are ultimately more important factors in determining directing effects (Scheme 3).

The tetrahedral nature of hydroxy groups provides unique access to previously unobtainable sites. For example, we compared menthol and an alkylated congener to form products **12** and **13** (Fig. 3B). The hydroxy group in the precursor to **12** is in the equatorial position, mandating the exocyclic isopropyl group as the reactive site (40% yield).²² In the precursor to **13**, the methyl and isopropyl substituent lock the hydroxy group


into the axial position, targeting its endocyclic tertiary site through a 1,3-diaxial relationship to afford fluorinated product in 57% yield (d.r. 99 : 1). In all, the comparison showcases the versatility in directing ability, offering a choice of regio- and stereoselectivity based on the stereochemistry of the hydroxy group. The directing system only necessitates two features based on our results: (1) the hydroxy group must be either secondary or tertiary (primary tends to favor oxidation) and (2) the oxygen atom must be within the range of 2.4–3.2 Å of the targeted secondary or tertiary hydrogen.

Among the several biologically active compounds we screened, caratol derivatives **14** and **15** were found to be attractive candidates that reveal directed fluorination to an exocyclic isopropyl group (Table 2). Comprising *ca.* 40% of carrot seed oil, caratol is its major constituent and has shown allelopathic interactions (*e.g.*, as insecticidal, antifungal, and herbicidal agents²³).²⁴ After extraction, isolation, and derivatization, molecules **14** and **15** are afforded in 65% and 83% yield (Table 2). Acetates and oxidized sulfur-containing functional groups were well tolerated.

Another natural product we modified to a tertiary alcohol was sclareolide, which has a rich history of fluorinating on its C2 and C3 positions due to polar effects, as reported by Tang,²⁵ Groves,²⁶ Britton,²⁶ and others.²⁷ The derived alcohol finally overrode this natural tendency and directed to the predicted position in 56% (d.r. 99 : 1) (product **16**). Smaller amounts of competitive polar effect fluorination were observed at the C2 and C3 positions, highlighting how challenging a problem the functionalization of the sclareolide core presents.^{28,29}

An altered dihydroactinidiolide was found to participate in the fluorination through a 1,3-diaxial guided HAT and fluorination in 55% yield (product **17**, d.r. 99 : 1). We next modeled several more substrates that participated in similar 1,3 relationships; however, each exhibited a variation from one another (*e.g.*, ring size or fused aromatic ring). Products **19** and **18** displayed the reaction's capability to direct to the desired positions with an expanded (65%; d.r. 99 : 1) and reduced (45%; d.r. 99 : 1) ring system when compared to the previous 6-membered ring examples. Additionally, we examined a methylated α-tetralone derivative. The desired 3-fluoro product **20** forms in 43% yield (d.r. 99 : 1), overriding benzylic fluorination (Scheme 4).³⁰ Under identical conditions α-tetralone provides 4-fluorotetralone in 48% yield. In similar motif, 1-phenylindanol, we intentionally targeted the benzylic position in a 90% and 10 : 1 d.r. (product **21**). Unlike the methylated α-tetralone derivative, the geometry of the starting material calculated at B3LYP/6-311++G** shows the hydroxy group is not truly axial and is 4.30 Å from the targeted C-H bond, explaining the dip in diastereoselectivity.

Next, we examined an isomer of borneol that is widely used in perfumery, fenchol.³¹ The secondary alcohol displays a diastereoselective fluorination in 38% (d.r. 99 : 1) (product **22**). Our last designed motif was ideally constructed to have a doubly-directing effect. Our observations show that a well-positioned hydroxy group not only provides sequential regioselective hydrogen atom abstraction but also displays a powerful demonstration of Selectfluor guidance to afford the *cis*-difluoro



Scheme 4 Comparing fluorination outcomes for different functional groups.



product (23) in 33% yield (85% brsm, d.r. 99 : 1). Spectroscopically (^1H , ^{13}C , and ^{19}F NMR), the product possesses apparent C_s symmetry and showcases close interactions (e.g., diagnostic couplings and chemical shifts). *cis*-Polyfluorocycloalkanes are of intense current interest in materials chemistry, wherein faces of differing polarity can complement one another.³²

All in all, this photochemical hydroxy-directed fluorination report represents one of the first steps in commanding diastereoselectivity within the field of radical fluorination. An ability to dictate regio- and stereoselectivity is demonstrated in a variety of substrates by simply switching the stereochemistry of the hydroxy group. Computations support the key role of Selectfluor coordination to the key hydroxy group in the fluorination step. Future studies will seek to uncover other compatible Lewis basic functional groups, expanding further the versatility of radical fluorination.

Data availability

Experimental and computational data is located in the ESI.†

Author contributions

S. Harry designed probes, synthesized compounds, and provided drafts of the manuscript. M. Xiang, E. Holt, A. Zhu, F. Ghorbani, and D. Patel synthesized compounds for isolation/characterization and assisted in compiling data. T. Lectka supervised the project and reviewed/edited the manuscript.

Conflicts of interest

The authors declare no competing financial interest.

Acknowledgements

T.L. thanks the National Science Foundation (NSF) (CHE grants 1800510 and 2102116) for financial support. Mass spectral data were obtained at the University of Delaware's mass spectrometry center.

References

- (a) P. Chauhan and S. S. Chimni, *RSC Adv.*, 2012, **2**, 737–758; (b) G. D. Rose and R. Wolfenden, *Annu. Rev. Biophys. Biomol. Struct.*, 1993, **22**, 381–415; (c) X. Wu and C. Zhu, *Chem. Commun.*, 2019, **55**, 9747–9756; (d) T. Steiner, *Crystallogr. Rev.*, 2003, **9**, 177–228; (e) S. Zhang, F. Zhang and Y. Tu, *Chem. Soc. Rev.*, 2011, **4**, 1937–1949; (f) A. S. Kertes and C. J. King, *Chem. Rev.*, 1987, **87**, 687–710; (g) W. Kroutil, H. Mang, K. Edegger and K. Faber, *Adv. Synth. Catal.*, 2004, **2**, 125–142.
- (a) C. Zhang, J. Ji and P. Sun, *J. Org. Chem.*, 2014, **79**, 3200–3205; (b) G. T. Hoang, Z. Pan, J. T. Brethorst and C. J. Douglas, *J. Org. Chem.*, 2014, **79**, 11383–11394; (c) A. M. Camelio, T. Barton, F. Guo, T. Shaw and D. Siegel, *Org. Lett.*, 2011, **13**, 1517–1519; (d) J. Sun, G. Zheng, T. Xiong, Q. Zhang, J. Zhao, Y. Li and Q. Zhang, *ACS Catal.*, 2016, **6**, 3674–3678; (e) X. Wu, H. Zhang, N. Tang, Z. Wu, D. Wang, M. Ji, Y. Xu, M. Wang and C. Zhu, *Nat. Commun.*, 2018, **9**, 3343.
- (a) T. Katsuki and K. B. Sharpless, *J. Am. Chem. Soc.*, 1980, **102**, 5976–5978; (b) H. Brunner and A. Sicheneder, *Angew. Chem., Int. Ed. Engl.*, 1988, **27**, 718–719; (c) V. Jäger, D. Schröter and B. Köppenhoefer, *Tetrahedron*, 1991, **47**, 2195–2210.
- (a) J. Brown, *Angew. Chem., Int. Ed. Engl.*, 1987, **26**, 190–203; (b) R. H. Crabtree and M. W. Davis, *Organometallics*, 1983, **2**, 682–684; (c) R. H. Crabtree and M. W. Davis, *J. Org. Chem.*, 1986, **51**, 2655–2661; (d) G. Stork and D. E. Kahne, *J. Am. Chem. Soc.*, 1983, **105**, 1072–2073.
- (a) T. P. Blaisdell and J. P. Morken, *J. Am. Chem. Soc.*, 2015, **137**, 8712–8715; (b) K. Manabe and S. Ishikawa, *Synthesis*, 2008, **16**, 2645–2649; (c) E. M. Simmons and J. F. Hartwig, *J. Am. Chem. Soc.*, 2010, **132**, 17092–17095.
- (a) E. M. Simmons and J. F. Hartwig, *Nature*, 2012, **483**, 70–73; (b) X. Wu, M. Wang, L. Huan, D. Wang, J. Wang and C. Zhu, *Angew. Chem., Int. Ed. Engl.*, 2018, **57**, 1640–1644; (c) T. P. Blaisdell, T. C. Caya, L. Zhang, A. Sanz-Marco and J. P. Morken, *J. Am. Chem. Soc.*, 2014, **136**, 9264–9267; (d) X. Wang, Y. Lu, H. Dai and J. Yu, *J. Am. Chem. Soc.*, 2010, **132**, 12203–12205; (e) Y. Lu, D. Wang, K. M. Engle and J. Yu, *J. Am. Chem. Soc.*, 2010, **132**, 5916–5921; (f) S. Saito, T. Ishikawa, A. Kuroda, K. Koga and T. Moriwake, *Tetrahedron*, 1992, **48**, 4067–4086.
- F. Ghorbani, S. A. Harry, J. N. Capilato, C. R. Pitts, J. Joram, G. Peters, J. Tovar, I. Smajlagic, M. Siegler, T. Dudding and T. Lectka, *J. Am. Chem. Soc.*, 2020, **142**, 14710–14724.
- (a) D. D. Bume, C. R. Pitts, F. Ghorbani, S. A. Harry, J. N. Capilato, M. A. Siegler and T. Lectka, *Chem. Sci.*, 2017, **8**, 6918–6923; (b) D. D. Bume, S. A. Harry, C. R. Pitts and T. Lectka, *J. Org. Chem.*, 2018, **83**, 1565–1575; (c) C. R. Pitts, D. D. Bume, S. A. Harry, M. A. Sielger and T. Lectka, *J. Am. Chem. Soc.*, 2017, **139**, 2208–2211.
- (a) Z. Yuan, H. Wang, X. Mu, P. Chen, Y. Guo and G. Liu, *J. Am. Chem. Soc.*, 2015, **137**, 2468–2471; (b) W. Zhang, Y. Gu, J. Lin and J. Xiao, *Org. Lett.*, 2020, **22**, 6642–6646; (c) Z. Li, Z. Wang, L. Zhu, X. Tan and C. Li, *J. Am. Chem. Soc.*, 2014, **136**, 16439–16443; (d) H. Shigehisa, E. Nishi, M. Fujisawa and K. Hiroya, *Org. Lett.*, 2013, **15**, 5158–5161; (e) W. Liu, X. Huang and J. T. Groves, *Nat. Protoc.*, 2013, **8**, 2348–2354; (f) W. Liu, X. Huang, M. J. Cheng, R. J. Nielsen, W. A. Goddard III and J. T. Groves, *Science*, 2012, **337**, 1322–1325.
- M. T. Reetz and S. Stanchev, *J. Chem. Soc., Chem. Commun.*, 1993, **3**, 328–330.
- G. A. Olah, *Angew. Chem.*, 1973, **85**, 183–225.
- (a) Y. Gu, T. Kar and S. Scheiner, *J. Am. Chem. Soc.*, 1999, **121**, 9411–9422; (b) S. G. Olesen and S. Hammerum, *J. Phys. Chem. A*, 2009, **113**, 7940–7944; (c) X. Wang, H. Woo, B. Kiran and L. Wang, *Angew. Chem.*, 2005, **117**, 5048–5052.
- (a) S. P. Bhatia, C. S. Letizia and A. M. Api, *Food Chem. Toxicol.*, 2008, **46**, 95–96; (b) M. Skold, A. Karlberg, M. Matura and A. Börje, *Food Chem. Toxicol.*, 2006, **44**, 538–545.



- 14 G. G. Henderson, R. Osborne, O. McCrone and J. M. Robertson, *J. Am. Chem. Soc.*, 1929, 1368–1372.
- 15 L. E. Combettes, P. Clausen-Thue, M. A. King, B. Odell, A. L. Thompson, V. Gouverneur and T. D. W. Claridge, *Chem.–Eur. J.*, 2012, 12, 13133–13141.
- 16 (a) S. L. Greenspan, D. S. Oppenheim and A. Klibanski, *Ann. Intern. Med.*, 1989, 110, 526–531; (b) J. M. Tanner, R. H. Whitehouse, P. C. R. Hughes and B. S. Carter, *J. Pediatr.*, 1976, 89, 1000–1008; (c) A. Y. Bagrov, J. I. Shapiro and O. V. Fedorova, *Pharmacol. Rev.*, 2009, 61, 9–38.
- 17 D. Tulchinsky and C. J. Hobel, *Am. J. Obstet. Gynecol.*, 1973, 117, 884–893.
- 18 Y. Itoh, Y. Nakashima, S. Tsukamoto, T. Kurohara, M. Suzuki, Y. Sakae, M. Oda, Y. Okamoto and T. Suzuki, *Sci. Rep.*, 2019, 9, 1–12.
- 19 J. Capilato, S. A. Harry, M. Seigler and T. Lectka, *Chem.–Eur. J.*, 2021, 28, e202103922.
- 20 D. H. Barton, R. Hesse, M. Pechet and L. Smith, *J. Chem. Soc., Perkin Trans. 1*, 1979, 115, 1159–1165.
- 21 G. Majetich and K. Wheless, *Tetrahedron*, 1995, 55, 7095–7129.
- 22 E. Hayashi, S. Hara, H. Shirato, T. Hatekeyama, T. Fukuhara and N. Yoneda, *Chem. Lett.*, 1995, 3, 205–206.
- 23 A. Cakir, H. Özer, T. Aydin, S. Kordali, A. T. Cavusoglu, T. Akcin, E. Mete and A. Akcin, *Rec. Nat. Prod.*, 2016, 10, 154–167.
- 24 S. Sridhar, R. V. Rajagopal, R. Rajavel, S. Masilamani and S. Narasimhan, *J. Agric. Food Chem.*, 2003, 51, 7596–7599.
- 25 X. Zhang, S. Guo and P. Tang, *Org. Chem. Front.*, 2015, 2, 806–810.
- 26 S. D. Halperin, H. Fan, S. Chang, R. E. Martin and R. Britton, *Angew. Chem.*, 2014, 126, 4778–4781.
- 27 (a) Y. Takahira, M. Chen, Y. Kawamata, P. Mykhailiuk, H. Nakamura, M. K. Peters, S. H. Reisberg, C. Li, L. Chen, T. Hoshikawa, T. Shibuguchi and P. S. Baran, *Synlett*, 2019, 30, 1178–1182; (b) S. Bloom, J. L. Knippel and T. Lectka, *Chem. Sci.*, 2014, 5, 1175–1178.
- 28 (a) C. Walling, *Free Radicals in Solution*, Wiley, New York, NY, 1957; (b) R. Bernardi, R. Galli and F. Minisci, *J. Chem. Soc. B*, 1968, 324–325; (c) F. Minisci, R. Galli and R. Bernardi, *Chem. Commun.*, 1967, 17, 903–904; (d) A. A. Zavitsas and J. A. Pinto, *J. Am. Chem. Soc.*, 1972, 94, 7390–7396; (e) T. Newhouse and P. S. Baran, *Angew. Chem., Int. Ed.*, 2011, 50, 3362–3374.
- 29 (a) Y. Amaoka, M. Nagamoto and M. Inoue, *Org. Lett.*, 2013, 15, 2160–2163; (b) C. R. Pitts, B. Ling, R. Woltornist, R. Liu and T. Lectka, *J. Org. Chem.*, 2014, 79, 8895–8899; (c) S. Bloom, C. R. Pitts, D. Miller, N. Haselton, M. G. Holl, E. Urheim and T. Lectka, *Angew. Chem., Int. Ed.*, 2012, 51, 10580–10583; (d) J.-B. Xia, C. Zhu and C. Chen, *Chem. Commun.*, 2014, 50, 11701–11704; (e) J. G. West, T. A. Bedell and E. J. Sorensen, *Angew. Chem., Int. Ed.*, 2016, 55, 8923–8927; (f) J.-B. Xia, Y. Ma and C. Chen, *Org. Chem. Front.*, 2014, 1, 468–472; (g) C. W. Kee, K. F. Chin, M. W. Wong and C.-H. Tan, *Chem. Commun.*, 2014, 50, 8211–8214; (h) S. D. Halperin, H. Fan, S. Chang, R. E. Martin and R. Britton, *Angew. Chem., Int. Ed.*, 2014, 53, 4690–4693; (i) P. Champagne, J. Desroches, J. Hamel, M. Vandamme and J.-F. Paquin, *Chem. Rev.*, 2015, 115, 9073–9174.
- 30 (a) A. M. Hua, D. N. Mai, R. Martinez and R. D. Baxter, *Org. Lett.*, 2017, 19, 2949–2952; (b) D. Cantillo, O. de Frutos, J. A. Rincon, C. Mateos and O. C. Kappe, *J. Org. Chem.*, 2014, 79, 8486–8490; (c) S. Bloom, C. R. Pitts, R. Woltornist, A. Griswold, M. G. Holl and T. Lectka, *Org. Lett.*, 2013, 15, 1722–1724; (d) W. Liu and J. T. Groves, *Angew. Chem., Int. Ed.*, 2013, 52, 6024–6027; (e) S. Bloom, S. A. Sharber, M. G. Holl, J. L. Knippel and T. Lectka, *J. Org. Chem.*, 2013, 78, 11082–11086; (f) J.-B. Xia, C. Zhu and C. Chen, *J. Am. Chem. Soc.*, 2013, 135, 17494–17500; (g) S. Bloom, M. McCann and T. Lectka, *Org. Lett.*, 2014, 16, 6338–6341; (h) M. B. Nodwell, A. Bagai, S. D. Halperin, R. E. Martin, H. Knust and R. Britton, *Chem. Commun.*, 2015, 51, 11783–11786; (i) D. D. Bume, C. R. Pitts, R. T. Jokhai and T. Lectka, *Tetrahedron*, 2016, 72, 6031–6036; (j) A. Koperniku, H. Liu and P. B. Hurley, *Eur. J. Org. Chem.*, 2016, 871–886.
- 31 G. K. Oloyede, *Arabian J. Chem.*, 2016, 9, 840–845.
- 32 (a) M. Zupan, A. Gregorcic and A. Pollak, *J. Org. Chem.*, 1976, 42, 1562–1565; (b) M. Kaj, S. Banik and E. Jacobsen, *Org. Lett.*, 2019, 21, 4919–4923; (c) Q. Huchet, B. Kuhn, B. Wagner, N. Kratochwil, H. Fischer, M. Kansy, D. Zimmerli, E. Carreira and K. Müller, *J. Med. Chem.*, 2015, 58, 9041–9060; (d) C. Thiehoff, Y. Rey and R. Gilmour, *Isr. J. Chem.*, 2017, 57, 92–100; (e) I. Yamamoto, M. Jordan, N. Gavande, M. Doddareddy, M. Chebib and L. Hunter, *Chem. Commun.*, 2012, 48, 829–831; (f) X. Hu, D. Thomas, R. Griffith and L. Hunter, *Angew. Chem., Int. Ed.*, 2014, 53, 6176–6179; (g) M. Schüler, D. O'Hagan and A. Slawin, *Chem. Commun.*, 2005, 4324–4326; (h) I. G. Molnar, C. Thiehoff, M. C. Holland and R. Gilmour, *ACS Catal.*, 2016, 6, 7167–7173.

