

Cite this: *Chem. Sci.*, 2022, 13, 14382

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 3rd September 2022  
Accepted 4th November 2022

DOI: 10.1039/d2sc04909k

rsc.li/chemical-science

# Selective editing of a peptide skeleton *via* C–N bond formation at N-terminal aliphatic side chains†

Yujie Han, Junjie Shi, Songrong Li, Tingting Dan, Wenwen Yang and Mingyu Yang \*

The applications of peptides and peptidomimetics have been demonstrated in the fields of therapeutics, diagnostics, and chemical biology. Strategies for the direct late-stage modification of peptides and peptidomimetics are highly desirable in modern drug discovery. Transition-metal-catalyzed C–H functionalization is emerging as a powerful strategy for late-stage peptide modification that is able to construct functional groups or increase skeletal diversity. However, the installation of directing groups is necessary to control the site selectivity. In this work, we describe a transition metal-free strategy for late-stage peptide modification. In this strategy, a linear aliphatic side chain at the peptide N-terminus is cyclized to deliver a proline skeleton *via* site-selective  $\delta$ -C(sp<sup>3</sup>)–H functionalization under visible light. Natural and unnatural amino acids are demonstrated as suitable substrates with the transformations proceeding with excellent regio- and stereo-selectivity.


## Introduction

Late-stage modification of peptides and peptidomimetics exhibited its potential utility in the field of therapeutic and diagnostic agents, drug candidates, and biological tool compounds (Scheme 1).<sup>1–6</sup> Although peptides are abundant in nature, non-natural peptides often demonstrate improved biological and pharmaceutical properties compared with their parent analogs. Therefore, straightforward and efficient methods for the late-stage modification of diverse peptides are highly desirable. To date, various methodologies,<sup>7–17</sup> including methylation,<sup>12</sup> phosphorylation,<sup>13</sup> S-alkylation/arylation,<sup>14,15</sup> and glycosylation<sup>16</sup> of polar functional groups, have been revealed for the successful chemical modification of natural peptides. Inspired by the development of direct C–H bond functionalization methods, site-specific C–H functionalization has recently emerged as a valuable tool for the late-stage modification of peptides and proteins.<sup>18–38</sup> For examples, electrophilic aromatic substitution,<sup>18,19</sup> Rh-catalyzed carbenoid insertion,<sup>21</sup> Pd-catalyzed arylation/alkenylation,<sup>22–26</sup> and photo-induced His-alkylation<sup>27</sup> have all been used in this capacity.

The modification of aliphatic side chains is a greater challenge than the modification of aromatic residues. In pioneering work by Li and coworkers, copper-catalyzed alkynylation of the  $\alpha$ -C–H bond of *N*-aryl substituted glycine was used to selectively modify short peptides in a racemic fashion.<sup>39</sup> Wang and Xu reported the alkylation of *N*-phenyl glycine residues in peptides *via* the decarboxylation of *N*-hydroxyphthalimide esters using a copper

photocatalyst.<sup>40</sup> Shi and coworkers reported a chemoselective peptide modification strategy involving a photocatalytic tryptophan  $\beta$ -position conjugation;<sup>41</sup> oxidation of indole nitrogen initiates a single-electron transfer from the tryptophan  $\beta$ -position to achieve the conjugation. However, these methods are mainly limited to weak C–H bonds, such as benzylic C–H bonds or those adjacent to a heteroatom (Scheme 2a).

Recently, directing group assisted transition-metal catalyzed functionalization of inert C(sp<sup>3</sup>)–H bonds within aliphatic residues of peptides has shown enormous potential, with applications in biochemistry and drug discovery.<sup>44–59</sup> Innovative contributions have been made by the groups of Yu,<sup>42,43</sup> Ackermann,<sup>44–46</sup> and Shi.<sup>47,48</sup> Both radical-induced and directing group assisted transition-metal catalyzed modifications functionalize the peptide at a single site (*i.e.*, point modifications) (Scheme 2b). To improve the skeletal diversity of peptides (Scheme 2c), Noisier/Albericio<sup>49</sup>



Scheme 1 Natural peptides/peptidomimetics.

Key Laboratory of Applied Surface and Colloid Chemistry of MOE & School of Chemistry and Chemical Engineering Shaanxi Normal University, 620 West Chang'an Ave, Xi'an, 710119, China. E-mail: yangmy05@snnu.edu.cn

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2sc04909k>



<sup>a</sup> Reaction was conducted on a 0.1 mmol scale. <sup>b</sup> Yield was determined by <sup>1</sup>H NMR. <sup>c</sup> Isolated yield in parentheses.

<sup>a</sup> Unless otherwise noted, the reaction conditions were as follows: **SM** (0.2 mmol), trichloroisocyanuric acid (TCCA, 1.5 equiv.), DCM 2.0 mL (*c* = 0.1 M), 0 °C, and 8 h; **1** (0.1 mmol), DCE 1.0 mL, blue LED (100 W), 45 °C, and 9 h; then NaOH (1.2 equiv.), CH<sub>3</sub>CN (2.0 mL), rt, and 5 h. <sup>b</sup> 2.5 mmol of **SM** (*n*) was used, *c* = 0.1 M. <sup>c</sup> In parentheses was the calculated total yield. <sup>d</sup> *dr* was determined by crude <sup>1</sup>H NMR. <sup>e</sup> Dichloro-product **1j** was obtained.



dipeptide **SMj** was used as a substrate, the cyclization reaction occurred exclusively at the N-terminal position; the lysine side chain was untouched by the reaction, thereby demonstrating the unique selectivity of this transformation (**2j**). Furthermore, the aliphatic residues at the N-terminal positions were also examined (**SMo–SMq**). Cyclization occurred at both primary and secondary C–H bonds, selectively yielding the corresponding proline skeleton derivatives. Significantly, the linear unnatural amino acid 2-aminopentanoic acid was converted into a natural proline derivative, albeit in moderate yield, indicating the power of this method (**2r**).

Encouraged by the successful results obtained for dipeptides, the application of the reaction to oligopeptides consisting of three to five amino acid units was explored (Table 3, **SMs–z**, and **SMAa–ae**). Because of the larger number of peptide bonds, selective modification of longer peptides is more challenging. However, for all peptides, the transformation proceeded in moderate to good yield with excellent diastereoselectivity under the standard conditions (**2s–2z** and **2aa–2ae**). Unprotected serine or threonine moieties adjacent to the N-terminal position were not detrimental to the reaction (**2w** and **2x**). Furthermore, the phenylalanine moiety, with its highly active benzylic C–H bonds, was untouched during the transformation (**2v**).

To further understand the reactivity of this catalytic system, the reactivity of an isoleucine derivative **Ile–NCl** and the isoleucine-derived dipeptide **1a** was compared. After irradiation with visible light, dipeptide **1a** was successfully converted to the corresponding product **2a'** in quantitative yield, whereas the amino acid only yielded the corresponding product **Ile–Cl** in 56% yield; this highlights the superior reactivity of peptide substrates over single amino acids (Scheme 3a, details in the ESI†).<sup>58</sup> Such a superior reactivity of peptides may have resulted from the hydrogen bonding between amide and chloride in dipeptides, which could assist the C–Cl bond formation.<sup>59</sup> Finally, gram-scale experiments were conducted to evaluate the scalability of the transformation. Within two tripeptide substrates, the N-terminal amino acid moiety was transformed into the corresponding cyclic proline derivatives in acceptable yields (Scheme 3b).

Based on our previous studies,<sup>52–55</sup> and other reported research,<sup>60</sup> we considered that such transformation proceeded through a 1,5-hydrogen atom transfer (1,5-HAT) process as the Hofmann–Löffler reaction reported.<sup>61,62</sup> Under the irradiation of visible light, the homolytic cleavage of the N–Cl bond occurred to generate a nitrogen radical and chloride radical (trapped by TEMPO and detected by ESI-MS, see the ESI†). Subsequently, 1,5-HAT takes place yielding the carbon-centered radical, which could be trapped by the chloride radical to form a C–Cl bond. Then, an intramolecular nucleophilic substitution occurred under the base conditions to form the target product.

## Conclusions

In summary, we have developed a visible light-promoted selective editing of peptide skeletons *via* C–N bond formation at N-terminal aliphatic side chains. A proline skeleton was constructed in peptides under such transition metal free conditions. The excellent diastereoselectivity and broad substrate scope of the reaction

highlight its potential utility in biochemistry, medicinal chemistry, and/or drug discovery.

## Author contributions

Y. H. and S. L. performed the experiments and analyzed the data. J. S., T. D. and W. Y. assisted the purification of compounds and analysis of data. M. Y. designed and directed the whole project and wrote the manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We gratefully acknowledge the financial support by the National Natural Science Foundation of China (22171175), Fundamental Research Funds for the central Universities (GK202103049), and start-up funds from Shaanxi Normal University, and Shaanxi Provincial Natural Science Foundation (2021JM-195).

## Notes and references

- 1 K. L. Dunbar, D. H. Scharf, A. Litomska and C. Hertweck, *Chem. Rev.*, 2017, **117**, 5521–5577.
- 2 S. Sato and H. Nakamura, *Molecules*, 2019, **24**, 3980–3996.
- 3 J. C. Lewis, P. S. Coelho and F. H. Arnold, *Chem. Soc. Rev.*, 2011, **40**, 2003–2021.
- 4 A. Henninot, J. C. Collins and J. M. Nuss, *Med. Chem.*, 2018, **61**, 1382–1414.
- 5 J. Lu, Y. Li, Z. Bai, H. Lv and H. Wang, *Nat. Prod. Rep.*, 2021, **38**, 981–992.
- 6 S. Xu, Z. Zhao and J. Zhao, *Chin. Chem. Lett.*, 2018, **29**, 1009–1016.
- 7 T. Morioka, A. Nishizawa, K. Nakamura, M. Tobisu and N. Chatani, *Chem. Lett.*, 2015, **44**, 1729–1731.
- 8 J. N. deGruyter, L. R. Malins and P. S. Baran, *Biochemistry*, 2017, **56**, 3863–3873.
- 9 C. D. Spicer and B. G. Davis, *Nat. Commun.*, 2014, **5**, 4740.
- 10 L. R. Malins, *Curr. Opin. Chem. Biol.*, 2018, **46**, 25–32.
- 11 O. Boutureira and G. J. L. Bernardes, *Chem. Rev.*, 2015, **115**, 2174–2195.
- 12 K. K. Biggar and S. S. Li, *Nat. Rev. Mol. Cell Biol.*, 2015, **16**, 5–17.
- 13 R. Klingberg, J. O. Jost, M. Schumann, K. A. Gelato, W. Fischle, E. Krause and D. Schwarzer, *ACS Chem. Biol.*, 2015, **10**, 138–145.
- 14 D. Wang, M. Yu, N. Liu, C. Lian, Z. Hou, R. Wang, R. Zhao, W. Li, Y. Jiang, X. Shi, S. Li, F. Yin and Z. Li, *Chem. Sci.*, 2019, **10**, 4966–4972.
- 15 E. V. Vinogradova, C. Zhang, A. M. Spokoyny, B. L. Pentelute and S. L. Buchwald, *Nature*, 2015, **526**, 687–691.
- 16 D. P. Gamblin, E. M. Scanlan and B. G. Davis, *Chem. Rev.*, 2009, **109**, 131–163.
- 17 M. Yang, X. Wang and J. Zhao, *ACS Catal.*, 2020, **10**, 5230–5235.





- 18 A. K. Mishra, R. Tessier, D. P. Hari and J. Waser, *Angew. Chem., Int. Ed.*, 2021, **60**, 17963–17968.
- 19 M. W. Jones, G. Mantovani, C. A. Blindauer, S. M. Ryan, X. Wang, D. J. Brayden and D. M. Haddleton, *J. Am. Chem. Soc.*, 2012, **134**, 7406–7413.
- 20 A. E. Mangubat-Medina, S. C. Martin, K. Hanaya and Z. T. Ball, *J. Am. Chem. Soc.*, 2018, **140**, 8401–8404.
- 21 J. M. Antos and M. B. Francis, *J. Am. Chem. Soc.*, 2004, **126**, 10256–10257.
- 22 Q. Bai, Z. Bai and H. Wang, *Org. Lett.*, 2019, **21**, 8225–8228.
- 23 X. Li, L. Qi, B. Li, Z. Zhao, G. He and G. Chen, *Org. Lett.*, 2020, **22**, 6209–6213.
- 24 J. Tan, J. Wu, S. Liu, H. Yao and H. Wang, *Sci. Adv.*, 2019, **5**, eaaw0323.
- 25 J. Tang, H. Chen, Y. He, W. Sheng, Q. Bai and H. Wang, *Nat. Commun.*, 2018, **9**, 3383.
- 26 Z. Bai and H. Wang, *Synlett*, 2020, **31**, 199–204.
- 27 X. Chen, F. Ye, X. Luo, X. Liu, J. Zhao, S. Wang, Q. Zhou, G. Chen and P. Wang, *J. Am. Chem. Soc.*, 2019, **141**, 18230–18237.
- 28 J. Ruiz-Rodríguez, F. Albericio and R. Lavilla, *Chem.–Eur. J.*, 2010, **16**, 1124–1127.
- 29 L. Mende-Tapia, S. Preciado, J. García, R. Ramón, N. Kielland, F. Albericio and R. Lavilla, *Nat. Commun.*, 2015, **6**, 7160.
- 30 W. Gong, G. Zhang, T. Liu, R. Giri and J. Q. Yu, *J. Am. Chem. Soc.*, 2014, **136**, 16940–16946.
- 31 B.-B. Zhan, Y. Li, J.-W. Xu, X.-L. Nie, J. Fan, L. Jin and B.-F. Shi, *Angew. Chem., Int. Ed.*, 2018, **57**, 5858–5862.
- 32 Y. Zhu, M. Bauer, J. Ploog and L. Ackermann, *Chem.–Eur. J.*, 2014, **20**, 13099–13102.
- 33 J. Tang, Y. He, H. Chen, W. Sheng and H. Wang, *Chem. Sci.*, 2017, **8**, 4565–4570.
- 34 Y. Zheng and W. Song, *Org. Lett.*, 2019, **21**, 3257–3260.
- 35 M. Yang, X. Jiang and Z.-J. Shi, *Org. Chem. Front.*, 2015, **2**, 51–54.
- 36 E. Hernando, J. Villalva, Á. M. Martínez, I. Alonso, N. Rodríguez, R. G. Arrayás and J. C. Carretero, *ACS Catal.*, 2016, **6**, 6868–6882.
- 37 B. Li, X. Li, B. Han, Z. Chen, X. Zhang, G. He and G. Chen, *J. Am. Chem. Soc.*, 2019, **141**, 9401–9407.
- 38 T. J. Osberger, D. C. Rogness, J. T. Kohrt, A. F. Stepan and M. C. White, *Nature*, 2016, **537**, 214–219.
- 39 L. Zhao, O. Baslé and C. J. Li, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 4106–4111.
- 40 C. Wang, R. Qi, H. Xue, Y. Shen, M. Chang, Y. Chen, R. Wang and Z. Xu, *Angew. Chem., Int. Ed.*, 2020, **59**, 7461–7466.
- 41 Y. Yu, L.-K. Zhang, A. V. Buevich, G. Li, H. Tang, P. Vachal, S. L. Colletti and Z.-C. Shi, *J. Am. Chem. Soc.*, 2018, **140**, 6797–6800.
- 42 W. Gong, G. Zhang, T. Liu, R. Giri and J.-Q. Yu, *J. Am. Chem. Soc.*, 2014, **136**, 16940–16946.
- 43 T. Liu, J.-X. Qiao, M. A. Poss and J.-Q. Yu, *Angew. Chem., Int. Ed.*, 2017, **56**, 10924–10927.
- 44 M. Bauer, W. Wang, M. M. Lorion, C. Dong and L. Ackermann, *Angew. Chem., Int. Ed.*, 2018, **57**, 203–207.
- 45 W. Wang, M. M. Lorion, O. Martinazzoli and L. Ackermann, *Angew. Chem., Int. Ed.*, 2018, **57**, 10554–10558.
- 46 J. Wu, N. Kaplaneris, S. Ni, F. Kaltenhäuser and L. Ackermann, *Chem. Sci.*, 2020, **11**, 6521–6526.
- 47 B.-B. Zhan, J. Fan, L. Jin and B.-F. Shi, *ACS Catal.*, 2019, **9**, 3298–3303.
- 48 J.-W. Xu, Z.-Z. Zhang, W.-H. Rao and B.-F. Shi, *J. Am. Chem. Soc.*, 2016, **138**, 10750–10753.
- 49 A. F. M. Noisier, J. García, I. A. Ionut and F. Albericio, *Angew. Chem., Int. Ed.*, 2017, **56**, 314–318.
- 50 J. Tang, Y. He, H. Chen, W. Sheng and H. Wang, *Chem. Sci.*, 2017, **8**, 4565–4570.
- 51 X. Zhang, G. Lu, M. Sun, M. Mahankali, Y. Ma, M. Zhang, W. Hua, Y. Hu, Q. Wang, J. Chen, G. He, X. Qi, W. Shen, P. Liu and G. Chen, *Nat. Chem.*, 2018, **10**, 540–548.
- 52 M. Yang, B. Su, Y. Wang, K. Chen, X. Jiang, Y.-F. Zhang, X.-S. Zhang, G. Chen, Y. Cheng, Z. Cao, Q.-Y. Guo, L. Wang and Z.-J. Shi, *Nat. Commun.*, 2014, **5**, 4707.
- 53 D. Meng, Y. Tang, J. Wei, X. Shi and M. Yang, *Chem. Commun.*, 2017, **53**, 5744–5747.
- 54 Y. Tang, Y. Qin, D. Meng, C. Li, J. Wei and M. Yang, *Chem. Sci.*, 2018, **9**, 6374–6378.
- 55 Y. Qin, Y. Han, Y. Tang, J. Wei and M. Yang, *Chem. Sci.*, 2020, **11**, 1276–1282.
- 56 C. Che, Y.-N. Li, X. Cheng, Y.-N. Lu and C.-J. Wang, *Angew. Chem., Int. Ed.*, 2021, **60**, 4698–4704.
- 57 W. Wang, J. Wu, R. Kuniyil, A. Kopp, R. N. Lima and L. Ackermann, *Chem.*, 2020, **6**, 3428–3439.
- 58 Q. Qin and S. Yu, *Org. Lett.*, 2015, **17**, 1894–1897.
- 59 K. J. Winstanley and D. K. Smith, *J. Org. Chem.*, 2007, **72**, 2803–2815.
- 60 A. E. Bosnidou, T. Duhamel and K. Muñiz, *Eur. J. Org. Chem.*, 2020, **40**, 6361–6365.
- 61 A. W. Hofmann, *Chem. Ber.*, 1883, **16**, 558–560.
- 62 K. Löffler and C. Freytag, *Chem. Ber.*, 1909, **42**, 3427–3431.

