


 Cite this: *Soft Matter*, 2022, 18, 7897

Correction: Interpenetration of fractal clusters drives elasticity in colloidal gels formed upon flow cessation

 Noémie Dagès,^a Louis V. Bouthier,^b Lauren Matthews,^c Sébastien Manneville,^a Thibaut Divoux,^a Arnaud Poulesquen^d and Thomas Gibaud^{id}*^a

DOI: 10.1039/d2sm90130g

rsc.li/soft-matter-journal

 Correction for 'Interpenetration of fractal clusters drives elasticity in colloidal gels formed upon flow cessation' by Noémie Dagès *et al.*, *Soft Matter*, 2022, 18, 6645–6659, <https://doi.org/10.1039/D2SM00481J>.

The authors regret a mistake in the derivation of the interpenetration ϕ -power law model in the published article. This mistake appears in eqn (8) and has repercussions on the final equation giving the gel elasticity G'_∞ in eqn (1) and (15). The corrected calculation of G'_∞ corroborates the experimental data presented in Fig. 4c of the published article. The paper's conclusion remains therefore unchanged.

In the published article, in eqn (8), the conservation of mass is expressed as

$$\phi \left(\frac{\xi_s}{r_0} \right)^{3-d_f} = \left(\frac{\xi_s}{L} \right)^{3-\text{dim}} \quad (1)$$

This is true if the size ξ_s is the only length scale playing a role in the clusters and the network. However, in our case, there are two length scales ξ_s and ξ_c which brings

$$\phi = \left(\frac{r_0}{\xi_s} \right)^3 \left(\frac{\xi_c}{r_0} \right)^{d_f} \left(\frac{\xi_s}{L} \right)^{3-\text{dim}} \quad (2)$$

This has an impact on eqn (15) which becomes

$$G'_\infty = \underbrace{\frac{U}{r_0 \delta^2}}_{G_{CB}} \underbrace{\frac{1}{2} \left(\frac{\xi_c}{r_0} \right)^{d_f} \left(1 + \frac{\xi_s}{2\xi_c} \right)^{\frac{d_f}{3}} \left(1 - \frac{\xi_s}{\xi_c} \right)^{\frac{2d_f}{3}}}_{g_{\text{Interp}}} \underbrace{\phi \left(\frac{\xi_s}{r_0} \right)^2 \left(\frac{\xi_c}{r_0} \right)^{-d_f}}_{g_{\text{Net}}} \quad (3)$$

or in a compact form

$$G'_\infty = \frac{U}{r_0 \delta^2} \frac{1}{2} \left(1 + \frac{\xi_s}{2\xi_c} \right)^{\frac{d_f}{3}} \left(1 - \frac{\xi_s}{\xi_c} \right)^{\frac{2d_f}{3}} \phi \left(\frac{\xi_s}{r_0} \right)^2 \quad (4)$$

Fig. 4c in the published article displays the variations of the gel crossover elasticity G_c as a function of the shear rate intensity $\dot{\gamma}_0$ applied before flow cessation and the fit using the interpenetration ϕ -power law model. We note that $G'_\infty = 0.3G_c$. The fit of G_c in this figure needs to be modified accordingly to eqn (4) above and replaced by Fig. 1 shown here. In this Fig. 1, the corrected model fits well the data. The conclusion of the paper remains therefore unchanged.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Univ Lyon, Ens de Lyon, Univ Claude Bernard, CNRS, Laboratoire de Physique, F69342 Lyon, France. E-mail: thomas.gibaud@ens-lyon.fr

^b Groupe CFL, CEMEF, Mines Paristech, 1 Rue Claude Daunesse, 06904 Sophia Antipolis, France

^c ESRF – The European Synchrotron, 38043 Grenoble Cedex, France

^d CEA, DES, ISEC, DE2D, SEAD, LCBC, Université of Montpellier, Marcoule, France






Fig. 1 Evolution of G_c vs. $\dot{\gamma}_0$. The red line is the best fit of the data using eqn (4) here and the structural information reported in Fig. 6a of the published article. The best fit is obtained with a single adjustable parameter, namely the prefactor $G_{CB} = 9$ Pa.

